

16.04.2004



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

EP04/4076

| | |
|-------------------|-----|
| REC'D 28 JUN 2004 | |
| WIPO | PCT |

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03008753.0

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 03008753.0
Demande no:

Anmeldetag:
Date of filing: 17.04.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

NeuroNova AG
Kraepelinstrasse 2
80804 München
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se referer à la description.)

Means and methods for diagnosing and treating affective disorders

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C07K14/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Means and methods for diagnosing and treating affective disorders

The present invention relates to nucleic acid molecules, preferably genomic sequences, encoding an ATP-gated ion channel P2X7R which contain a mutation in the 5'UTR or 3'UTR regions, a mutation in exon 3, 5, 6, 8 or 13 or in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 or a deletion in exon 13, which allow to diagnose affective disorders. The invention further relates to polypeptides encoded by said nucleic acid molecules vectors and host cells comprising said nucleic acid molecules as well as to methods for producing polypeptides encoded by said nucleic acid molecules. The present invention also provides antibodies specifically directed to polypeptides encoded by said nucleic acid molecules and aptamers specifically binding said nucleic acid molecules.

Additionally, primers for selectively amplifying said nucleic acid molecules are provided in the present invention as well as kits, compositions, particularly pharmaceutical and diagnostic compositions comprising said nucleic acid molecules, vectors, polypeptides, aptamers, antibodies and/or primers. Moreover, the present invention relates to methods for diagnosing affective disorders associated with a non-functional P2X7R protein, an altered ATP-gating of the P2X7R protein, an over- or underexpression of the P2X7R protein or associated with the presence of any one of the aforementioned nucleic acid molecules or polypeptides encoded thereby. Additionally, the present invention relates to uses and methods for treating affective disorders employing a functional or non-functional ATP-gated ion-channel P2X7R.

The present invention also relates to uses of modulators of P2X7R activity for treating affective diseases.

Furthermore, the present invention also relates to methods for identifying and characterizing compounds which are capable of specifically interacting with or altering the characteristics of the polypeptides of the present invention as well as to methods for the production of pharmaceutical compositions.

Up to 10% of persons visiting a physician are afflicted with an affective disorder (also known as behavioural disorder, mood disorder). Nonetheless, most cases remain

undiagnosed or inadequately treated. Affective disorders include among others, depression, anxiety, and bipolar disorder. These diseases are well described in the literature; see, for example, Diagnostic and Statistical Manual of Mental Disorders-4th Edition Text Revision (DMS-IV-TR), American Psychiatric Press, 2000.

Depression, also known as unipolar affective disorder, is characterized by a combination of symptoms such as lowered mood, loss of energy, loss of interest, feeling of physical illness, poor concentration, altered appetite, altered sleep and a slowing down of physical and mental functions resulting in a relentless feeling of hopelessness, helplessness, guilt, and anxiety. The primary subtypes of this disease are major depression, dysthymia (milder depression), and atypical depression. Other important forms of depression are premenstrual dysphoric disorder and seasonal affective disorder. Present treatment of depression consists of psychotherapy, antidepressant drugs, or a combination of both. Most antidepressive drugs target the transport of the neurotransmitters serotonin and/or norepinephrine, or the activity of the enzyme monoamine oxidase. They include: Selective serotonin-reuptake inhibitors (e.g., fluoxetine, paroxetine, sertraline, fluvoxamine), tricyclic antidepressants (e.g., amitriptyline, imipramine, desipramine, nortriptyline), monoamine oxidase inhibitors (e.g., phenelzine, isocarboxazid, tranylcypromine), and designer antidepressants such as mirtazapine, reboxetine, nefazodone. However, all existing antidepressive drugs possess shortcomings such as long latency until response, high degree of non-responders and undesirable side effects (Holsboer, Biol. Psychol. 57 (2001), 47-65). Therefore, a need exists in the medical community for new antidepressive drugs with improved pharmacological profile (Baldwin, Hum. Psychopharmacol. Clin. Exp. 16 (2001), S93-S99).

Anxiety disorders are defined by an excessive or inappropriate aroused state characterized by feelings of apprehension, uncertainty, or fear. They are classified according to the severity and duration of their symptoms and specific affective characteristics. Categories include: (1) Generalized anxiety disorder, (2) panic disorder, (3) phobias, (4) obsessive-compulsive disorder, (5) post-traumatic stress disorder, and (6) separation anxiety disorder. The standard treatment for most anxiety disorders is a combination of cognitive-behavioural therapy with antidepressant medication. Additional medications include benzodiazepines and buspirone.

Bipolar disorder, also known as manic-depression, is characterized by mood swings between periods of mania (i.e. mood elevation including exaggerated euphoria, irritability) and periods of depression. Bipolar disorder is classified according to the severity of the symptoms. Patients diagnosed with bipolar disorder type I suffer from manic or mixed episodes with or without major depression. In Bipolar Disorder type II, patients have episodes of hypomania and episodes of major depression. With hypomania the symptoms of mania (euphoria or irritability) appear in milder forms and are of shorter duration. The current drugs used to treat bipolar disorders are lithium, valproate and lamotrigine, which stimulates the release of the neurotransmitter glutamate. As with antidepressive drugs, they take weeks to become effective and can result in undesirable side effects, for example, high levels of lithium in the blood can be fatal.

Compelling evidence suggest that affective disorders are biological diseases. However, there are no laboratory tests or other procedures that a common physician can use to make a definitive diagnosis. Instead, a specially trained physician or psychiatrist must diagnose the illness based on a group of symptoms that occur together. This process is often time consuming and laborious requiring several visits for the physician to perform a careful history of the symptoms that the patient is currently experiencing as well as any symptoms he or she has had in the past. Therefore, an easy and effective method for the accurate diagnosis of affective disorders is of high interest to the medical community (Wittchen et al., J. Clin. Psychiatry 62, suppl. 26 (2001), 23-28).

Most patients afflicted with affective disorders have family antecedents and identical twins studies suggest a strong genetic component. For example, genetic mapping on an isolated population of the central valley of Costa Rica suggests a locus for severe bipolar disorder at chromosome 18q22-q23 (Freimer et al., Nature Genetics 12 (1996), 436-441). Moreover, genetic studies performed on the Old Order Amish population suggest that genes on chromosomes 6, 13, and 15 may contribute to the susceptibility of bipolar affective disorder (Ginns et al., Nature Genetics 12 (1996), 431-435). Recently, a genome-wide search in a homogenous population found in the Saguenay/Lac-St-Jean region of Quebec suggests the presence of a major locus for bipolar disorder on chromosome 12q23-q24 (Morissette et al., Am. J. Med. Genet. (Neuropsychiatr. Genet.) 88 (1999), 567-587). Susceptibility loci on chromosomes 5

and 21 were also found in this study. Other groups report minimal evidence for linkage in the region of 12q23 (Kelsoe et al., Proc. Natl. Acad. Sci. USA 98 (2001), 585-590; Skar, Annu. Rev. Genomics Hum. Genet. 3 (2002), 371-413). Given the various loci mentioned in the above studies (e.g., links to chromosomes 5, 6, 12, 13, 15, 18, 21), a definite genetic link for affective diseases remains to be found.

Thus, although several genes have been assumed to be linked with affective disorders as mentioned hereinabove, however, no clear correlation has so far been shown. Since no well-suited medication nor diagnosis on a molecular level for affective disorders is available, there is a need for identifying a gene whose mutations cause the whole spectrum of affective disorders as well as for providing medicaments and methods for diagnosis and treatment of affective disorders.

Thus, the technical problem underlying the present invention is to provide means and methods for diagnosis and treating affective disorders.

The solution to said technical problem is achieved by providing the embodiments characterized in the claims.

Accordingly, the present invention relates to a nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:

- (a) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 5'UTR region corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;
- (b) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of the ATP-gated ion channel P2X7R, wherein in the exon as indicated in column "Exon" of the following Table A the amino acid residue as indicated in column "Amino acid residue" of Table A corresponding to the position as indicated in column "Position in wild-type" of Table A of the wild-type ATP-gated ion channel P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue

Table A

| Exon | Amino acid residue | Position in wild-type |
|-------------|---------------------------|------------------------------|
| exon 3 | R (Arg) | 117 |
| exon 5 | G (Gly) | 150 |
| exon 6 | E (Glu) | 186 |
| exon 6 | L (Leu) | 191 |
| exon 8 | R (Arg) | 270 |
| exon 13 | I (Ile) | 568 |
| exon 13 | R (Arg) | 578 |

- (c) a nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in exon 5 or 8 corresponding to position 32548 or position 37633 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide
- (d) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted;
- (e) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R, wherein in the intron as indicated in column "Intron" of the following Table B the nucleotide as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1 is replaced by another nucleotide

Table B

| Intron | REPLACED NUCLEOTIDE | Position in wild-type |
|---------------|----------------------------|------------------------------|
| intron 1 | G | 3166 |
| intron 1 | C | 24778 |
| intron 1 | C | 24830 |
| intron 3 | A | 26308 |
| intron 3 | G | 26422 |
| intron 4 | G | 32394 |
| intron 4 | T | 32434 |
| intron 5 | A | 32783 |
| intron 6 | G | 35641 |
| intron 6 | A | 35725 |
| intron 6 | T | 36001 |

| | | |
|-----------|---|-------|
| intron 7 | G | 36378 |
| intron 7 | T | 36387 |
| intron 7 | G | 36398 |
| intron 9 | C | 47214 |
| intron 11 | T | 47563 |
| intron 12 | C | 54307 |
| intron 12 | G | 54308 |

- (f) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 3'UTR region corresponding to position 55169, 55170, 55171, 55917 or 54925 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;
- (g) a nucleotide sequence comprising at least 20 or 21 nucleotides and comprising the mutations or deletions as defined in any one of (a) to (f);
- (h) a nucleic acid sequence comprising a nucleotide sequence as shown in any one of SEQ ID NOs: 13 to 51;
- (i) a nucleic acid sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NOs: 5 to 12;
- (j) a nucleotide sequence which hybridizes to a nucleotide sequence defined in any one of (a) to (g) or to the nucleotide sequence of (h) and having a mutation as defined in any one of (a) to (f); and
- (k) a nucleic acid sequence being degenerate as a result of the genetic code to the nucleic acid sequence as defined in (j).

It has surprisingly been found that mutations in the P2X7R gene which encodes the ATP-gated ion channel P2X7R can cause the whole spectrum of affective disorders. Six different mutations in the 5'UTR of the P2X7R gene, seven different mutations in exons 3, 5, 6, 8 and 13 of the P2X7R gene leading to an amino acid replacement of the corresponding amino acid in the wild-type sequence of P2X7R depicted in SEQ ID NO: 3 or 4 and two mutations in exons 5 and 8 of said gene, respectively, leading to a replacement of a nucleotide by another nucleotide, a deletion of nucleotides in exon 13 of said gene, 18 mutations in introns 1, 3, 4, 5, 6, 7, 9, 11 and 12 and 5 mutations in the 3'UTR of the P2X7R gene have been identified to co-segregate with the affection status in 41 unrelated families affected with affective disorders. The

term "affective disorder" when used in the context of the present invention means to include, but is not limited to, depression, anxiety, unipolar disorder, bipolar disorder type I, bipolar disorder type II, mania, attention deficit hyperactive disorder, substance abuse, and any other disorders affecting the normal behaviour, or mood of an individual.

Each mutation causes alterations that can explain affective disorders as shown in the Examples hereinbelow.

P2X7R is an ATP-gated ion channel belonging to the P2X ionotropic channel family. The gene was first isolated from rat brain (Surprenant et al., (1996), 272, 735-738; Genbank accession number NM_019256) and subsequently from a human monocyte library (Rassendren et al., J. Biol. Chem. 272 (1997), 5482-5486; Genbank accession numbers NM_002562, Y09561) by virtue of its sequence homology with the other members of the P2X family. It was later found that P2X7R corresponded to the unidentified P2Z receptor which mediates the permeabilising action of ATP on mast cells and macrophages (Dahlgvist and Diamant, Acta Physiol. Scand. 34 (1974), 368-384; Steinberg and Silverstein, J. Biol. Chem. 262 (1987), 3118-3122; Gordon, Biochem. J. 233 (1986), 309-319). The P2X7R has two hydrophobic membrane-spanning domains, an extracellular loop, and forms transmembrane ion channels. P2X7 receptors seem to function only in homooligomeric form and bear a pharmacological profile markedly different from other P2X homo- or heteromers (North and Surprenant, Annual Rev. Pharmacology Toxicology 40 (2000), 563-580). P2X7R requires levels of ATP in excess of 1 mM to achieve activation, whereas other P2X receptors activate at ATP concentrations of $\leq 100 \mu\text{M}$ (Steinberg et al., J. Biol. Chem. 262 (1987), 8884-8888; Greenberg et al., J. Biol. Chem. 263 (1988), 10337-10343) 32). While all P2X receptors demonstrate non-selective channel-like properties following ligation, the channels formed by the P2X7R can rapidly transform into pores that can allow the passage of molecules of up to 900 Dalton (Virginio et al., J. Physiol. 519 (1999), 335-346).

P2X7R is expressed in hematopoietic cells, mast cells and macrophages (Surprenant et al., Science 272 (1996), 3118-3122), where it is organized in tetrameric or hexameric form (Kim et al., J. Biol. Chem. 276 (2001), 23262-23267). P2X7R is inter alia involved in the regulation of the immune function and inflammatory response.

Activation of P2X7R by ATP in macrophages is associated with mitogenic stimulation of T cells (Baricordi et al., Blood 87 (1996), 682-690), the release of cytokines such as interleukin-1 β (Griffiths et al., J. Immunol. 154 (1995), 2821-2828), and formation of macrophage polykaryons (Falzone et al., J. Clin. Invest. 95 (1995), 1207-1216). Stimulation of the P2X7R with ATP can also result in cell death by triggering massive transmembrane ion fluxes (particularly influx of Ca²⁺ and Na⁺, and efflux of K⁺) and the formation of non-selective plasma membrane pores (Di Virgilio et al., Cell Death Differ. 5 (1998), 191-199).

In the brain, P2X7R was originally thought to be restricted to microglia (resident macrophage of the brain) and ependymal cells rather than neurons (Collo et al., Neuropharmacology 36 (1997), 1277-1283) suggesting a role of P2X7R in neurodegeneration. However, P2X7R has since been found in neurons of the rat retina (Brandle et al., Brain Research Molecular Brain Res. 62 (1998), 106-109), cochlear ganglion cells (Brandle et al., Neuroscience Letters 273 (1999), 105-108), and presynaptic terminals of neurons throughout the brainstem and spinal cord (Deuchards et al., J. Neurosci. 21 (2001), 7143-7152). Subsequent studies also suggest that P2X7R regulates the release of neurotransmitters such as glutamate and GABA in neurons of the hippocampus (Armstrong et al., J. Neuroscience 22 (2002), 5938-5945, Sperlágh et al., J. Neurochem. 81 (2002), 1196-1211). Organisation of P2X7R in glial cells and astrocytes of the brain appears monomeric (Kim et al., J. Biol. Chem. 276 (2001), 23262-23267).

Several agonists and antagonists of P2X7R have been identified. Brilliant Blue (Jiang et al., Mol. Pharmacol. 58 (2000), 82-88), the isoquinolines 1-[N,O-Bis(5-isoquinolinesulfonyl)-N-methyl-L-tyrosyl]-4-phenylpiperazine and N-[1-[N-methyl-p-(5-isoquinolinesulfonyl) benzyl]-2-(4-phenylpiperazine)ethyl]-5-isoquinolinesulfonamide (Humphreys et al., Mol. Pharmacol., 54 (1998), 22-32), adamantane derivatives (WO 99/29660, WO 99/29661, WO 00/61569, WO 01/42194, WO 01/44170, WO 01/44213), substituted phenyl compounds (WO 00/71529), piperidine and piperazine derivatives (WO 01/46200) are antagonists of P2X7R while Oxidized ATP (oATP) acts as an irreversible inhibitor of the receptor (Chen et al., J. Biol. Chem., 268 (1993), 8199-8203). Some of these antagonists are presently being evaluated for the treatment of inflammatory, immune, and cardiovascular diseases. BzATP (2- and 3-O-(4-benzoylbenzoyl)adenosine 5-triphosphate) acts as agonist of P2X7R (North

and Surprenant, Annu. Rev. Pharmacol. Toxicol. 40 (2000), 563-580). WO 99/55901 describes a method for identifying compounds that modulate the activity of a mammalian purinoreceptor selected from the group consisting of P2X2, P2X3, P2X4, P2X5, P2X6 and P2X7 and suggests a role of said purinoreceptors in therapy of behavioural disorders such as epilepsy, depression and aging-associated degenerative diseases.

Mutant mice lacking P2X7R are healthy, fertile and demonstrate no overt phenotype. However, in contrast to their wild-type counterparts, LPS-activated peritoneal macrophages from *P2X7R*^{-/-} animals fail to generate mature interleukin-1 β (IL-1 β) when challenged with ATP suggesting an inability of peritoneal macrophages to release IL-1 in response to ATP (Solle et al, J. Biol. Chem. 276 (2001), 125-132). A detailed behavioural study of the P2X7R^{-/-} mice was not performed. In humans, a Glu-496 to Ala polymorphism leads to the loss of P2X7 function (Gu et al., J. Biol. Chem. 276 (2001), 11135–11142) and is associated with B-cell chronic lymphocytic leukaemia (Thunberg, et al, The Lancet 360 (2002), 1935-1939). Additional polymorphs in the putative P2X7R promoter region, and coding region have been reported (Li et al., FEBS Lett. 531 (2002), 127-131; EP 1199372).

Despite the abundant literature concerning P2X7R, a role in affective disorders has never been suggested or alluded to in the prior art.

Before the present invention is described in detail, it is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described herein as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art.

Preferably, the terms used herein are defined as described in "A multilingual glossary of biotechnological terms: (IUPAC Recommendations)", Leuenberger, H.G.W, Nagel, B. and Kölbl, H. eds. (1995), Helvetica Chimica Acta, CH-4010 Basel, Switzerland).

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step.

Several documents are cited throughout the text of this specification. Each of the documents cited herein (including all patents, patent applications, scientific publications, manufacturer's specifications, instructions, etc.), whether supra or infra, are hereby incorporated by reference in their entirety. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and "the", include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to "a reagent" includes one or more of such different reagents, and reference to "the method" includes reference to equivalent steps and methods known to those of ordinary skill in the art that could be modified or substituted for the methods described herein.

In accordance with the present invention, the term "nucleic acid sequence" means the sequence of bases comprising purine- and pyrimidine bases which are comprised by nucleic acid molecules, whereby said bases represent the primary structure of a nucleic acid molecule. Nucleic acid sequences include DNA, cDNA, genomic DNA, RNA, synthetic forms and mixed polymers, both sense and antisense strands, or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those skilled in the art.

When used herein, the term "polypeptide" means a peptide, a protein, or a polypeptide which encompasses amino acid chains of a given length, wherein the amino acid residues are linked by covalent peptide bonds. However, peptidomimetics of such proteins/polypeptides wherein amino acid(s) and/or peptide bond(s) have been replaced by functional analogs are also encompassed by the invention as well as other than the 20 gene-encoded amino acids, such as selenocysteine. Peptides, oligopeptides and proteins may be termed polypeptides. The terms polypeptide and

protein are often used interchangeably herein. The term polypeptide also refers to, and does not exclude, modifications of the polypeptide, e.g., glycosylation, acetylation, phosphorylation and the like. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature.

The term "position" used in accordance with the present invention means the position of either an amino acid within an amino acid sequence depicted herein or the position of a nucleotide within a nucleic acid sequence depicted herein.

The term "ATP-gated ion channel P2X7R", in accordance with this invention, denotes a polypeptide which can be classified as a member of the P2X ionotropic receptor family. They are also known as purinergic receptors. P2X receptors are ligand-gated ion channels. The ligand for these receptors may be ATP and/or another natural nucleotide such as ADP, UTP and UDP, or a synthetic nucleotide such as 2-methylthioATP. The criteria for the classification are: (1) a sequence homology that is higher than 39% across the family or different species; (2) signal transduction mechanism involving ion conductance (Khakh et al., Pharmacol Rev. 253 (2001), 107-18). Accordingly, the term "ATP-gated ion channel P2X7R" is interchangeable with the terms "ionotropic receptor" or "purinergic receptor". Preferably, the term "ATP-gated ion channel P2X7R" denotes a polypeptide which can be classified as an ATP-gated ion channel P2X7R on the basis of one or more structural and/or functional characteristics, preferably those described above. Structural characteristics refer to certain structural features which allow to classify a polypeptide as being a P2X7R protein. One such feature is the amino acid sequence. In the context of the present invention a polypeptide is classified as an ATP-gated ion channel P2X7R if it shows a certain degree of sequence identity over its own length to the amino acid sequence of the human P2X7R protein depicted in SEQ ID NO: 3 or 4. This degree of sequence identity is at least 40%, more preferably at least 50%, even more preferably at least 60%, at least 70%, at least 80%, at least 90% or at least 95%. It is particularly preferred that the degree of sequence identity is at least 65%.

Moreover, structural characteristics of P2X7R proteins are two hydrophobic membrane-spanning domains, an extra cellular loop which could be analyzed by using the program TMPRED (Hofmann Biol. Chem. 347 (1993), 166) or TMHMM (Krogh J. Mol. Bio. 305 (2001), 567-580). Additionally, P2X7R may exist as a single polypeptide, as dimer, tetramer or the like.

Thus, in the context of the present invention a protein is preferably classified as a P2X7R protein if it displays at least one of the above-mentioned structural characteristics. Functional characteristics refer to properties related to the biological activity of the P2X7R protein. In particular, P2X7R is an ATP-gated ion channel which allows calcium and sodium ions to pass from extracellular solution to intracellular solution, and allows potassium ions to pass from intracellular to extracellular solution. Moreover, the ATP-gated ion channel P2X7R forms naturally a homooligomeric form. The characteristics of P2X7R receptor proteins can be determined as mentioned hereinbelow. The term "ATP-gated ion channels P2X7R" comprises functional and non-functional forms of the ATP-gated ion channels P2X7R. A functional ATP-gated ion channel P2X7R is understood to be a P2X7R protein which has at least one of the above-mentioned functional characteristics which can be measured by methods known in the art. A non-functional ATP-gated ion channel P2X7R is a protein which can be classified as a P2X7R protein due to structural characteristics as described above but which has lost at least one, preferably all, functional characteristics of a P2X7R protein as described above. Non-functionality of the P2X7R protein can, e.g., be determined by measuring whether calcium and sodium ions can flow into cells or whether potassium ions can exit from cells. Thus, it is possible to determine the occurrence of a mutation in the ATP-gated ion channel P2X7R by measuring either calcium and/or sodium influx or efflux of cells. Cells harbouring a mutation in the P2X7R gene show an altered ion influx and/or efflux in comparison to cells harbouring a wild-type P2X7R protein. Additionally, there are different methods that could be used to determine whether the P2X7R is functional or non-functional, for example, altered. One method consists of measuring the rate of ATP-induced incorporation of ethidium into cells, e.g. cells isolated from an individual. Ethidium is incorporated into the cells through P2X7R pores, when the pore formation is activated by ATP. Cells are then incubated with or

without ATP in the presence of ethidium, then they are analyzed by flow cytometry. Ethidium fluorescence is measured and compared in the presence or absence of ATP. If the P2X7R has lower activity, the ethidium fluorescence induced by ATP will be lower than in control cells. Such a method was used to verify P2X7R activity in isolated B-lymphocytes and T-lymphocytes from leukaemia patients (Wiley et al., *Lancet* 359 (2002), 1114-1119). Briefly, isolated cells are incubated in 1 ml of Hepes buffered potassium chloride at 37°C with continuous stirring. Ethidium is then added at a concentration of 25 mol/l, followed 40 seconds later by the addition of 10 µl of 100 mmol/l ATP stock. Cells are analyzed at 1,000 events/s by flow cytometry using a Coulter Elite flow cytometer (Coulter, Hialeah, FL) with argon laser excitation at 488 nm. Fluorescent emission was collected using a 590-nm long-pass filter. The linear mean channel fluorescence intensity for each gated subpopulation over successive 5-s intervals was analyzed with the use of Win-MDI software (Joseph Trotter, version 2.7) and plotted against time.

Another method of determining P2X7R activity is to measure calcium entry into isolated cells incubated with fluorescent dyes that emit only upon binding to calcium. The cells have to be loaded with the dye and then the calcium entry has to be stimulated. Examples of such dyes include Fura-2, Calcium green, calcium orange, calcium crimson (all available from Molecular Probes). Methods of measuring calcium transport are well known in the art; see for example, Takahashi et al., *Physiol Rev.* 79 (1999), 1089-1125. Furthermore, calcium entry into the cells produces changes in the membrane electric potential. This changes can be measured by electrophysiology (patch clamp) or by using dyes which are sensitive to voltage change. Such methods are also well known in the art, see for example, Gonzalez et al., *DDT* 4 (1999), 431-439; González and Tsien, *Chemistry & Biology* 4 (1997), 269-277; González and Tsien, *Biophysical Journal* 69 (1995), 1272-1280.

Yet another method is to measure uptake of ¹³³Ba21. Ba21 is a good surrogate for Ca21 and once inside the cell is neither pumped nor sequestered by transport ATPases. Ba21 uptake can be measured over 60 s using ¹³³BaCl2 (final concentration, 0.2 mM). At time 0, a prewarmed stock solution of ¹³³Ba21 (0.4 mM and 1 µCi/ml) is added in equal volumes to prewarmed isolated cells in 150 mM KCl with HEPES (pH 7.4) at 37°C. ATP (1 mM) is added either 10 minutes before or simultaneously with the ¹³³Ba21 isotope. Aliquots of 0.8 ml are taken at time points

between 0 and 60 s and are immediately mixed with 0.2 ml of ice-cold 50 mM MgCl₂ (in KCl-HEPES medium) that had been previously layered over 250 µl of oil mixture (di-n-butyl phthalate and di-iso-octyl phthalate, 7:3 vol/vol) and then centrifuged at 8,000 g for 30 s. The supernatants and the oil are aspirated, and the cell pellets are counted in a Wallac Wizard 3 automatic gamma-counter or in any other suitable gamma measuring unit.

The present invention is based on the finding that mutations of different kinds in the P2X7R gene are linked to the occurrence of affective disorders. The first type of mutations are mutations in the 5'UTR. Examples of such mutations are single nucleotide replacements at positions corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as described in SEQ ID NO: 1.

The position with respect to nucleotide sequences mentioned herein refer to the sequence shown in SEQ ID NO: 1. This sequence represents the nucleic acid sequence of the P2X7R gene encoding the ATP-gated ion channel P2X7R. It is possible for the skilled person to identify the position in the genomic sequence corresponding to a position in SEQ ID NO: 1 by aligning the sequences. Moreover, the exact locations of the exons and introns are indicated in SEQ ID NO: 1 hereinbelow. Additionally, the person skilled in the art is able to identify exons and introns of the P2X7R gene by comparing SEQ ID NO: 1 with SEQ ID NO: 2 which shows the cDNA sequence of the P2X7R gene.

Preferably, at position 362 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a thymine (T) is replaced by another nucleotide, preferably a purine base. More preferably, at said position said thymidine is replaced by a pyrimidine base. Particularly preferred, said thymine is replaced by a cytosine (C).

At position 532 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a thymine (T) is preferably replaced by another nucleotide, preferably a pyrimidine base. More preferably, at said position said thymine is replaced by a purine base. Particularly preferred, said thymidine is replaced by a guanine (G).

The adenine (A) residues at positions 1100 and 1122, respectively, in the 5'UTR of the genomic sequence of the P2X7R gene is preferably replaced by a pyrimidine base. More preferably, said adenine is replaced by a purine base and particularly preferred said adenine is replaced by a guanine (G).

At position 1171 in the 5'UTR of the genomic sequence of the P2X7R gene depicted in SEQ ID NO: 1 a cytidine (C) is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, said cytidine is replaced by a purine base and even more preferred, said cytidine is replaced by a guanine (G).

The guanine at position 1702 in the 5'UTR of the genomic sequence of the gene P2X7R depicted in SEQ ID NO: 1 is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, said guanine is replaced by a purine base and particularly preferred it is replaced by an adenine (A).

A second type of mutation found in the P2X7R gene are mutations in exons which lead to amino acid substitutions in the corresponding amino acid sequence. These are the mutations listed under item (b), supra. In this context, the term "an amino acid residue as indicated in column 'Amino acid residue' of Table A corresponding to position X of the wild-type ATP-gated ion channel P2X7R as depicted in column 'Position in wild-type' " has the following meaning: The amino acid residue in question would be located at position X in the sequence of SEQ ID NO: 3 or 4 if the sequence in which said amino acid residue occurs is compared and aligned with the amino acid sequence of SEQ ID NO: 3 or 4. The amino acid sequence shown in SEQ ID NO: 3 or 4 is the amino acid sequence of the human P2X7R gene and is used as a reference sequence in the present invention.

In order to determine whether an amino acid residue or nucleotide residue in a given P2X7R sequence corresponds to a certain position in the amino acid sequence or nucleotide sequence of SEQ ID NO: 1, 3 or 4, the skilled person can use means and methods well-known in the art, e.g., alignments, either manually or by using computer programs such as those mentioned further down below in connection with the definition of the term "hybridization" and degrees of homology.

For example, BLAST2.0, which stands for Basic Local Alignment Search Tool (Altschul, Nucl. Acids Res. 25 (1997), 3389-3402; Altschul, J. Mol. Evol. 36 (1993), 290-300; Altschul, J. Mol. Biol. 215 (1990), 403-410), can be used to search for local sequence alignments. BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying similar sequences. The fundamental unit of BLAST algorithm output is the High-scoring Segment Pair (HSP). An HSP consists of two sequence fragments of arbitrary but equal lengths whose alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cutoff score set by the user. The BLAST approach is to look for HSPs between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence whose match satisfies E is reported in the program output.

Analogous computer techniques using BLAST (Altschul (1997), loc. cit.; Altschul (1993), loc. cit.; Altschul (1990), loc. cit.) are used to search for identical or related molecules in nucleotide databases such as GenBank or EMBL. This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score which is defined as:

$$\frac{\% \text{sequence identity} \times \% \text{maximum BLAST score}}{100}$$

100

and it takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error, and at 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

As mentioned above, the second group of mutations identified in the P2X7R gene are mutations in the exons of the P2X7R gene which lead to amino acid substitutions. In this respect SEQ ID NO 2 shows the cDNA sequence of the P2X7R gene. In exon 3 at position 117 of the corresponding wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 an arginine (R) residue is replaced by another amino acid residue, preferably by an aliphatic, acidic or basic amino acid residue. More preferably, by an aromatic amino acid residue which is particularly preferred to be a tryptophane (W). The resulting polypeptide is shown in SEQ ID NO: 5.

In exon 5 at position 150 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 a glycine (G) residue is replaced by another amino acid residue, preferably by an aliphatic, aromatic or acidic amino acid residue. More preferably, by a basic amino acid residue and particularly preferred by an arginine (R). The resulting polypeptide is shown in SEQ ID NO: 6.

At position 186 in exon 6 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 a glutamate residue (E) is replaced by another amino acid residue, preferably by an aliphatic, aromatic or acidic amino acid residue. More preferably, said glutamate is replaced by a basic amino acid residue which is particularly preferred a lysine (K). The resulting polypeptide is shown in SEQ ID NO: 7.

In exon 6 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 at position 191 a leucine residue (L) is replaced by another amino acid residue. Said amino acid residue is preferably an aliphatic, acidic or basic amino acid residue. More preferably, said amino acid residue is an aromatic amino acid residue which is particularly preferred to be a proline (P). The resulting polypeptide is shown in SEQ ID NO: 8.

In exon 8 of the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 at position 270 an arginine residue (R) is replaced by another amino acid residue. Said amino acid residue is preferably an aromatic, acidic or basic amino acid residue. More preferably, said amino acid residue is an aliphatic amino acid residue

which is particularly preferred to be a cysteine (C). The resulting polypeptide is shown in SEQ ID NO: 9.

At position 568 in exon 13 of the wild-type amino acid residue of P2X7R depicted in SEQ ID NO: 3 or 4 an isoleucine (I) residue is replaced by another amino acid residue. More preferably, said isoleucine is replaced by an aromatic, basic or acidic amino acid residue. Even more preferred, said isoleucine is replaced by an aliphatic amino acid residue which is particularly preferred to be an asparagine (N). The resulting polypeptide is shown in SEQ ID NO: 10.

In exon 13 at position 578 in the wild-type amino acid sequence of P2X7R depicted in SEQ ID NO: 3 or 4 an arginine residue (R) is replaced by another amino acid residue. Said amino acid residue is preferably an aromatic, acidic or basic amino acid residue. More preferably, it is an aliphatic amino acid residue and particularly preferred it is a glutamine (Q) residue. The resulting polypeptide is shown in SEQ ID NO: 12.

It is envisaged that the above-mentioned mutations in the exons of the P2X7R gene occur due to point mutations caused by, e.g. chemical and/or physical means or inaccuracy of the replication complex followed by a failure of the reparation machinery of a cell, a change of a single codon occur can be achieved. Possible types of point mutations are transitions, i.e. change of a purine or pyrimidine base for another purine or pyrimidine base, e.g. adenine to guanine or thymidine to cytosine or transversions, i.e. change of a purine or pyrimidine base for another pyrimidine or purine base, e.g., adenine to thymidine or guanine to cytosine. Additionally a point mutation can also be caused by insertion or deletion of one or more nucleotides.

The mutations leading to the replacement of the amino acids as mentioned hereinabove and hereinbelow are indicated in Table 1 hereinbelow.

The third group of mutations in the P2X7R gene has been identified to be in exons 5 and 8 of the P2X7R gene depicted in SEQ ID NO: 1 and to be silent, i.e. they do not lead to amino acid changes. In particular, at position 32548 in exon 5 of the wild-type

genomic sequence P2X7R gene depicted in SEQ ID NO: 1 a cytidine residue is replaced by another nucleotide. Said nucleotide is preferably a pyrimidine base and particularly preferred a thymine. The exchange of the cytidine residue at position 32548 in exon 5 of the P2X7R gene by another nucleotide preferably does not lead to the replacement of the amino acid cysteine by another amino acid residue.

In exon 8 of the wild-type P2X7R gene depicted in SEQ ID NO: 1 at position 37633 a cytidine residue is replaced by another nucleotide residue. Said nucleotide residue is preferably a pyrimidine base and particularly preferred thymine. Due to this replacement the amino acid aspartate (D) encoded by the respective codon in which at position 37633 a replacement has taken place is preferably not replaced by another amino acid residue.

The above-mentioned mutations in exons 5 and 8 at positions 32548 and 37633, respectively, of the wild-type P2X7R gene depicted in SEQ ID NO:1 are mutations at the third position of a triplet codon, i.e. at the wobble base, which lead to so-called silent mutations. Silent mutations do normally not lead to a change of the amino acid due to the degeneracy of the genetic code, i.e. 64 triplets encode at all 20 naturally occurring amino acids. However, said silent mutations lead to a change in the codon encoding its respective amino acids insofar that the newly generated codon may not fit so well into the codon usage of an organism. Namely, the newly generated codon is not translated by the ribosome with the same efficiency as the "old" codon. This may lead to insufficient amounts of the corresponding polypeptide causing an distinct phenotype.

The fourth group of mutations in the P2X7R gene described hereinabove in item (d) is a deletion of 7 amino acids corresponding to positions 488 to 494 of the wild-type P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4. Thus, the present invention also relates to nucleic acid sequences encoding a P2X7R protein in which amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted. This means, according to the present invention, that a fragment encompassing amino acid positions 488 to 494 of the corresponding wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4 is deleted which results in a shortened polypeptide. An example

for such a shortened polypeptide is depicted in SEQ ID NO: 11. This type of mutation as described herein preferably encodes a non-functional ATP-gated ion channel P2X7R. In the present invention the deletion of a fragment encompassing amino acids 488 to 494 of the wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4 is the result of a deletion in exon 13. The resulting protein depicted in SEQ ID NO: 11 lacks amino acids 488 to 494 of the corresponding wild type amino acid sequence depicted in SEQ ID NO: 3 or 4 such that amino acid position 494 of the deleted polypeptide depicted in SEQ ID NO: 11 corresponds to amino acid position 502 of the wild-type amino acid sequence depicted in SEQ ID NO: 3 or 4. Preferably, the nucleic acid sequence of the invention encodes a P2X7R polypeptide in which exactly amino acids corresponding to positions 488 to 494 of SEQ ID NO: 3 or 4 are deleted. However, also mutants are comprised in which either more or less amino acids within the P2X7R amino acid sequence set forth in SEQ ID NO: 3 or 4 may be deleted due to, for example, atypical splicing or deletion of nucleotides of the nucleic acid molecule encoding P2X7R or wrong posttranslational processes, as long as the P2X7R ATP-gated ion channel is non-functional. For example, it is also possible that further amino acids preceding amino acid position 488 or amino acids succeeding amino acid position 494 may be deleted or that less amino acids are deleted.

Preferably at least one, more preferably at least two, even more preferably at least three and most preferably at least 5 amino acid residues are further deleted upstream from the position corresponding to amino acid residue 488 and/or downstream of the position corresponding to amino acid residue 494 of SEQ ID NO: 3 or 4.

However, it is preferred that not more than 20, preferably not more than 15, even more preferably not more than 10 and most preferably not more than 7 amino acid residues are further deleted upstream of the position corresponding to amino acid residue 488 of SEQ ID NO: 3 or 4 or downstream of the position corresponding to amino acid residue 494 of SEQ ID NO: 3 or 4.

Another group of mutation (mentioned in item (e), supra) resides in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 of the wild-type genomic sequence of P2X7R depicted in SEQ ID NO: 1. Said mutations in said introns are point mutations as shown in Table B hereinabove and in Table 1, hereinbelow.

At the respective position indicated in the column "Position in wild-type" in Table B or indicated in the column "Polymorphism" in Table 1 the position of the nucleotide residue in the respective intron which is replaced by another nucleotide residue is shown. Accordingly, the term "a nucleotide as indicated in column "Intron" of the Table B corresponding to the position as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B is replaced by another nucleotide means that a nucleotide residue in a P2X7R encoding sequence would be located at position Y in SEQ ID NO: 1 when the P2X7R sequence is compared and aligned with the sequence of SEQ ID NO: 1.

If the nucleotide at the respective position is a purine base such as adenine or guanine it is preferred that due to a transition it is replaced by another purine base. For example, an adenine is replaced by a guanine or a guanine is replaced by an adenine. If the nucleotide at the respective position is a pyrimidine base it is preferred that due to a transition it is replaced by another pyrimidine base. For example, thymine is replaced by a cytidine and a cytidine is replaced by a thymine.

It is also preferred that due to a transversion a purine base is replaced by a pyrimidine base or vice versa. For example, an adenine is replaced by a thymine and a guanine is replaced by a cytidine. Particularly preferred, said nucleotide in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 of the P2X7R gene depicted in SEQ ID NO: 1 is replaced by the nucleotide depicted in column "Polymorphism" of Table 1, hereinbelow.

A last group of mutations that has been identified relates to mutations which reside in the 3'UTR of the wild-type P2X7R gene depicted in SEQ ID NO: 1. The mutations were found at positions 54925, 55169, 55170, 55171 or 55917 respectively, of the wild-type P2X7R gene depicted in SEQ ID NO: 1.

At position 54925 a guanine residue was found to be replaced by another nucleotide. Preferably, said guanine residue is replaced by a pyrimidine base, more preferably by a purine base and particularly preferred by an adenine.

At position 55169 a cytidine residue is replaced by another nucleotide, preferably by a pyrimidine base. More preferably, it is replaced by a purine base and particularly preferred, it is replaced by an adenine.

At positions 55170 and 55171 an adenine residue is replaced by another nucleotide residue, preferably by a purine base. More preferably, said adenine residue is replaced by a pyrimidine base and particularly preferred said adenine residue is replaced by a cytidine residue. It was also found that at position 55917 a cytidine residue is replaced by another nucleotide. Preferably, said nucleotide residue is a purine base, more preferably a pyrimidine base and particularly preferable a thymine.

As is evident from the above, not all identified mutations are located in exons or lead to a change in the amino acid sequence. Some of the mutations are located in the 5'UTR, the 3'UTR or in introns.

It is known that polymorphisms in promoter and enhancer regions can affect gene function by modulating transcription, particularly if they are situated at recognition sites for DNA binding proteins (Fishman et al., J. Clin. Invest. 102 (1998), 1369-1376). The term "polymorphism" which is used in the present invention means single nucleotide substitution, nucleotide insertion and nucleotide deletion which in the case of insertion and deletion includes insertion or deletion of one or more nucleotides at a position of a gene and corresponding alterations in expressed proteins. Polymorphisms in the 5' untranslated region (5'UTR) of genes can affect the efficiency with which proteins are translated. A representative example of this is in the c-myc gene where a C-G SNP that creates an internal ribosome entry site is associated with increased efficiency of c-myc translation and myeloma (Chappell et al., Oncogene 19 (2000), 4437-4440). Polymorphisms in the 3'UTR can affect gene function by altering the secondary structure of RNA and efficiency of translation or by affecting motifs in the RNA that bind proteins which regulate RNA degradation. Polymorphisms within introns can affect gene function by affecting RNA splicing resulting in aberrant polypeptides. Another way in which intronic polymorphisms can affect gene function is when they affect regulatory motifs within introns. Examples are the Sp1 binding site polymorphism within intron 1 of the COLIA1 gene (Mann et al., J. Clin. Invest 107 (2001), 899-907) and a repeat polymorphisms within the IL-1Ra

gene (Keen et al., Bone 23 (1998), 367-371). Further examples between intronic SNPs and gene function are described in Caceres and Komblitt, Trends Genet. 4 (2002), 186-93. Example 4 on page 52, line 30 to page 53, line 51 of the text describes potential alternative splicing events and aberrant protein production associated with three SNPs disclosed in the application.

The nucleic acid sequences described hereinabove may comprise at least 56580 nucleotides, preferably at least 10000 nucleotides, at least 5000 nucleotides, at least 1000 nucleotides, at least 500 nucleotides, at least 100 nucleotides. More preferably, said nucleic acid sequences comprise at least 50 nucleotides and particularly preferred they comprise at least 20 or 21 nucleotides comprising the mutations or deletions as described hereinabove. Most preferably such a nucleic acid sequence has a sequence as depicted in any one of SEQ ID NOs: 13 to 51.

The nucleic acid sequences described hereinabove which comprise mutations in exons leading to a replacement of the corresponding amino acid sequence of the P2X7R wild-type polypeptide depicted in SEQ ID NO: 3 or 4 encode polypeptides shown in SEQ ID NOs: 5 to 10 and 12

Additionally, the nucleic acid sequences described hereinabove which comprise a deletion leading to a truncated polypeptide in comparison to the full-length polypeptide of the wild-type P2X7R polypeptide shown in SEQ ID NO: 3 or 4 is shown in SEQ ID NO: 11.

The present invention also relates to nucleic acid molecules which hybridize to one of the above described nucleic acid molecules and which shows a mutation as described hereinabove.

The term "hybridizes" as used in accordance with the present invention may relate to hybridizations under stringent or non-stringent conditions. If not further specified, the conditions are preferably non-stringent. Said hybridization conditions may be established according to conventional protocols described, for example, in Sambrook, Russell "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory, N.Y. (2001); Ausubel, "Current Protocols in Molecular Biology", Green Publishing Associates and Wiley Interscience, N.Y. (1989), or Higgins and Hames

(Eds.) "Nucleic acid hybridization, a practical approach" IRL Press Oxford, Washington DC, (1985). The setting of conditions is well within the skill of the artisan and can be determined according to protocols described in the art. Thus, the detection of only specifically hybridizing sequences will usually require stringent hybridization and washing conditions such as 0.1xSSC, 0.1% SDS at 65°C. Non-stringent hybridization conditions for the detection of homologous or not exactly complementary sequences may be set at 6xSSC, 1% SDS at 65°C. As is well known, the length of the probe and the composition of the nucleic acid to be determined constitute further parameters of the hybridization conditions. Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility. Hybridizing nucleic acid molecules also comprise fragments of the above described molecules. Such fragments may represent nucleic acid sequences which code for a non-functional ATP-gated ion channel P2X7R or a non-functional fragment thereof, and which have a length of at least 12 nucleotides, preferably at least 15, more preferably at least 18, more preferably of at least 21 nucleotides, more preferably at least 30 nucleotides, even more preferably at least 40 nucleotides and most preferably at least 60 nucleotides. Furthermore, nucleic acid molecules which hybridize with any of the aforementioned nucleic acid molecules also include complementary fragments, derivatives and allelic variants of these molecules. Additionally, a hybridization complex refers to a complex between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., Cot or Rot analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips, pins or glass slides to which, e.g., cells have been fixed). The terms complementary or

complementarity refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, the sequence "A-G-T" binds to the complementary sequence "T-C-A". Complementarity between two single-stranded molecules may be "partial", in which only some of the nucleic acids bind, or it may be complete when total complementarity exists between single-stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

The term "hybridizing sequences" preferably refers to sequences which display a sequence-identity of at least 40%, preferably at least 50%, more preferably at least 60%, even more preferably at least 70%, particularly preferred at least 80%, more particularly preferred at least 90%, even more particularly preferred at least 95% and most preferably at least 97% identity with a nucleic acid sequence as described above encoding a P2X7R protein having a described mutation. Moreover, the term "hybridizing sequences" preferably refers to sequences encoding a P2X7R protein having a sequence identity of at least 40%, preferably at least 50%, more preferably at least 60%, even more preferably at least 70%, particularly preferred at least 80%, more particularly preferred at least 90%, even more particularly preferred at least 95% and most preferably at least 97% identity with an amino acid sequence of a P2X7R mutant as described herein above.

In accordance with the present invention, the term "identical" or "percent identity" in the context of two or more nucleic acid or amino acid sequences, refers to two or more sequences or subsequences that are the same, or that have a specified percentage of amino acid residues or nucleotides that are the same (e.g., 60% or 65% identity, preferably, 70-95% identity, more preferably at least 95% identity), when compared and aligned for maximum correspondence over a window of comparison, or over a designated region as measured using a sequence comparison algorithm as known in the art, or by manual alignment and visual inspection. Sequences having, for example, 60% to 95% or greater sequence identity are considered to be substantially identical. Such a definition also applies to the complement of a test sequence. Preferably the described identity exists over a region that is at least about 15 to 25 amino acids or nucleotides in length, more preferably,

over a region that is about 50 to 100 amino acids or nucleotides in length. Those having skill in the art will know how to determine percent identity between/among sequences using, for example, algorithms such as those based on CLUSTALW computer program (Thompson Nucl. Acids Res. 2 (1994), 4673-4680) or FASTDB (Brutlag Comp. App. Biosci. 6 (1990), 237-245), as known in the art.

Although the FASTDB algorithm typically does not consider internal non-matching deletions or additions in sequences, i.e., gaps, in its calculation, this can be corrected manually to avoid an overestimation of the % identity. CLUSTALW, however, does take sequence gaps into account in its identity calculations. Also available to those having skill in this art are the BLAST and BLAST 2.0 algorithms (Altschul Nucl. Acids Res. 25 (1977), 3389-3402). The BLASTN program for nucleic acid sequences uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (W) of 3, and an expectation (E) of 10. The BLOSUM62 scoring matrix (Henikoff Proc. Natl. Acad. Sci., USA, 89, (1989), 10915) uses alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

Moreover, the present invention also relates to nucleic acid molecules the sequence of which is degenerate in comparison with the sequence of an above-described hybridizing molecule. When used in accordance with the present invention the term "being degenerate as a result of the genetic code" means that due to the redundancy of the genetic code different nucleotide sequences code for the same amino acid.

The present invention also related to nucleic acid molecules which comprise one or more of the above-described mutations or deletions.

The nucleic acid molecules according to the invention may be derived from any organism encoding corresponding P2X7R ATP-gated ion channels. For example, P2X7R ATP-gated ion channels have been reported in various organisms, for example, rat (see, Suprenant (1996), loc. cit.), mouse (Genbank Accession No. AJ 489297), xenopus (Genbank Accession No. AJ 345114), chicken (Genbank Accession No. BM 491404) or Bos Taurus (Genbank Accession No. AF 083073) . In

a preferred embodiment the nucleic acid molecule of the invention is derived from a vertebrate, preferably from a mammal, even more preferably the nucleic acid molecule is derived from rabbit or guinea pig, and most preferably the nucleic acid is derived from mouse, rat or human.

The nucleic acid molecule according to the invention may be any type of nucleic acid, e.g. DNA, RNA or PNA (peptide nucleic acid).

For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by Nielsen et al., *Science* 254:1497 (1991); and Egholm et al., *Nature* 365:666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ($T_{sub.m}$) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

The DNA may, for example, be cDNA. In a preferred embodiment it is a genomic DNA. The RNA may be, e.g., mRNA. The nucleic acid molecule may be natural, synthetic or semisynthetic or it may be a derivative, such as peptide nucleic acid (Nielsen, *Science* 254 (1991), 1497-1500) or phosphorothioates. Furthermore, the nucleic acid molecule may be a recombinantly produced chimeric nucleic acid molecule comprising any of the aforementioned nucleic acid molecules either alone or in combination.

Preferably, the nucleic acid molecule of the present invention is part of a vector. Therefore, the present invention relates in another embodiment to a vector comprising the nucleic acid molecule of this invention. Such a vector may be, e.g., a plasmid, cosmid, virus, bacteriophage or another vector used e.g. conventionally in genetic engineering, and may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions.

The nucleic acid molecules of the present invention may be inserted into several commercially available vectors. Nonlimiting examples include plasmid vectors compatible with mammalian cells, such as pUC, pBluescript (Stratagene), pET (Novagen), pREP (Invitrogen), pCRTopo (Invitrogen), pcDNA3 (Invitrogen), pCEP4 (Invitrogen), pMC1 neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), EBO-pSV2neo, pBPV-1, pdBPVMMTneo, pRSVgpt, pRSVneo, pSV2-dhfr, pUCTag, pZD35, pLXIN and pSIR (Clontech) and pIRES-EGFP (Clontech). Baculovirus vectors such as pBlueBac, BacPacz Baculovirus Expression System (CLONTECH), and MaxBacTM Baculovirus Expression System, insect cells and protocols (Invitrogen) are available commercially and may also be used to produce high yields of biologically active protein. (see also, Miller (1993), Curr. Op. Genet. Dev., 3, 9; O'Reilly, Baculovirus Expression Vectors: A Laboratory Manual, p. 127). In addition, prokaryotic vectors such as pcDNA2; and yeast vectors such as pYes2 are nonlimiting examples of other vectors suitable for use with the present invention. For vector modification techniques, see Sambrook and Russel (2001), loc. cit. Vectors can contain one or more replication and inheritance systems for cloning or expression, one or more markers for selection in the host, e. g., antibiotic resistance, and one or more expression cassettes.

The coding sequences inserted in the vector can be synthesized by standard methods, isolated from natural sources, or prepared as hybrids. Ligation of the coding sequences to transcriptional regulatory elements (e. g., promoters, enhancers, and/or insulators) and/or to other amino acid encoding sequences can be carried out using established methods.

Furthermore, the vectors may, in addition to the nucleic acid sequences of the invention, comprise expression control elements, allowing proper expression of the coding regions in suitable hosts. Such control elements are known to the artisan and may include a promoter, translation initiation codon, translation and insertion site or internal ribosomal entry sites (IRES) (Owens, Proc. Natl. Acad. Sci. USA 98 (2001), 1471-1476) for introducing an insert into the vector. Preferably, the nucleic acid molecule of the invention is operatively linked to said expression control sequences allowing expression in eukaryotic or prokaryotic cells. Particularly preferred are in this context control sequences which allow for correct expression in neuronal cells and/or cells derived from nervous tissue.

Control elements ensuring expression in eukaryotic and prokaryotic cells are well known to those skilled in the art. As mentioned above, they usually comprise regulatory sequences ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in for example mammalian host cells comprise the CMV-HSV thymidine kinase promoter, SV40, RSV-promoter (Rous sarcome virus), human elongation factor 1 α -promoter, CMV enhancer, CaM-kinase promoter or SV40-enhancer.

For the expression for example in nervous tissue and/or cells derived therefrom, several regulatory sequences are well known in the art, like the minimal promoter sequence of human neurofilament L (Charron, J. Biol. Chem. 270 (1995), 25739-25745). For the expression in prokaryotic cells, a multitude of promoters including, for example, the tac-lac-promoter, the lacUV5 or the trp promoter, has been described. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pRc/CMV, pcDNA1, pcDNA3 (Invitrogen, as used, inter alia in the appended examples), pSPORT1 (GIBCO BRL) or pGEMHE (Promega), or prokaryotic expression vectors, such as lambda gt11.

An expression vector according to this invention is at least capable of directing the replication, and preferably the expression, of the nucleic acids and protein of this invention. Suitable origins of replication include, for example, the Col E1, the SV40 viral and the M 13 origins of replication. Suitable promoters include, for example, the cytomegalovirus (CMV) promoter, the iacZ promoter, the gai10 promoter and the Autographa californica multiple nuclear polyhedrosis virus (AcMNPV) polyhedral promoter. Suitable termination sequences include, for example, the bovine growth hormone, SV40, iacZ and AcMNPV polyhedral polyadenylation signals. Examples of selectable markers include neomycin, ampicillin, and hygromycin resistance and the like. Specifically-designed vectors allow the shuttling of DNA between different host cells, such as bacteria-yeast, or bacteria-animal cells, or bacteria-fungal cells, or bacteria invertebrate cells.

Beside the nucleic acid molecules of the present invention, the vector may further comprise nucleic acid sequences encoding for secretion signals. Such sequences are well known to the person skilled in the art. Furthermore, depending on the expression system used leader sequences capable of directing the expressed polypeptide to a cellular compartment may be added to the coding sequence of the nucleic acid molecules of the invention and are well known in the art. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a part thereof, into, inter alia, the extracellular membrane. Optionally, the heterologous sequence can encode a fusion protein including an C- or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. Once the vector has been incorporated into the appropriate host, the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and, as desired, the collection and purification of the proteins, antigenic fragments or fusion proteins of the invention may follow. Of course, the vector can also comprise regulatory regions from pathogenic organisms.

Furthermore, said vector may also be, besides an expression vector, a gene transfer and/or gene targeting vector. Gene therapy, which is based on introducing therapeutic genes (for example for vaccination) into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable

vectors, vector systems and methods for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, Nature Medicine 2 (1996), 534-539; Schaper, Circ. Res. 79 (1996), 911-919; Anderson, Science 256 (1992), 808-813, Isner, Lancet 348 (1996), 370-374; Muhlhauser, Circ. Res. 77 (1995), 1077-1086; Wang, Nature Medicine 2 (1996), 714-716; WO 94/29469; WO 97/00957; Schaper, Current Opinion in Biotechnology 7 (1996), 635-640 or Verma, Nature 389 (1997), 239-242 and references cited therein. The nucleic acid molecules of the invention and vectors as described herein above may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Additionally, baculoviral systems or systems based on vaccinia virus or Semliki Forest Virus can be used as eukaryotic expression system for the nucleic acid molecules of the invention. In addition to recombinant production, fragments of the protein, the fusion protein or antigenic fragments of the invention may be produced by direct peptide synthesis using solid-phase techniques (cf Stewart et al. (1969) Solid Phase Peptide Synthesis; Freeman Co, San Francisco; Merrifield, J. Am. Chem. Soc. 85 (1963), 2149-2154). In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City CA) in accordance with the instructions provided by the manufacturer. Various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

The present invention in addition relates to a host transformed with a vector of the present invention or to a host comprising the nucleic acid molecule of the invention. Said host may be produced by introducing said vector or nucleotide sequence into a host cell which upon its presence in the cell mediates the expression of a protein encoded by the nucleotide sequence of the invention or comprising a nucleotide sequence or a vector according to the invention wherein the nucleotide sequence and/or the encoded polypeptide is foreign to the host cell.

By "foreign" it is meant that the nucleotide sequence and/or the encoded polypeptide is either heterologous with respect to the host, this means derived from a cell or organism with a different genomic background, or is homologous with respect to the

host but located in a different genomic environment than the naturally occurring counterpart of said nucleotide sequence. This means that, if the nucleotide sequence is homologous with respect to the host, it is not located in its natural location in the genome of said host, in particular it is surrounded by different genes. In this case the nucleotide sequence may be either under the control of its own promoter or under the control of a heterologous promoter. The location of the introduced nucleic acid molecule or the vector can be determined by the skilled person by using methods well-known to the person skilled in the art, e.g., Southern Blotting. The vector or nucleotide sequence according to the invention which is present in the host may either be integrated into the genome of the host or it may be maintained in some form extrachromosomally. In this respect, it is also to be understood that the nucleotide sequence of the invention can be used to restore or create a mutant gene via homologous recombination.

Said host may be any prokaryotic or eukaryotic cell. Suitable prokaryotic/bacterial cells are those generally used for cloning like *E. coli*, *Salmonella typhimurium*, *Serratia marcescens* or *Bacillus subtilis*. Said eukaryotic host may be a mammalian cell, an amphibian cell, a fish cell, an insect cell, a fungal cell, a plant cell or a bacterial cell (e.g., *E. coli* strains HB101, DH5a, XL1 Blue, Y1090 and JM101). Eukaryotic recombinant host cells are preferred. Examples of eukaryotic host cells include, but are not limited to, yeast, e.g., *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis* or *Pichia pastoris* cells, cell lines of human, bovine, porcine, monkey, and rodent origin, as well as insect cells, including but not limited to, *Spodoptera frugiperda* insect cells and *Drosophila*-derived insect cells as well as zebra fish cells. Mammalian species-derived cell lines suitable for use and commercially available include, but are not limited to, L cells, CV-1 cells, COS-1 cells (ATCC CRL 1650), COS-7 cells (ATCC CRL 1651), HeLa cells (ATCC CCL 2), C1271 (ATCC CRL 1616), BS-C-1 (ATCC CCL 26) and MRC-5 (ATCC CCL 171).

In a particularly preferred embodiment said mammalian cell is a neuronal cell and/or a cultured cell like, inter alia, a HEK 293 (human embryonic kidney) cell, a CHO,

HeLa, NIH3T3, BHK, PC12 cell or a neuronal stem cell preferably derived from a mammal and more preferably from a human.

In another more preferred embodiment said amphibian cell is an oocyte. In an even more preferred embodiment said oocyte is a frog oocyte, particularly preferred a *Xenopus laevis* oocyte.

In a more preferred embodiment, the host according to the invention is a non-human transgenic organism. Said non-human organism may be a mammal, amphibian, a fish, an insect, a fungus or a plant. Particularly preferred non-human transgenic animals are *Drosophila* species, *Caenorhabditis elegans*, *Xenopus* species, zebra fish, *Spodoptera frugiperda*, *Autographa californica*, mice and rats. Transgenic plants comprise, but are not limited to, wheat, tobacco, parsley and *Arabidopsis*. Transgenic fungi are also well known in the art and comprise, inter alia, yeasts, like *S. pombe* or *S. cerevisiae*, or *Aspergillus*, *Neurospora* or *Ustilago* species or *Pichia* species.

In another embodiment, the present invention relates to a method for producing the polypeptide encoded by a nucleic acid molecule of the invention comprising culturing/raising the host of the invention and isolating the produced polypeptide.

A large number of suitable methods exist in the art to produce polypeptides in appropriate hosts. If the host is a unicellular organism or a mammalian or insect cell, the person skilled in the art can revert to a variety of culture conditions that can be further optimized without an undue burden of work. Conveniently, the produced protein is harvested from the culture medium or from isolated (biological) membranes by established techniques. Furthermore, the produced polypeptide may be directly isolated from the host cell. Said host cell may be part of or derived from a part of a host organism, for example said host cell may be part of the CNS of an animal or the harvestable part of a plant. Additionally, the produced polypeptide may be isolated from fluids derived from said host, like blood, milk or cerebrospinal fluid.

Additionally the present invention relates to polypeptides depicted in SEQ ID NOs: 5 to 12 which are encoded by the nucleic acid molecules of the invention or produced by the method of the invention. The polypeptide of the invention may accordingly be produced by microbiological methods or by transgenic mammals. It is also envisaged

that the polypeptide of the invention is recovered from transgenic plants. Alternatively, the polypeptide of the invention may be produced synthetically or semi-synthetically.

For example, chemical synthesis, such as the solid phase procedure described by Houghton Proc. Natl. Acad. Sci. USA (82) (1985), 5131-5135, can be used. Another method is in vitro translation of mRNA. A preferred method involves the recombinant production of protein in host cells as described above. For example, nucleotide acid sequences comprising all or a portion of any one of the nucleotide sequences according to the invention can be synthesized by PCR, inserted into an expression vector, and a host cell transformed with the expression vector. Thereafter, the host cell is cultured to produce the desired polypeptide, which is isolated and purified. Protein isolation and purification can be achieved by any one of several known techniques; for example and without limitation, ion exchange chromatography, gel filtration chromatography and affinity chromatography, high pressure liquid chromatography (HPLC), reversed phase HPLC, preparative disc gel electrophoresis. In addition, cell-free translation systems can be used to produce the polypeptides of the present invention. Suitable cell-free expression systems for use in accordance with the present invention include rabbit reticulocyte lysate, wheat germ extract, canine pancreatic microsomal membranes, E. coli S30 extract, and coupled transcription/translation systems such as the TNT-system (Promega). These systems allow the expression of recombinant polypeptides or peptides upon the addition of cloning vectors, DNA fragments, or RNA sequences containing coding regions and appropriate promoter elements. As mentioned supra, protein isolation/purification techniques may require modification of the proteins of the present invention using conventional methods. For example, a histidine tag can be added to the protein to allow purification on a nickel column. Other modifications may cause higher or lower activity, permit higher levels of protein production, or simplify purification of the protein.

In a further embodiment, the present invention relates to an antibody specifically directed to a polypeptide of the invention, wherein said antibody specifically reacts with an epitope generated and/or formed by the mutation in the ATP-gated ion channel P2X7R selected from the group consisting of:

- (i) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein the R (Arg), G (Gly), E (Glu), L (Leu), R (Arg), I (Ile) or R (Arg) residue corresponding to position 117, 150, 186, 191, 270, 568 or 578 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue; and
- (ii) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted.

With respect to preferred embodiments of (i) and (ii) the same applies as described above in connection with the nucleic acid molecules. The term "specifically" in this context means that the antibody reacts with the mutant P2X7R protein but not with a wild-type P2X7R protein. Preferably this term also means that such an antibody does not bind to other mutant forms of the P2X7R protein, in particular those described herein. Whether the antibody specifically reacts as defined herein above can easily be tested, inter alia, by comparing the reaction of said antibody with a wild-type ATP-gated ion channel P2X7R (or a subunit or a fragment thereof) with the reaction of said antibody with a mutant P2X7R polypeptide of the invention.

The antibody of the present invention can be, for example, polyclonal or monoclonal. The term "antibody" also comprises derivatives or fragments thereof which still retain the binding specificity. Techniques for the production of antibodies are well known in the art and described, e.g. in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. These antibodies can be used, for example, for the immunoprecipitation and immunolocalization of the polypeptides of the invention as well as for the monitoring of the presence of such polypeptides, for example, in recombinant organisms or in diagnosis. They can also be used for the identification of compounds interacting with the proteins according to the invention (as mentioned herein below). For example, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the polypeptide of the invention (Schier, Human

Antibodies Hybridomas 7 (1996), 97-105; Malmborg, J. Immunol. Methods 183 (1995), 7-13).

The present invention furthermore includes chimeric, single chain and humanized antibodies, as well as antibody fragments, like, inter alia, Fab fragments. Antibody fragments or derivatives further comprise F(ab')₂, Fv or scFv fragments; see, for example, Harlow and Lane, loc. cit.. Various procedures are known in the art and may be used for the production of such antibodies and/or fragments. Thus, the (antibody) derivatives can be produced by peptidomimetics. Further, techniques described for the production of single chain antibodies (see, inter alia, US Patent 4,946,778) can be adapted to produce single chain antibodies to polypeptide(s) of this invention. Also, transgenic animals may be used to express humanized antibodies to polypeptides of this invention. Most preferably, the antibody of this invention is a monoclonal antibody. For the preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples for such techniques include the hybridoma technique (Köhler and Milstein Nature 256 (1975), 495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor, Immunology Today 4 (1983), 72) and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc. (1985), 77-96). Techniques describing the production of single chain antibodies (e.g., US Patent 4,946,778) can be adapted to produce single chain antibodies to immunogenic polypeptides as described above. Furthermore, transgenic mice may be used to express humanized antibodies directed against said immunogenic polypeptides. It is in particular preferred that the antibodies/antibody constructs as well as antibody fragments or derivatives to be employed in accordance with this invention or capable to be expressed in a cell. This may, inter alia, be achieved by direct injection of the corresponding proteinaceous molecules or by injection of nucleic acid molecules encoding the same. Furthermore, gene therapy approaches are envisaged. Accordingly, in context of the present invention, the term "antibody molecule" relates to full immunoglobulin molecules as well as to parts of such immunoglobulin molecules. Furthermore, the term relates, as discussed above, to modified and/or altered antibody molecules, like chimeric and humanized antibodies. The term also relates to monoclonal or polyclonal antibodies as well as to recombinantly or

synthetically generated/synthesized antibodies. The term also relates to intact antibodies as well as to antibody fragments thereof, like, separated light and heavy chains, Fab, Fab/c, Fv, Fab', F(ab')₂. The term "antibody molecule" also comprises bifunctional antibodies and antibody constructs, like single chain Fvs (scFv) or antibody-fusion proteins. It is also envisaged in context of this invention that the term "antibody" comprises antibody constructs which may be expressed in cells, e.g. antibody constructs which may be transfected and/or transduced via, inter alia, viruses or vectors. It is in particular envisaged that such antibody constructs specifically recognize the polypeptides of the present invention. It is, furthermore, envisaged that said antibody construct is employed in gene therapy approaches.

The present invention relates also to an aptamer specifically binding to a polypeptide according to the invention wherein said aptamer reacts with an epitope of a polypeptide of the present invention. The present invention furthermore relates to an aptamer specifically directed to a corresponding nucleic acid molecule according to the invention.

In accordance with the present invention, the term "aptamer" means nucleic acid molecules that can bind to target molecules. Aptamers commonly comprise RNA, single stranded DNA, modified RNA or modified DNA molecules. The preparation of aptamers is well known in the art and may involve, inter alia, the use of combinatorial RNA libraries to identify binding sides (Gold, Ann. Rev. Biochem. 64 (1995), 763-797).

Furthermore, the present invention relates to a primer or pair of primers capable of specifically amplifying the nucleic acid molecules of the present invention. The term "primer" when used in the present invention means a single-stranded nucleic acid molecule capable of annealing the nucleic acid molecule of the present application and thereby being capable of serving as a starting point for amplification. Said term also comprises oligoribo- or desoxyribonucleotides which are complementary to a region of one of the strands of a nucleic acid molecule of the present invention. According to the present invention the term "pair of primers" means a pair of primers that are with respect to a complementary region of a nucleic acid molecule directed

in the opposite direction towards each other to enable, for example, amplification by polymérase chain reaction (PCR).

The term "amplifying" refers to repeated copying of a specified sequence of nucleotides resulting in an increase in the amount of said specified sequence of nucleotides. and allows the generation of a multitude of identical or essentially identical (i.e. at least 95% more preferred at least 98%, even more preferred at least 99% and most preferred at least 99.5% such as 99.9% identical) nucleic acid molecules or parts thereof. Such methods are well established in the art; see Sambrook et al. "Molecular Cloning, A Laboratory Manual", 2nd edition 1989, CSH Press, Cold Spring Harbor. They include polymerase chain reaction (PCR) and modifications thereof, ligase chain reaction (LCR) to name some preferred amplification methods.

When used in the context of primers the term "specifically" means that only the nucleic acid molecules as described herein above are amplified and nucleic acid molecules encoding the wild-type P2X7R ATP-gated receptor as depicted in SEQ ID NO: 1 are not amplified. Thus, a primer according to the invention is preferably a primer which binds to a region of a nucleic acid molecule of the invention which is unique for this molecule and which is not present in the wild-type P2X7R encoding sequence, i.e. the primer binds in a region in which one of the above described mutations occur. In connection with a pair of primers according to the invention it is possible that one of the primers of the pair is specific in the above described meaning or both of the primers of the pair are specific. In both cases, the use of such a pair of primers would allow to specifically amplify a mutant of the invention as described herein-above but not the wild-type P2X7R encoding sequence.

The 3'-OH end of a primer is used by a polymerase to be extended by successive incorporation of nucleotides. The primer or pair of primers of the present invention can be used, for example, in primer extension experiments on template RNA according to methods known by the person skilled in the art. Preferably, the primer or pair of primers of the present invention are used for amplification reactions on template RNA or template DNA, preferably cDNA or genomic DNA. The terms "template DNA" or "template RNA" refers to DNA or RNA molecules or fragments thereof of any source or nucleotide composition, that comprise a target nucleotide sequence as defined above. The primer or pair of primers can also be used for

hybridization experiments as known in the art. Preferably, the primer or pair of primers are used in polymerase chain reactions to amplify sequences corresponding to a sequence of the nucleic acid molecule of the present invention. It is known that the length of a primer results from different parameters (Gillam, Gene 8 (1979), 81-97; Innis, PCR Protocols: A guide to methods and applications, Academic Press, San Diego, USA (1990)). Preferably, the primer should only hybridize or bind to a specific region of a target nucleotide sequence. The length of a primer that statistically hybridizes only to one region of a target nucleotide sequence can be calculated by the following formula: $(\frac{1}{4})^x$ (whereby x is the length of the primer). For example a hepta- or octanucleotide would be sufficient to bind statistically only once on a sequence of 37 kb. However, it is known that a primer exactly matching to a complementary template strand must be at least 9 base pairs in length, otherwise no stable-double strand can be generated (Goulian, Biochemistry 12 (1973), 2893-2901). It is also envisaged that computer-based algorithms can be used to design primers capable of amplifying the nucleic acid molecules of the invention. Preferably, the primers of the invention are at least 10 nucleotides in length, more preferred at least 12 nucleotides in length, even more preferred at least 15 nucleotides in length, particularly preferred at least 18 nucleotides in length, even more particularly preferred at least 20 nucleotides in length and most preferably at least 25 nucleotides in length. The invention, however, can also be carried out with primers which are shorter or longer.

It is also envisaged that the primer or pair of primers is labeled. The label may, for example, be a radioactive label, such as ^{32}P , ^{33}P or ^{35}S . In a preferred embodiment of the invention, the label is a non-radioactive label, for example, digoxigenin, biotin and fluorescence dye or a dye.

In another preferred embodiment said primers are selected from the group consisting of SEQ ID NOs: 52 to 111.

In yet another embodiment, the present invention relates to a composition comprising a nucleic acid molecule, a vector, a polypeptide, an antibody, an aptamer and/or a primer or pair of primers of the invention.

The term "composition", as used in accordance with the present invention, relates to compositions which comprise at least one nucleic acid molecule, vector, polypeptide,

an antibody and/or primer or pair of primers of this invention. It may, optionally, comprise further molecules capable of altering the characteristics of the component of the invention thereby, for example, suppressing, blocking, modulating and/or activating their function which have neuroprotective, nootropic and/or antidepressive properties. The composition may be in solid, liquid or gaseous form and may be, inter alia, in the form of (a) powder(s), (a) tablet(s), (a) solution(s) or (an) aerosol(s).

In a preferred embodiment the composition according to the invention is a diagnostic composition, optionally further comprising suitable means for detection. As described above, the present invention is based on the surprising finding that mutations in the P2X7R protein are connected with affective disorders. Thus, this knowledge now allows to diagnose affective disorders in an easy way. The diagnostic composition comprises at least one of the aforementioned compounds of the invention. The diagnostic composition may be used, inter alia, for methods for determining the presence and/or expression of the nucleic acids and/or polypeptides of the invention. This may be effected by detecting, e.g., the presence of a corresponding gene in the genetic material of an individual or the presence of the corresponding mRNA which comprises isolation of DNA or RNA from a cell derived from said individual, contacting the DNA or RNA so obtained with a nucleic acid probe as described above under hybridizing conditions, and detecting the presence of mRNAs hybridized to the probe. Alternatively, the diagnostic composition may also be used for detecting the presence of a nucleic acid molecule of the invention by PCR. Furthermore, polypeptides of the invention can be detected with methods known in the art, which comprise, inter alia, immunological methods, like, RIA, FIA, ELISA, FACS or Western blotting.

Furthermore, the diagnostic composition of the invention may be useful, inter alia, in detecting the prevalence, the onset or the progress of a disease related to the expression of a polypeptide of the invention. Accordingly, the diagnostic composition of the invention may be used, inter alia, for assessing the prevalence, the onset and/or the disease status of affective disorders, as defined herein above. It is also contemplated that the diagnostic composition of the invention may be useful in discriminating (the) stage(s) of a disease.

The diagnostic composition optionally comprises suitable means for detection. The nucleic acid molecule(s), vector(s), host(s), antibody(ies), aptamer(s), polypeptide(s) described above are, for example, suitable for use in immunoassays in which they can be utilized in liquid phase or bound to a solid phase carrier. Examples of well-known carriers include glass, polystyrene, polyvinyl ion, polypropylene, polyethylene, polycarbonate, dextran, nylon, amyloses, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble or insoluble for the purposes of the invention.

Solid phase carriers are known to those in the art and may comprise polystyrene beads, latex beads, magnetic beads, colloid metal particles, glass and/or silicon chips and surfaces, nitrocellulose strips, membranes, sheets, duracytes and the walls of wells of a reaction tray, plastic tubes or other test tubes. Suitable methods of immobilizing nucleic acid molecule(s), vector(s), host(s), antibody(ies), aptamer(s), polypeptide(s), etc. on solid phases include but are not limited to ionic, hydrophobic, covalent interactions or (chemical) crosslinking and the like. Examples of immunoassays which can utilize said compounds of the invention are competitive and non-competitive immunoassays in either a direct or indirect format. Commonly used detection assays can comprise radioisotopic or non-radioisotopic methods. Examples of such immunoassays are the radioimmunoassay (RIA), the sandwich (immunometric assay) and the Northern or Southern blot assay. Furthermore, these detection methods comprise, inter alia, IRMA (Immune Radioimmunometric Assay), EIA (Enzyme Immuno Assay), ELISA (Enzyme Linked Immuno Assay), FIA (Fluorescent Immuno Assay), and CLIA (Chemiluminescent Immune Assay). Furthermore, the diagnostic compounds of the present invention may be employed in techniques like FRET (Fluorescence Resonance Energy Transfer) assays.

Appropriate labels and methods for labeling are known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include inter alia, fluorochromes (like fluorescein, rhodamine, Texas Red, etc.), enzymes (like horse radish peroxidase, β -galactosidase, alkaline phosphatase), radioactive isotopes (like ^{32}P , ^{33}P , ^{35}S or ^{125}I), biotin, digoxigenin, colloidal metals, chemi- or bioluminescent compounds (like dioxetanes, luminol or acridiniums).

A variety of techniques are available for labeling biomolecules, are well known to the person skilled in the art and are considered to be within the scope of the present invention and comprise, inter alia, covalent coupling of enzymes or biotinyl groups, phosphorylations, biotinylations, random priming, nick-translations, tailing (using terminal transferases). Such techniques are, e.g., described in Tijssen, "Practice and theory of enzyme immunoassays", Burden and von Knippenburg (Eds), Volume 15 (1985); "Basic methods in molecular biology", Davis LG, Dibner MD, Battey Elsevier (1990); Mayer, (Eds) "Immunochemical methods in cell and molecular biology" Academic Press, London (1987); or in the series "Methods in Enzymology", Academic Press, Inc.

Detection methods comprise, but are not limited to, autoradiography, fluorescence microscopy, direct and indirect enzymatic reactions, etc.

Said diagnostic composition may be used for methods for detecting the presence and/or abundance of a nucleic acid molecule of the invention in a biological and/or medical sample and/or for detecting expression of such a nucleic acid molecule (e.g. by determining the mRNA or the expressed polypeptide). Furthermore, said diagnostic composition may also be used in methods of the present invention, inter alia, for the detection of specific antagonists or agonists for P2X7R ATP-gated ion channels (see herein below).

In a further embodiment the present invention provides a method of diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R protein expressed in the cells of said individual is non-functional, shows an altered ATP-gating in comparison to the wild-type P2X7R protein or is over- or under-expressed in comparison to the P2X7R protein level of an unaffected individual.

The term "over- or under-expressed in comparison to the P2X7R protein level" in the context of the present invention means that the P2X7R protein level is higher or lower than the P2X7R level of a healthy individual, i.e. an individual not affected with an affective disorder. The over-expression may result, e.g. from an increased amount of P2X7R mRNA caused by enhanced transcription rates due to increased activity of the RNA-polymerase II. The amount of mRNA may accordingly lead to an increased translation and, thus, to a higher protein level of P2X7R. It may also

be possible that a higher amount of P2X7R protein is caused by increased stability of the protein. An under-expression of P2X7R protein may be caused by low transcription rates of the P2X7R gene and, thus, insufficient amounts of P2X7R mRNA give only rise to a low P2X7R protein amount. Another reason may be that the P2X7R protein is unstable and, thus, is not present in amounts comparable to the wild-type protein level.

The under- or over-expression of P2X7R protein may be determined by methods well-known to the person skilled in the art. These include, but are not limited, to methods for determining the amount of mRNA or the amount and/or activity of the protein. Examples are Northern Blot analysis or immuno based techniques, such as Western Blotting.

"Non-functional" means that the P2X7R protein has lost at least one functional property displayed by the wild-type P2X7R protein as described herein above. Preferably, "non-functional" means that the P2X7R protein does no longer function as a channel. Non-functionality may, e.g., be caused by the fact that one allele occurring in an individual codes for a P2X7R protein which leads to non-functional dimers (dominant negative mutation). Whether a P2X7R protein in an individual is functional or non-functional can be determined by the methods described herein above and in the examples.

The term "altered ATP-gating" means that the respective P2X7R protein reacts in a different way to ATP than the wild-type P2X7R protein. This can be determined as described in the appended examples or as described hereinabove.

In the context of diagnosis, not only the activity of the P2X7R could be of diagnostic value but also the amount of expression. For example, if a polymorphism affects RNA stability or translation efficiency, this could lead to lower expression of the P2X7 protein not only in the hippocampus but also in the blood. Therefore, one could speculate that a lower amount of P2X7 detected by western blot in blood cells could be related to depression.

Another aspect of the present invention is a method for diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R gene

sequence or encoded protein thereof comprises a mutation in comparison to the wild-type P2X7R sequence.

A preferred embodiment of the present invention is a method, wherein a mutation is a mutation in a P2X7R sequence as defined hereinabove and/or a nucleotide replacement or deletion selected from the following Table C indicating in column "Region of P2X7R" the region of the P2X7R genomic nucleotide sequence in which the replacement or deletion occurs, in column "Nucleotide" of Table C the nucleotide which is replaced by another nucleotide or the nucleotides which are deleted and in column "Position in wild-type" of Table C the corresponding position in the nucleotide sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1

Table C

| Region of P2X7R | Nucleotide | Position in wild-type |
|------------------------|-------------------|------------------------------|
| 5'UTR | T | 362 |
| 5'UTR | T | 532 |
| 5'UTR | A | 1100 |
| 5'UTR | A | 1122 |
| 5'UTR | C | 1171 |
| 5'UTR | T | 1351 |
| 5'UTR | G | 1702 |
| 5'UTR | T | 1731 |
| 5'UTR | C | 1860 |
| 5'UTR | C | 2162 |
| 5'UTR | C | 2238 |
| 5'UTR | A | 2373 |
| 5'UTR | G | 2569 |
| 5'UTR | G | 2702 |
| intron 1 | G | 3166 |
| intron 1 | C | 24778 |
| intron 1 | C | 24830 |
| exon 2 | T | 24942 |
| exon 3 | C | 26188 |
| exon 3 | A | 26308 |
| exon 3 | G | 26422 |
| intron 4 | G | 32394 |
| intron 4 | T | 32434 |
| exon 5 | G | 32493 |
| exon 5 | G | 32506 |

| | | |
|-----------|--------------------------------------|----------------|
| exon 5 | C | 32507 |
| exon 5 | C | 32548 |
| intron 5 | A | 32783 |
| intron 5 | T | 35309 |
| intron 5 | C | 35374 |
| intron 5 | A | 35378 |
| exon 6 | G | 35438 |
| exon 6 | T | 35454 |
| intron 6 | T | 35549 |
| intron 6 | G | 35641 |
| intron 6 | A | 35725 |
| intron 6 | T | 36001 |
| intron 6 | A | 36064 |
| intron 6 | deletion of GTTT | 36091 to 36094 |
| intron 6 | C | 36108 |
| intron 7 | C | 36374 |
| intron 7 | G | 36378 |
| intron 7 | T | 36387 |
| intron 7 | G | 36398 |
| intron 7 | C | 37439 |
| intron 7 | T | 37513 |
| exon 8 | C | 37604 |
| exon 8 | G | 37605 |
| exon 8 | G | 37623 |
| exon 8 | C | 37633 |
| intron 9 | C | 47214 |
| exon 11 | G | 47383 |
| exon 11 | C | 47411 |
| intron 11 | T | 47563 |
| intron 12 | C | 54307 |
| intron 12 | G | 54308 |
| exon 13 | C | 54399 |
| exon 13 | A | 54480 |
| exon 13 | C | 54523 |
| exon 13 | deletion of CCCTGAGAGCCACAGGTGCCT | 54562 to 54582 |
| exon 13 | A | 54588 |
| exon 13 | C | 54664 |
| exon 13 | G | 54703 |
| exon 13 | A | 54804 |
| exon 13 | G | 54834 |
| exon 13 | G | 54847 |
| 3'UTR | G | 54925 |
| 3'UTR | C | 55169 |
| 3'UTR | A | 55170 |
| 3'UTR | A | 55171 |
| 3'UTR | C | 55917 |

As indicated hereinabove, if the respective nucleotide which is replaced by another nucleotide is a purine base, it is preferred to be replaced by another purine base. If it is a pyrimidine base, it is preferred to be replaced by another pyrimidine base. It is also preferred that a purine base is replaced by a pyrimidine base and that a pyrimidine base is replaced by a purine base. Most preferably, the nucleotides indicated in Table C are replaced by the nucleotides indicated at the respective position in Table 12 hereinbelow (see Example 3).

In a preferred embodiment the present invention relates to diagnostic composition designed for use in a method in which the occurrence of the mutation in the ATP-gated ion channel P2X7R gene is determined by PCR, immunological methods and/or electrophysiological methods as described herein below and in the appended Examples. Additionally, it is possible to determine the occurrence of a mutation in the ATP-gated ion channel P2X7R as described hereinabove.

In yet another aspect the present invention relates to the use of a nucleic acid molecule, a vector, a polypeptide, an antibody, aptamer and/or a primer or pair of primers of the present invention for the preparation of a diagnostic composition for the detection of an affective disorder.

It is also envisaged that the present invention relates to methods of diagnosing an affective disorder of an individual comprising:

- (a) isolating DNA from cells obtained from an individual;
- (b) determining all or part of the nucleotide composition of the P2X7R gene; and
- (c) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism, mutation or allelic variation.

The term "gene" means a nucleotide sequence associated with the production of a protein, including promoter sequences, enhancer sequences, intron sequences, exon sequences, coding regions, 5' untranslated region (5'UTR), 3' untranslated region (3'UTR), and splice variants.

In a preferred embodiment of the described method the individual is a mammal and more preferably human. Moreover, the cells are preferably derived from skin, blood, urine or cerebral spinal fluid.

The method of the present invention allows for the diagnosis of an affective disorder according to the composition of a genetic marker corresponding to the P2X7R gene. As is demonstrated by the appended examples, polymorphisms in the P2X7R are genetically linked to patients suffering from an affective disorder.

In accordance with this embodiment of the present invention, the diagnosis of an affective disorder can, e.g., be effected by isolating cells from an individual, and isolating the genomic DNA of said cells. Such cells can be collected from body fluids, skin, hair, biopsies and other sources. Collection and analysis of cells from bodily fluids such as blood, urine and cerebrospinal fluid is well known to the art; see for example, Rodak, "Haematology: Clinical Principles & Applications" second ed., WB Saunders Co, 2002; Brunzel, "Fundamentals of Urine and Body Fluids Analysis", WB Saunders Co, 1994; Herndon and Brumback (Ed.), "Cerebrospinal Fluid", Kluwer Academic Pub., 1989. In addition, methods for DNA isolation are well described in the art; see, for example, Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd edition, Cold Spring Harbor Laboratory, 2001.

Once DNA has been isolated, various oligonucleotide primers spanning the P2X7R locus may be designed in order to amplify the genetic material by Polymerase Chain Reaction (PCR). Conventional methods for designing, synthesizing, producing said oligonucleotide primers and performing PCR amplification may be found in standard textbooks, see, for example Agrawal (Ed.), "Protocols for Oligonucleotides and Analogs: Synthesis and Properties (Methods in Molecular Biology, 20)", Humana Press, 1993; Innis et al. (Ed.), "PCR Applications: Protocols for Functional Genomics", Academic Press, 1999; Chen and Janes (Ed.), "PCR Cloning Protocols: From Molecular Cloning to Genetic", 2nd edition, Humana Press, 2002. Primers for the detection of P2X7R polymorphisms are also given in, but not limited to, SEQ ID NO: 52 to SEQ ID NO: 111. Once DNA has been amplified, nucleotide structure can be analysed by sequencing methods and compared to normal P2X7R DNA.

Sequencing may be performed manually by any molecular biologist of ordinary skills or by an automated sequencing apparatus. These procedures are common in the art, see, for example, Adams et al. (Ed.), "Automated DNA Sequencing and Analysis", Academic Press, 1994; Alphey, "DNA Sequencing: From Experimental Methods to Bioinformatics", Springer Verlag Publishing, 1997.

Detection and analysis of polymorphisms in P2X7R can also be performed using amplification refractory mutation system (ARMSTM), amplification refractory mutation system linear extension (ALEXTM), single-strand conformation polymorphism (SSCP), heteroduplex analysis, PCR-SSCP, fluorescent SSCP in an automated DNA sequencer, denaturing gradient gel electrophoresis, RNase protection assays, detection of mutations by sequence specific oligonucleotide hybridization, chemical cleavage methods, enzyme mismatch cleavage methods, cleavage fragment length methods, allele-specific oligonucleotide hybridization on DNA chips, and other such methods known in the art, see, for example Nollau et al, Clin. Chem. 43 (1997), 1114-1128; Burczak and Mardis (Ed.), "Polymorphism Detection & Analysis Techniques", Eaton Pub Co, 2000; Cotton et al. (Ed.), "Mutation Detection: A Practical Approach", Irl Press, 1998; Taylor (Ed.), "Laboratory Methods for the Detection of Mutations and Polymorphisms in DNA", CRC Press, 1997.

The present invention also relates to a method of diagnosing an affective disorder in an individual comprising:

- (a) isolating RNA from cells obtained from an individual;
- (b) converting the RNA into cDNA;
- (c) determining all or part of the nucleotide composition of the cDNA so obtained; and
- (c) analyzing said nucleotide composition for the presence of one or more polymorphism(s) or allelic variation.

With respect to the preferred embodiments the same applies as already described above.

Detection and analysis of polymorphisms in the P2X7R RNA can be performed according to the methods described above.

The present invention also relates to a method for diagnosing an affective disorder in an individual comprising:

- (a) isolating RNA or proteins from cells obtained from an individual;
- (b) determining the levels of P2X7R RNA or protein; and
- (c) comparing the levels of P2X7R RNA or protein with the corresponding levels from a normal individual not afflicted with an affective disorder.

With respect to the preferred embodiments the same applies as already described above.

As is demonstrated by the appended examples, a relationship exists between the expression, or protein level of P2X7R and an affective disorder. This and other embodiments of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

According to another aspect on the invention, there is provided a polynucleotide comprising at least 20 bases of the human P2X7R gene and comprising a mutation or polymorphism selected from any of the following:

Table 1: Novel polymorphisms in the human P2X7R

| Region in P2X7 | Polymorphism | Protein Modification |
|-----------------------|---------------------|-----------------------------|
| 5'UTR | 362 T-C | |
| 5'UTR | 532 T-G | |
| 5'UTR | 1100 A-G | |
| 5'UTR | 1122 A-G | |
| 5'UTR | 1171 C-G | |
| 5'UTR | 1702 G-A | |
| Intron01 | 3166 G-C | |
| Intron01 | 24778 C-T | |
| Intron01 | 24830 6C-T | |
| Exon03 | 26188 C-T | Arg117Trp |

| | | |
|----------|--|---|
| Intron03 | 26308 A-G | |
| Intron03 | 26422 G-A | |
| Intron04 | 32394 G-A | |
| Intron04 | 32434 T-C | |
| Exon05 | 32493 G-A | Gly150Arg |
| Exon05 | 32548 C-T | Silent Cys168 |
| Intron05 | 32783 A-C | |
| Exon06 | 35438 G-A | Glu186Lys |
| Exon06 | 35454 T-C | Leu191Pro |
| Intron06 | 35641 G-C | |
| Intron06 | 35725 A-C | |
| Intron06 | 36001 T-G | |
| Intron07 | 36378 G-A | |
| Intron07 | 36387 T-A | |
| Intron07 | 36398 G-C | |
| Exon08 | 37604 C-T | Arg270Cys |
| Exon08 | 37633 C-T | Silent Asp279 |
| Intron09 | 47214 C-T | |
| Intron11 | 47563 T-C | |
| Intron12 | 54307 C-T | |
| Intron12 | 54308 G-A | |
| Exon13 | 54562-54582 deletion of CCCTGAGA GCCACAGGTGCCT | deletion of 7aa 488 to 494 (PESHRCL) |
| Exon13 | 54804 A-T | Ile568Asn |
| Exon13 | 54834 G-A | Arg578Gln |
| 3'UTR | 55169 C-A | |
| 3'UTR | 55170 A-C | |
| 3'UTR | 55171 A-C | |
| 3'UTR | 55917 C-T | |
| 3'UTR | 54925 G-A | |

The polymorphism describes the position and the variation observed. The position and numbering of the polymorphism corresponds to the human P2X7R gene as defined in SEQ ID No 1. Primers used for SNP amplification and sequencing are shown in Table 1a and listed in SEQ ID NO: 52 to SEQ ID NO: 111.

Table 1a. Primer sequences for SNP amplification and sequencing

| Primer Name | Orientation | Sequence | Begin | End |
|--------------------|--------------------|---------------------------|--------------|------------|
| P2RX7_01.for | Sense | cgtaggacttggcgcttct | 2785 | 2803 |
| P2RX7_01.rev | Anti sense | gagcacgtctcagattcgaaa | 3224 | 3244 |
| P2RX7_02.for | Sense | ccatgaggcagggtatgactattc | 24665 | 24687 |
| P2RX7_02.rev | Antisense | ctcctggatctcaccaggtt | 25168 | 25187 |
| P2RX7_03.for | Sense | ctcgtccagctttgatattaagc | 25966 | 25988 |
| P2RX7_03.rev | Antisense | ggccctagtgctagaaccaga | 26426 | 26447 |
| P2RX7_04.for | Sense | attcatccgtcagtgcc | 30794 | 30811 |
| P2RX7_04.rev | Antisense | gccatgtgaattttctaccgat | 31277 | 31298 |
| P2RX7_05.for | Sense | ttcgttggtggttaggatggg | 32314 | 32333 |
| P2RX7_05.rev | Antisense | caaggatgctcagggtagtagc | 32805 | 32826 |
| P2RX7_06.for | Sense | cactaggtttgctgtatccatttct | 35277 | 35301 |
| P2RX7_06.rev | Antisense | gcaactgtgtgagagcttgg | 35731 | 35750 |
| P2RX7_07.for | Sense | tcaaccctgggtccagtgtg | 35950 | 35968 |
| P2RX7_07.rev | Antisense | caccaagtagctctcactcataagg | 36424 | 36448 |
| P2RX7_08.for | Sense | caataacacttggtgcgagttaggt | 37380 | 37403 |
| P2RX7_08.rev | Antisense | catcttggtgccttggaacc | 37750 | 37770 |
| P2RX7_09.for | Sense | gtgagtggtaatcctgctactgc | 45321 | 45343 |
| P2RX7_09.rev | Antisense | aggccactcctgtactcg | 45743 | 45761 |
| P2RX7_10_11.for | Sense | ccaagtcacagcatgaggc | 47119 | 47137 |
| P2RX7_10_11.rev | Antisense | accagcgacgtatccac | 47632 | 47649 |
| P2RX7_12.for | Sense | aagcatggggttccatttc | 50252 | 50268 |
| P2RX7_12.rev | Antisense | gcataaaagggactcctgctagta | 50691 | 50714 |
| P2RX7_13a.for | Sense | gcttacagaacacatgcatgg | 54232 | 54252 |
| P2RX7_13a.rev | Antisense | gcacctgtaggcacagtgc | 54739 | 54757 |
| P2RX7_13b.for | Sense | atcaccacctcagagctgttc | 54620 | 54640 |
| P2RX7_13b.rev | Antisense | gttaacatggctactgcagcc | 55203 | 55223 |
| P2XR7_13d.for | Sense | gcttagaaaggaggcgactcc | 54484 | 54504 |
| P2XR7_Pro13.for | Sense | ttgtgacatttgcaaggctgcc | 2617 | 2638 |
| P2XR7_Pro7.rev | Antisense | tctgaagctctgctcctgag | 1955 | 1974 |
| P2XR7_Pro8.rev | Antisense | ctcaccttctggcttccagt | 1611 | 1630 |
| P2XR7_Pro9.for | Sense | cttaccactcccaggactaa | 1496 | 1515 |
| P2XR7_Pro10.for | Sense | gtctgcctgttcaactgccat | 1149 | 1168 |
| P2XR7_Pro1.for | Sense | cagagaccttcagaaacttcg | 1841 | 1861 |
| P2XR7_Pro2.rev | Antisense | agatcaccagggacacagtg | 2261 | 2280 |
| P2XR7_Pro3.for | Sense | ctcaactccactttcctcgg | 2133 | 2152 |
| P2XR7_Pro4.rev | Antisense | cctttcacttttttgggtctcatg | 2655 | 2677 |
| P2XR7_Pro5.for | Sense | gggagaattctgaaaatgccc | 2691 | 2711 |
| P2XR7_Pro6.rev | Antisense | ggaccagagctctactcttc | 2951 | 2970 |
| P2XR7_Pro11.for | Sense | aggcatagatcgacctgcc | 2296 | 2315 |
| P2XR7_Pro12.rev | Antisense | aagaagcgccaagtcctacg | 2785 | 2804 |
| P2XR7_Pro14.for | Sense | gcaatccagactgaagttgac | 2051 | 2071 |
| P2XR7_Pro15.rev | Antisense | actctggtctgcagttggtg | 2428 | 2447 |
| P2XR7_Pro21.for | Sense | cctttaaaatcagagaccttcaga | 1831 | 1854 |
| P2XR7_Pro22.for | Sense | gccatcctctgaacacccat | 2708 | 2727 |
| P2XR7_3UTR10.for | Sense | cccttggaactcttgctatcg | 55804 | 55824 |
| P2XR7_3UTR1.for | Sense | ggcagtacagtggcttcaaga | 54858 | 54878 |
| P2XR7_3UTR2.rev | Antisense | gtgggacagtttgctgtgcct | 55150 | 55170 |

| | | | | |
|-----------------|-----------|-----------------------------|-------|-------|
| P2XR7_3UTR3.for | Sense | gagtccttaccaatagcagg | 55183 | 55202 |
| P2XR7_3UTR4.rev | Antisense | gtcaaagaatttgtggccacc | 55643 | 55663 |
| P2XR7_3UTR5.for | Sense | catgaactgtcttttaatgtgtaaag | 55515 | 55540 |
| P2XR7_3UTR6.rev | Antisense | gagatacgggtttcaccatggtg | 55955 | 55976 |
| P2XR7_3UTR7.for | Sense | aattagctgggcatggtgcg | 55992 | 56011 |
| P2XR7_3UTR8.rev | Antisense | ttgagatggagtctcgctctg | 56122 | 56140 |
| P2XR7_3UTR9.rev | Antisense | cactgtccacgtgactgctt | 56208 | 56227 |
| P2RX7_11.For | Sense | tcctacttcgggtctggttaagagatt | 47281 | 47305 |
| P2RX7_11.Rev | Antisense | gggcctaattttcgtgcát | 47591 | 47609 |
| P2RX7_13G.For | Sense | aagaacctagaacctgagggcctt | 54333 | 54355 |
| P2RX7_13G.Rev | Antisense | ttgagatgggaggcagctt | 54541 | 54559 |
| P2RX7_13H.For | Sense | ttcggctcccaggacat | 54773 | 54789 |
| P2RX7_13H.Rev | Antisense | cacagagctttgcaggtgaa | 55248 | 55267 |

Another aspect of the present invention is in the form of a diagnostic kit for affective disorders comprising a specific oligonucleotide probe, or primer corresponding to P2X7R polymorphisms. The diagnostic kit may comprise appropriate packaging and instructions for the use in the method of the invention. Said kit may further comprise appropriate buffer, and enzymes such as reverse transcriptase, and thermostable polymerases.

In a preferred embodiment of the invention, diagnosis can be performed on a mouse, rat or human. The invention is generally applied in vitro, e.g. using cells or other material obtained from an individual. However, it can also be applied on a living individual, or post mortem.

In accordance with the embodiments of the present invention, diagnosis of an affective disorder may be followed by prescription, or administration of an antidepressant drug. Administration and dosage of antidepressive drugs can vary between patients and are well known in the medical art, see, for example Benkert and Hippus, "Kompendium der Psychiatrischen Pharmakotherapie", Springer Verlag Publishing, 2000; Albers, "Handbook of Psychiatric Drugs: 2001-2002 Edition", Current Clinical Strategies Publishing, 2000. Preferred examples include between 5 mg and 80 mg per day, preferably 20 mg, fluoxetine; between 5 mg and 50 mg per day, preferably 20 mg, paroxetine; between 5 mg and 200 mg per day, preferably 50 mg, sertraline; between 5 mg and 300 mg per day, preferably 100 mg, fluvoxamine; between 5 mg and 100 mg per day, preferably 30 mg, mirtazapine; between 4 mg

and 50 mg, preferably 8 mg, reboxetine; between 5 mg and 600 mg per day, preferably 200 mg, nefazodone; between 450 mg and 1800 mg per day, preferably 900 mg, lithium carbonate.

The P2X7R protein is also useful for monitoring the efficacy and/or dosing of a drug or the likelihood of a patient to respond to a drug. Thus, in yet another embodiment the invention relates to a method for, monitoring the efficacy and/or dosing of a drug, e.g. an antidepressive drug, and/or the likelihood of a patient to respond to said drug which comprises determining the level of expression and/or activity of the P2X7R protein in a patient before and after administration of the respective drug. As presented in the examples below, treatment with an antidepressive drug results in an upregulation in P2X7R activity. In humans, P2X7R activity can be monitored by Positron Emission Tomography (PET) or Single Photon Emission Computerised Tomography (SPECT) using a radiolabelled ligand tracer for P2X7R. Examples of P2X7R ligands can be, but are not limited to, ATP, an antagonist binding P2X7R, an agonist binding P2X7R, or a small polynucleotide comprising at least 20 bases of the human P2X7R gene. A modulation of P2X7R activity, membrane distribution or expression levels would reflect the activity and potency of the antidepressive drug. Methods and techniques required for PET analysis are well known in the art, see, for example Paans and Vaalburg, Curr. Pharmac. Design 6 (2000), 1583-1591; van Waarde, Curr. Pharmac. Design. 6 (2000), 1593-1610; Paans et al, Methods 27 (2002), 195-207; Passchier et al., Methods 27 (2002), 278-286; Laruelle et al., Methods 27 (2002), 287-299.

In accordance with the present invention by the term "sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source containing polynucleotides or polypeptides or portions thereof. As indicated, biological samples include body fluids (such as blood, sera, plasma, urine, synovial fluid and spinal fluid) and tissue sources found to express the polynucleotides of the present invention. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. A biological sample which includes genomic DNA, mRNA or proteins is preferred as a source.

As described herein above, mutations of the P2X7R encoding gene can occur on DNA level or on mRNA level and may result in an altered expression of P2X7R or in the expression of P2X7R ATP-gated ion channels which show either an altered function or no function when compared to the wild-type P2X7R ATP-gated ion channel as described herein. Thus, various methods on DNA level, RNA level or protein level exist for determining whether the ATP-gated ion channel P2X7R gene shows a mutation as described herein above. Consequently, mRNA, cDNA, DNA and genomic DNA are the preferred nucleic acid molecules to be used in the below mentioned methods. Also polypeptides or fragments thereof are preferred if a mutation in the P2X7R ATP-gated ion channel protein as described herein is to be determined.

Preferably, a point mutation leading to the replacement of an amino acid residue at the positions as indicated in Table 1 of the corresponding wild-type P2X7R amino acid sequence depicted in SEQ ID NO: 3 or 4 by another amino acid can be determined by PCR. Said PCR is followed by a restriction fragment length polymorphism (RFLP) analysis if due to the point mutation a recognition site for a restriction endonuclease is generated which is not present in the wild-type nucleotide sequence or a recognition site for a restriction enzyme is created which does not occur in the wild-type P2X7R. More preferably said mutation can be determined by PCR using primers and conditions that allow only an amplification of the wild-type nucleotide sequence encoding the corresponding wild-type amino acid at the respective position, but not of the nucleotide sequence of a nucleic acid molecule encoding a different amino acid residue at the corresponding position. It is even more preferred that PCR is performed to determine a mutation using primers and conditions that allow no amplification if the wild-type nucleotide sequence is present, but only if another amino acid residue is encoded at the respective position. Particularly preferred is a method using PCR and primers under conditions that allow amplification of a fragment comprising at least the nucleotide residues encoding the amino acid residue corresponding to positions of SEQ ID NO: 1.

Said PCR is followed by e.g., sequencing and/or single strand conformation analysis (SSCA). Said fragment is preferably of at least 25 nucleotides in length, more preferred of at least 50 nucleotide in length, even more preferred of at least 75

nucleotides in length, particularly preferred of at least 100 nucleotides in length, more particularly preferred of at least 200 nucleotides in length, also more particularly preferred at least 250 nucleotides in length, even more particularly preferred at least 300 nucleotides in length and most particularly preferred at least 600 nucleotides in length. Said primers are preferably of at least 12 nucleotides in length, more preferred of at least 15 nucleotides in length, even more preferred of at least 18 nucleotides in length and most preferred of at least 21 nucleotides in length as depicted in SEQ ID NOs: 52 to 111. The temperature for annealing said primers is preferably at least 50°C, more preferred at least 55°C and most preferred at least 58°C. The temperature for denaturation is preferably at least 95°C for preferably at least 10 sec, more preferably at least 20 sec, even more preferred at least 30 sec and most preferred at least 60 sec. However, depending on the length and the G-C content of the nucleic acid sequence to be amplified the temperature for denaturation may be shorter or longer. The temperature for extension of the annealed primers is preferably at least 10 sec, more preferably at least 20 sec, even more preferred at least 30 sec and most preferred at least 60 sec. A PCR reaction comprising the aforementioned conditions is exemplified in the Examples herein below. The subsequent sequencing and/or SSCA is carried out as known in the art. Preferably, the PCR fragments are separated on a 10% polyacrylamide gel at 4°C or also preferred at room temperature. PCR fragments showing a SSCA band shift are amplified with the primers under conditions as mentioned above and are subsequently sequenced. Alternatively, it is also possible to directly sequence genomic DNA in order to determine whether a mutation in the CLCN2 gene has occurred. A direct genomic sequencing approach is, for example, demonstrated for baker's yeast in Horecka, Yeast 16 (2000), 967-970.

Preferably, a deletion is determined by using hybridization techniques as known in the art. In particular, a primer is designed as mentioned herein above that is capable to only hybridize to wild-type genomic DNA as depicted in SEQ ID NO: 1 but not to a nucleotide sequence comprising a deletion of a fragment between nucleotides 54562 and 54582 of SEQ ID NO: 1. Also preferred is the method of fluorescent in situ hybridization (FISH) for determining on whole chromosomes, in particular on chromosome 12q23-q24 that said chromosome has the above mentioned deletion. Even more preferred is that a deletion of nucleotide residues as described herein

may be determined by using PCR, wherein one primer of a pair of primers is located within the region of genomic DNA comprising said deletion. Preferably, said deletion is between nucleotide positions 54562 and 54582 as depicted in SEQ ID NO: 1. Thus, under the appropriate conditions no PCR fragment will result if the genomic DNA comprises said deletion. It is particularly preferred that PCR using primers which are located upstream or downstream of the deletion is performed to determine said deletion. Under appropriate conditions as mentioned herein above, both a fragment of genomic DNA of the wild-type nucleotide sequence as set forth in SEQ ID NO: 1 and a fragment of the nucleotide sequence comprising a deletion of preferably the nucleotides between positions 54562 and 54582 as depicted in SEQ ID NO: 1 will be amplified.

It is also possible to determine the above-described P2X7R mutations on the protein level. Some of the mutations described above lead to shortened versions of the P2X7R protein. Thus, it is conceivable to determine the occurrence of these mutations by determining the length or molecular weight of the P2X7R protein expressed in an individual, e.g. by SDS PAGE.

It is also possible to determine the mutations of the P2X7R ATP-gated channel as described herein by using the antibodies of the present invention. Said antibodies specific for said mutations of P2X7R proteins will be determined by assay techniques such as radioimmunoassays, competitive-binding assays, Western blot analysis and ELISA assay. Also preferred are classical immunohistological methods.

The finding, described in the present invention, that certain mutations in the P2X7R encoding gene and/or the corresponding protein are connected with affective disorder is indicative that the non- or dysfunction of the P2X7R protein is responsible for various forms of affective disorders. Thus, the finding of these mutations not only allows the diagnosis of affective disorders by determining whether the above-described mutations occur in an individual. It also allows to develop a treatment of affective disorders which has been diagnosed to be the result of a mutation in the P2X7R encoding gene. Such a treatment can, e.g., comprise the introduction of a nucleic acid molecule encoding a non-functional or functional wild-type P2X7R protein thereby restoring in said individual the P2X7R activity or the activation or

repression of (a) P2X7R gene(s) in vivo. The term "activation or repression" in this context means that the expression of the gene is either enhanced (activation) or reduced (repression). An enhancement of expression can, e.g., be achieved by increasing the efficiency of transcription initiation, for example, by using suitable compounds which have an activating effect on transcription. Alternatively, an enhancement can be achieved by replacing the naturally occurring promoter by a more efficient promoter.

A repression may be achieved by suppressing expression of the gene, e.g., by specifically suppressing transcription from the respective promoter by suitable compounds or by rendering the promoter less efficient or non-functional.

In another embodiment the present invention also relates to a pharmaceutical composition. In accordance with the present invention the term "pharmaceutical composition" relates to a composition comprising a nucleic acid molecule comprising a nucleotide sequence which encodes an ATP-gated ion channel P2X7R and which is selected from the group consisting of:

- (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as depicted in SEQ ID NO: 3 or 4;
- (b) a nucleotide sequence comprising the nucleotide sequence as depicted in SEQ ID NO: 1 or SEQ ID NO: 2;
- (c) a nucleotide sequence which hybridizes to the nucleotide sequence of (a) or (b); and
- (d) a nucleotide sequence which is degenerated as a result of the genetic code to the nucleotide sequence of (c).

Such pharmaceutical compositions comprise a therapeutically effective amount of a nucleic acid molecule encoding a functional P2X7R protein and, optionally, a pharmaceutically acceptable carrier. The pharmaceutical composition may be administered with a physiologically acceptable carrier to a patient, as described herein. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency or other generally recognized pharmacopoeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered.

Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium ion, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the aforementioned compounds, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In another preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilised powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition

is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The pharmaceutical composition of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

In vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems. Preferably, the pharmaceutical composition is administered directly or in combination with an adjuvant.

The pharmaceutical composition is preferably designed for the application in gene therapy. The technique of gene therapy has already been described above in connection with the nucleic acid molecules of the invention and all what has been said there also applies in connection with the pharmaceutical composition. For example, the nucleic acid molecule in the pharmaceutical composition is preferably in a form which allows its introduction, expression and/or stable integration into cells of an individual to be treated.

For gene therapy, various viral vectors which can be utilized, for example, adenovirus, herpes virus, vaccinia, or, preferably, an RNA virus such as a retrovirus. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can also incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a P2X7R sequence of interest encoding a functional P2X7R protein into the

viral vector, along with another gene which encodes, for example, the ligand for a receptor on a specific target cell, for example, the vector is now target specific.

Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a sugar, a glycolipid, or a protein. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing the inserted polynucleotide sequence.

Since recombinant retroviruses are preferably defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsidation. Helper cell lines which have deletions of the packaging signal include, but are not limited to w2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced. Alternatively, NIH 3T3 or other tissue culture cells can be directly transfected with plasmids encoding the retroviral structural genes gag, pol and env, by conventional calcium phosphate transfection. These cells are then transfected with the vector plasmid containing the genes of interest. The resulting cells release the retroviral vector into the culture medium. Another targeted delivery system for P2X7R polynucleotides is a colloidal dispersion system. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. The preferred colloidal system of this invention is a liposome. Liposomes are artificial membrane vesicles which are useful as delivery vehicles in vitro and in vivo. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley, et al., Trends Biochem. Sci.,

6:77, 1981). In addition to mammalian cells, liposomes have been used for delivery of polynucleotides in plant, yeast and bacterial cells. In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the genes of interest at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino, et al., *Biotechniques*, 6:682, 1988). The composition of the liposome is usually a combination of phospholipids, particularly high-phase-transition-temperature phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations. Examples of lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidylethanolamine, sphingolipids, cerebroside, and gangliosides. Particularly useful are diacylphosphatidylglycerols, where the lipid moiety contains from 14-18 carbon atoms, particularly from 16-18 carbon atoms, and is saturated. Illustrative phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine and distearoylphosphatidylcholine. The targeting of liposomes can be classified based on anatomical and mechanistic factors. Anatomical classification is based on the level of selectivity, for example, organ-specific, cell-specific, and organelle-specific. Mechanistic targeting can be distinguished based upon whether it is passive or active. Passive targeting utilizes the natural tendency of liposomes to distribute to cells of the reticulo-endothelial system (RES) in organs which contain sinusoidal capillaries.

In another aspect the present invention relates to a method of treating an affective disorder comprising administering a therapeutically effective amount of the pharmaceutical composition comprising a nucleotide sequence encoding a functional ATP-gated ion channel as described herein above to a subject suffering from said disorder.

In the context of the present invention the term "subject" means an individual in need of a treatment of an affective disorder. Preferably, the subject is a vertebrate, even more preferred a mammal, particularly preferred a human.

The term "administered" means administration of a therapeutically effective dose of the aforementioned nucleic acid molecule encoding a functional P2X7R protein to an individual. By "therapeutically effective amount" is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art and described above, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art.

The methods are applicable to both human therapy and veterinary applications. The compounds described herein having the desired therapeutic activity may be administered in a physiologically acceptable carrier to a patient, as described herein. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways as discussed below. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt %. The agents may be administered alone or in combination with other treatments.

The administration of the pharmaceutical composition can be done in a variety of ways as discussed above, including, but not limited to, orally, subcutaneously, intravenously, intra-arterial, intranodal, intramedullary, intrathecal, intraventricular, intranasally, intrabronchial, transdermally, intranodally, intrarectally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the candidate agents may be directly applied as a solution dry spray.

The attending physician and clinical factors will determine the dosage regimen. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A typical dose can be, for example, in the range of 0.001 to 1000 μg ; however, doses below or above this exemplary range are envisioned, especially considering the aforementioned factors.

The dosages are preferably given once a week, however, during progression of the treatment the dosages can be given in much longer time intervals and in need can be given in much shorter time intervals, e.g., daily. In a preferred case the immune response is monitored using herein described methods and further methods known to those skilled in the art and dosages are optimized, e.g., in time, amount and/or composition. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately 10^6 to 10^{12} copies of the DNA molecule. If the regimen is a continuous infusion, it should also be in the range of 1 μ g to 10 mg units per kilogram of body weight per minute, respectively. Progress can be monitored by periodic assessment. The pharmaceutical composition of the invention may be administered locally or systemically. Administration will preferably be parenterally, e.g., intravenously. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium ion solution, Ringer's dextrose, dextrose and sodium ion, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like.

It is also envisaged that the pharmaceutical compositions are employed in co-therapy approaches, i.e. in co-administration with other medicaments or drugs, for example other drugs for preventing, treating or ameliorating affective disorders.

Another aspect of the present invention is a pharmaceutical composition comprising a compound, the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells or comprising a nucleic acid molecule the expression of which in cells or the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells. Said pharmaceutical

composition may be useful for treating individuals having an increased amount of the P2X7R protein or expression level as described hereinabove.

Preferably, the above-mentioned pharmaceutical composition is an antisense nucleic acid, a ribozyme, a co-suppressive nucleic acid, iRNA or siRNA.

An siRNA approach is, for example, disclosed in Elbashir ((2001), Nature 411, 494-498)). It is also envisaged in accordance with this invention that for example short hairpin RNAs (shRNAs) are employed in accordance with this invention as pharmaceutical composition. The shRNA approach for gene silencing is well known in the art and may comprise the use of st (small temporal) RNAs; see, inter alia, Paddison (2002) Genes Dev. 16, 948-958.

As mentioned above, approaches for gene silencing are known in the art and comprise "RNA"-approaches like RNAi or siRNA. Successful use of such approaches has been shown in Paddison (2002) loc. cit., Elbashir (2002) Methods 26, 199-213; Novina (2002) Mat. Med. June 3, 2002; Donze (2002) Nucl. Acids Res. 30, e46; Paul (2002) Nat. Biotech 20, 505-508; Lee (2002) Nat. Biotech. 20, 500-505; Miyagishi (2002) Nat. Biotech. 20, 497-500; Yu (2002) PNAS 99, 6047-6052 or Brummelkamp (2002), Science 296, 550-553. These approaches may be vector-based, e.g. the pSUPER vector, or RNA polIII vectors may be employed as illustrated, inter alia, in Yu (2002) loc. cit.; Miyagishi (2002) loc. cit. or Brummelkamp (2002) loc. cit.

A compound which leads to a reduction of the expression of the P2X7R gene may, e.g., be a compound which acts on the regulatory region of the gene and thereby reduces the level of transcription. Such compounds can be identified by methods as described herein below.

The invention also relates to the use of a nucleic acid molecule encoding a functional P2X7R protein as described herein above in connection with the pharmaceutical composition for the preparation of a pharmaceutical composition for treating an affective disorder.

Furthermore, the present invention relates to a method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic

acid molecule according to the invention or a therapeutically effective amount of the corresponding encoded polypeptide to a subject suffering from said disorder.

In another preferred embodiment the present invention relates to a pharmaceutical composition comprising, inter alia, the polynucleotides according to the present invention, i.e. polynucleotides having mutations and/or deletions as described hereinabove. Such pharmaceutical compositions may, e.g., be useful for treating individuals having an increased or amount of the P2X7R protein or having a P2X7R protein showing an increased activity which can be determined as described hereinabove. It is envisaged that, e.g. a non-functional P2X7R protein comprised by said pharmaceutical composition is incorporated in a P2X7R complex which naturally exists in cells as described hereinabove. It is also envisaged that the above-described techniques for gene therapy can be used for treating an individual with the nucleic acid molecules of the present invention, *mutatis mutandis*.


With respect to the possible modes of administration and preferred embodiments the same applies as has been set forth above.

Additionally, the present invention also envisages the use of the nucleic acid molecules, the vectors, the polypeptides, the antibody and/or the aptamer according to the invention for the preparation of a pharmaceutical composition for the treatment of an affective disorder.

A further aspect of the present invention is the use of a modulator of P2X7R activity or expression for the preparation of a pharmaceutical composition for treating an affective disorder. In the context of the present invention the term "modulator" means (a) compound(s), a complex of compounds, (a) substance(s) or complex of substances which can modify, i.e. modulate the activity of P2X7R or the expression of P2X7R. The modulation can, for example, occur at the protein level. Particularly, the P2X7R protein may interfere with the modulator such that it is either more active or less active. The modulation can also occur on nucleic acid level. Namely, the gene is transcribed more frequently or less frequently giving rise to more or less protein. Modulation can also influence RNA or protein stability.

In a preferred embodiment said modulator is selected from the group consisting of piperidine and piperazine derivatives, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate and 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate as, for example, described in WO 99/29660, WO 99/29661; WO 99/296896; WO 00/61569; WO 01/42194; WO 01/44170; WO 01/44213; WO 00/71529; WO 01/46200. The following compounds illustrate compounds which are also preferred to be used as modulators of P2X7R activity.


A compound of general formula:



where A is phenyl or a 5- or 6-membered heterocyclic ring containing one or two heteroatoms selected from O, N or S; and optionally substituted by C₁₋₆alkyl, halogen, nitro, amino, alkylamino, CF₃, SO₂Me, NHSO₂Me or cyano; B is C=O, NH or SO₂; X is C=O, CH(Me), O or (CH₂)_p where p is 0 or 1; Y is O, CH₂, NH or S; Z is C=O or SO₂, provided that when Z is C=O, then Y is O, CH₂ or S; R is hydrogen or C₁₋₆alkyl; R¹ is hydrogen, halogen; R² is phenyl optionally substituted by CO₂H, CO₂alkyl, CONH₂ or R² is OH, NHR³, NHCH(R⁴)(CHR⁵)_nR⁶, NH-R⁷-R⁸, SO₂NHalkyl, NHCOalkyl, NHSO₂alkyl, morpholine, NR⁹R¹⁰, piperazine substituted by phenyl, alkoxyphenyl, pyridyl or fluorophenyl; n is 0, 1 or 2; R³ is hydrogen, a bi- or tricyclic saturated ring system optionally containing a nitrogen atom, piperidiny, alkylpyrrolidine, ethynylcyclohexyl, a 5-membered aromatic ring containing 2 or 3 heteroatoms, C₄₋₆ cycloalkyl optionally substituted by alkyl, cyano or hydroxy, or C₁₋₈ alkyl optionally containing an oxygen atom in the alkyl chain and being optionally substituted by one or more substituents selected from ethynyl, cyano, fluoro, dialkylamino, hydroxy, thioalkyl, CO₂R¹¹ or CONH₂; R⁴ is hydrogen or alkyl optionally substituted by hydroxy or alkoxy; R⁵ is hydrogen or hydroxy; R⁶ is CO₂R¹¹, NHCO₂R¹², CONH₂ or a 5 or 6-membered saturated ring containing an oxygen atom, a 5-membered heterocyclic ring containing one or two heteroatoms selected from O,

N or S, or phenyl optionally substituted by one or more groups selected from alkyl, hydroxy, amino, alkoxy, or nitro; R^6 is alkyl; R^7 is a cyclopentane ring; R^8 is phenyl; R^9 and R^{10} are independently hydrogen, benzyl, alkenyl, cycloalkyl, alkyl optionally substituted by hydroxy, alkoxy, cyano, dialkylamino, phenyl, pyridyl or CO_2R^{11} or R^9 and R^{10} together form a 5- to 7-membered saturated or partially saturated ring optionally containing a further heteroatom and optionally substituted by one or more groups selected from alkyl (optionally containing an oxygen atom in the chain and optionally substituted 14 by hydroxy), COalkyl, CO_2R^{11} , $COR^{13}R^{14}$, CHO or piperidine, R^{11} is hydrogen or alkyl; R^{12} is alkyl; and R^{13} and R^{14} are independently hydrogen or alkyl, is or a pharmaceutically acceptable salt or solvate thereof.


A compound of general formula:



wherein A represents a group CH_2 or an oxygen atom; B represents a hydrogen or halogen atom; D represents a group CH_2 , OCH_2 , $NHCH_2$ or CH_2CH_2 ; R represents a phenyl, benzothiazolyl, indolyl, indazolyl, purinyl, pyridyl, pyrimidinyl or thiophenyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or a cyano, carboxyl, hydroxyl, nitro, halo- C_1 - C_6 -alkyl, $-N(R^1)-C(=O)-R^2$, $-C(O)-NR^3R^4$, $-NR^5R^6$, C_3 - C_8 -cycloalkyl, 3- to 8-membered heterocyclyl, C_3 - C_8 -cycloalkyloxy, C_1 - C_6 -alkylcarbonyl, phenoxy, benzyl, C_1 - C_6 -alkylthio, phenylthio, C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -alkylsulphinyl or C_1 - C_6 -alkylsulphonyl group, or a C_1 - C_6 -alkyl or C_1 - C_6 -alkoxy group optionally substituted by one or more substituents independently selected from a halogen atom or an amino, carboxyl, hydroxyl, C_1 - C_6 -alkoxy, (di) C_1 - C_6 -alkylamino, C_1 - C_6 -alkoxycarbonyl, imidazolyl, morpholinyl, piperidinyl or pyrrolidinyl group; R^1 represents a hydrogen atom or a C_1 - C_6 -alkyl or C_3 - C_8 -cycloalkyl group; R^2 represents a C_1 - C_6 -alkyl or C_3 - C_8 -cycloalkyl group; and R^3 , R^4 , R^5 and R^6 each independently represent a hydrogen




atom or a C_1-C_6 -alkyl or C_3-C_8 -cycloalkyl group; with the provisos that when A is CH_2 , B is H and D is CH_2 , then R does not represent a phenyl, ortho-carboxyphenyl, methylphenyl or para-phenoxyphenyl group, and that when A is CH_2 , D is CH_2 or CH_2CH_2 and R represents a substituted phenyl group, the substituent or substituents present do not comprise, in an ortho position, a C_1-C_6 -alkoxy group substituted by an amino, (di) C_1-C_6 -alkylamino, imdazolyl, morpholiny, piperidiny or pyrrolidiny group; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




(I)

wherein x represents 1 or 2; A represents a group CH_2 or an oxygen atom; B represents a hydrogen or halogen atom; R represents a phenyl, pyridyl, indolyl, indazolyl, pyrimidinyl or thiophenyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or an amino, cyano, carboxyl, hydroxyl, nitro, C_1-C_6 -alkyl, halo- C_1-C_6 -alkyl, $-N(R^1)-C(=O)-R^2$, $-C(O)NR^3R^4$, $-NR^5R^6$, C_3-C_8 -cycloalkyl, 3- to 8-membered heterocyclyl, C_3-C_8 -cycloalkyloxy, C_1-C_6 -alkylcarbonyl, C_1-C_6 -alkoxycarbonyl, C_1-C_6 -alkylsulphiny or C_1-C_6 -alkylsulphonyl group, or a C_1-C_6 -alkoxy, C_1-C_6 -alkylamino, phenoxy, benzyl, C_1-C_6 -alkylthio or phenylthio group optionally substituted by one or more substituents independently selected from a 15 halogen atom or an amino, cyano, carboxyl, hydroxyl, nitro, 1-pyrrolidiny, 1-piperidiny, C_1-C_6 -alkyl, C_1-C_6 -alkoxy, (di) C_1-C_6 -alkylamino, halo- C_1-C_6 -alkyl, C_1-C_6 -alkoxycarbonyl or one of the following groups:



R^1 represents a hydrogen atom or a $\text{C}_1\text{-C}_6$ -alkyl or $\text{C}_3\text{-C}_8$ -cycloalkyl group; R^2 represents a $\text{C}_1\text{-C}_6$ -alkyl or $\text{C}_3\text{-C}_8$ -cycloalkyl group; R^3 and R^4 each independently represent a hydrogen atom or a $\text{C}_1\text{-C}_6$ -alkyl or $\text{C}_3\text{-C}_8$ -cycloalkyl group; R^5 represents a hydrogen atom or a $\text{C}_1\text{-C}_6$ -alkyl or $\text{C}_3\text{-C}_8$ -cycloalkyl group; R^6 represents a $\text{C}_3\text{-C}_8$ -cycloalkyl group and, additionally, a $\text{C}_1\text{-C}_6$ -alkyl group when R^5 is not a hydrogen atom; R^7 represents a hydrogen atom or a $\text{C}_1\text{-C}_6$ -alkyl or $\text{C}_3\text{-C}_8$ -cycloalkyl group; R^8 represents a $\text{C}_1\text{-C}_6$ -alkyl or $\text{C}_3\text{-C}_8$ -cycloalkyl group; R^9 represents a hydrogen atom or a hydroxyl group; and R^{10} represents a hydrogen atom or a phenyl or imidazolyl group; with the provisos that R does not represent an unsubstituted pyridyl group when A represents a group CH_2 and B represents a hydrogen atom, and that when R represents a substituted phenyl, indolyl or indazolyl group, the substituent or substituents present do not comprise an amido, carboxyl, (di) $\text{C}_1\text{-C}_6$ -alkylamido or $\text{C}_1\text{-C}_6$ -alkoxycarbonyl group in an ortho position; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




wherein D represents CH_2 or CH_2CH_2 ; E represents $\text{C}(\text{O})\text{NH}$ or $\text{NHC}(\text{O})$; R^1 and R^2 each independently represent a hydrogen or halogen atom, or an amino, nitro, $\text{C}_1\text{-C}_6$ -alkyl or trifluoromethyl group; R^3 represents a group of formula:




X represents an oxygen or sulphur atom or a group NH, SO or SO₂; Y represents an oxygen or sulphur atom or a group NR¹¹, SO or SO₂; Z represents a group -OH, -SH, -CO₂H, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylsulphinyl, C₁-C₆-alkylsulphonyl, -NR⁶R⁷, -C(O)NR⁸R⁹, imidazolyl, 1-methylimidazolyl, -N(R¹⁰)C(O)-C₁-C₆-alkyl, C₁-C₆-alkylcarbonyloxy, C₁-C₆-alkoxycarbonyloxy, -OC(O)NR¹²R¹³, -OCH₂OC(O)R¹⁴, OCHOC(O)OR¹⁵ or -OC(O)OCH₂OR¹⁶, R⁴ represents a C₂-C₆-alkyl group; R⁵ represents a C₁-C₆-alkyl group; R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹², and R¹³ each independently represent a hydrogen atom, or a C₁-C₆-alkyl group optionally substituted by at least one hydroxyl group; R¹¹ represents a hydrogen atom, or a C₁-C₆-alkyl group optionally substituted by at least one substituent independently selected from hydroxyl and C₁-C₆-alkoxy; and R¹⁴, R¹⁵ and R¹⁶ each independently represent a C₁-C₆-alkyl group; with the provisos that (i) when E represents NEC(O), X represents O, S or NH and Y represents O, then Z represents -NR⁶R⁷ where R⁶ represents a hydrogen atom and R⁷ represents either a hydrogen atom or a C₁-C₆-alkyl group substituted by at least one hydroxyl group, and (ii) when E represents NHC(O), X represents O, S or NH, Y represents NH, and R⁵ represents CH₂CH₂, then Z is not -OH or imidazolyl; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:



wherein D represents CH₂ or CH₂CH₂; E represents C(O)NH or NHC(O); R¹ and R² each independently represent hydrogen, halogen, amino, nitro, C₁-C₆-alkyl or

trifluoromethyl, but R^1 and R^2 may not both simultaneously represent hydrogen; R^3 represents a group of formula




R^4 represents a C_1 - C_6 -alkyl group; X represents an oxygen or sulphur atom or a group NR^{13} , SO or SO_2 ; R^5 represents hydrogen, or R^5 represents C_1 - C_6 -alkyl or C_2 - C_6 -alkenyl, each of which may be optionally substituted by at least one substituent selected from halogen, hydroxyl, (di) C_1 - C_6 -alkylamino, $-Y-R^6$,




NH_2 , and a 5- or 6-membered heteroaromatic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur which heteroaromatic ring may itself be optionally substituted by at least one substituent selected from halogen, hydroxyl and C_1 - C_6 -alkyl; Y represents an oxygen or sulphur atom or a group NH , SO or SO_2 ; R^6 represents a group $-R^7Z$ where R^7 represents a C_2 - C_6 -alkyl group and Z represents an $-OH$, $-CO_2H$, $-NR^8R^9$, $-C(O)NR^{10}R^{11}$ or $-N(R^{12})C(O)-C_1$ - C_6 -alkyl group, and, in the case where Y represents an oxygen or sulphur atom or a group NH , R^6 additionally represents hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, $-C(O)NR^{14}R^{15}$, $-CH_2OC(O)R^{16}$, $-CH_2OC(O)OR^{17}$ or $-C(O)OCH_2OR^{18}$; R^8 , R^9 , R^{10} , R^{11} and R^{12} each independently represent a hydrogen atom or a C_1 - C_6 -alkyl group; R^{13} represents hydrogen, C_3 - C_8 -cycloalkyl, C_3 - C_8 -cycloalkylmethyl, or R^{13} represents a C_1 - C_6 -alkyl group optionally substituted by at least one substituent selected from hydroxyl to and C_1 - C_6 -alkoxy; and R^{14} , R^{15} , R^{16} , R^{17} , and R^{18} each independently represent a C_1 - C_6 -alkyl group; with the proviso that when E is $C(O)NH$, X is O , NH or $N(C_1$ - C_6 -alkyl), then R^5 is other than a hydrogen atom or an unsubstituted C_1 - C_6 -alkyl group; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




wherein m represents 1, 2 or 3; each R^1 independently represents a hydrogen or halogen atom; A represents $C(O)NH$ or $NHC(O)$; Ar represents a group



X represents a bond, an oxygen atom or a group CO , $(CH_2)_{1-6}$, $CH=$, $(CH_2)_{1-6}O$, $O(CH_2)_{1-6}$, $O(CH_2)_{2-6}O$, $O(CH_2)_{2-3}O(CH_2)_{1-3}$, $CR'(OH)$, $(CH_2)_{1-3}O(CH_2)_{1-3}$, $(CH_2)_{1-3}O(CH_2)_{2-3}O$, NR^5 , $(CH_2)_{1-6}NR^5$, $NR^5(CH_2)_{1-6}$, $(CH_2)_{1-3}NR^5(CH_2)_{1-3}$, $O(CH_2)_{2-6}NR^5$, $O(CH_2)_{2-3}NR^5(CH_2)_{1-3}$, $(CH_2)_{1-3}NR^5(CH_2)_{2-3}O$, $NR^5(CH_2)_{2-6}O$, $NR^5(CH_2)_{2-3}O(CH_2)_{1-3}$, $CONR^5$, NR^5CO , $S(O)_n$, $S(O)_nCH_2$, $CH_2S(O)_n$, SO_2NR^5 or NR^5SO_2 ; n is 0, 1 or 2; R^1 represents a hydrogen atom or a C_1 - C_6 -alkyl group; one of R^2 and R^3 represents a halogen, cyano, nitro, amino, hydroxyl, or a group selected from (i) C_1 - C_6 -alkyl optionally substituted by at least one C_3 - C_6 -cycloalkyl, (ii) C_3 - C_8 -cycloalkyl, (iii) C_1 - C_6 -alkyloxy optionally substituted by at least one C_3 - C_6 -cycloalkyl, and (iv) C_3 - C_8 -cycloalkyloxy, each of these groups being optionally substituted by one or more fluorine atoms, and the other of R^2 and R^3 represents a hydrogen or halogen atom; either R^4 represents a 3- to 9-membered saturated or unsaturated aliphatic heterocyclic ring system containing one or two nitrogen atoms and optionally an oxygen atom, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, carboxyl, cyano, C_1 - C_6 -alkyl, C_1 - C_6 -hydroxyalkyl, $-NR^6R^7$, $-(CH_2)_rNR^6R^7$ and $CONR^6R^7$, r is 1, 2, 3, 4, 5 or 6; R^5 represents a hydrogen atom or a C_1 - C_6 -alkyl or C_3 - C_8 -cycloalkyl group; R^6 and R^7 each independently represent a hydrogen atom or a C_1 - C_6 -alkyl, C_2 - C_6 -hydroxyalkyl or C_3 - C_8 -cycloalkyl group, or R^6 and R^7 together with the nitrogen

atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring; with the provisos that, (a) when A represents C(O)NH and R⁴ represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and (b) when A represents C(O)NH and X represents a group (CH₂)₁₋₆ or O(CH₂)₁₋₆, then R⁴ does not represent an unsubstituted imidazolyl, unsubstituted morpholinyl, unsubstituted piperidinyl or unsubstituted pyrrolidinyl group, and (c) when A represents NHC(O) and R⁴ represents an unsubstituted 3- to 8-membered saturated aliphatic heterocyclic ring system containing one nitrogen atom, then X is other than a bond, and (d) when A represents NHC(O) and X represents O(CH₂)₁₋₆, NH(CH₂)₁₋₆ or SCH₂, then R⁴ does not represent an unsubstituted 1-piperidinyl or unsubstituted 1-pyrrolidinyl group, and (e) when A represents NHC(O) and X represents O(CH₂)₂₋₃NH(CH₂)₂, then R⁴ does not represent an imidazolyl group; or a pharmaceutically acceptable salt or solvate thereof.


A compound of general formula:




X represents a nitrogen atom or a group C(R⁵); Y represents an oxygen or sulphur atom or a group NR⁶; either R¹ and R² each independently represent a hydrogen atom or a C₁-C₄-alkyl group but do not both simultaneously represent a hydrogen atom, or R¹ and R² together represent a group -CH₂ZCH₂-; Z represents a bond, an oxygen or sulphur atom or a group CH₂ or NR⁷; m is 0 or 1; R³ represents a 5- to 10-membered unsaturated ring system which may comprise from 1 to 4 ring heteroatoms independently selected from nitrogen, oxygen and sulphur, the ring system being optionally substituted by one or more substituents independently selected from halogen, nitro, cyano, NR⁸R⁹, C₁-C₄-alkyl-C(O)NH-, NHR¹²C(O)-, C₁-C₄-alkyl-SO₂-, C₁-C₄-alkyl-SO₂NH-, C₁-C₄-alkyl-NHSO₂-, C₁-C₄-alkoxy, and C₁-C₄-alkyl optionally substituted by one or more fluorine atoms; R⁴ represents a phenyl or pyridinyl group, each of which is substituted in an ortho position with a substituent

selected from halogen, C₁-C₄-alkoxy, C₁-C₄-alkylthio, and C₁-C₄-alkyl optionally substituted by one or more fluorine atoms, the phenyl or pyridinyl group being optionally further substituted by one or more substituents independently selected from halogen, cyano, hydroxyl, C₁-C₄-alkylthio, C₁-C₄-alkyl-NH-, NHR¹³-C₁-C₄-alkyl-, C₁-C₄-alkyl-SO₂-, C₁-C₄-alkyl-SO₂NH-, C₁-C₄-alkyl-NHSO₂-, C₁-C₄-alkyl-C(O)NH-, C₁-C₄-alkyl-NHC(O)-, -D-G, C₁-C₄-alkoxy optionally substituted by -NR¹⁴R¹⁵ or by R¹⁶, and C₁-C₄-alkyl optionally substituted by one or more fluorine atoms or by one or more hydroxyl groups, or R⁴ represents a 9- or 10-membered unsaturated bicyclic ring system which may comprise from 1 to 4 ring heteroatoms independently selected from nitrogen, oxygen and sulphur, the bicyclic ring system being optionally substituted by one or more substituents independently selected from halogen, oxo, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio and -NR¹⁰R¹¹; D represents an oxygen atom or a group (CH₂)_{sub.n} or CH₂NH; n is 1, 2 or 3; G represents a piperazinyl, morpholinyl or 2,5-diazabicyclo[2.2.1]heptyl group, or G represents a piperidinyl group optionally substituted by amino; R⁵ represents a hydrogen atom, or a hydroxyl or C₁-C₄-alkoxy group; R⁶ represents a hydrogen atom, or a cyano, nitro, hydroxyl, C₁-C₄-alkyl or C₁-C₄-alkoxy group; R⁷, R⁸ and R⁹ each independently represent a hydrogen atom or a C₁-C₄-alkyl group; R¹⁰ and R¹¹ each independently represent a hydrogen atom or a C₁-C₄-alkyl group, or R¹⁰ and R¹¹ together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocyclic ring comprising one or two ring nitrogen atoms; R¹² represents a hydrogen atom, or a C₁-C₄-alkyl group optionally substituted by amino; R¹³ represents a hydrogen atom, or a C₁-C₄-alkyl group optionally substituted by hydroxyl; R¹⁴ and R¹⁵ each independently represent a hydrogen atom or a C₁-C₄-alkyl group optionally substituted by hydroxyl, or R¹⁴ and R¹⁵ together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated heterocyclic ring comprising one or two ring nitrogen atoms; and R¹⁶ represents a 1-(C₁-C₄-alkyl)-piperidinyl group; with the proviso that when m is 0, X is N and Y is O, then R⁴ does not represent 2-benzothiazolyl; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:




wherein: each R^1 independently represents a hydrogen or halogen atom, or a trifluoromethyl, cyano, nitro, C_1 - C_6 -alkyl or C_1 - C_6 -alkoxy group; T represents an oxygen atom or a group NH; U represents an oxygen or sulphur atom or a group NH; Ar represents a group:



X represents a bond, an oxygen atom or a group CO , CH_2 , CH_2O , $O(CH_2)_m$, CH_2OCH_2 , NR^5 , CH_2NR^5 , NR^5CH_2 , $CH_2NR^5CH_2$, $CONR^5$, $S(O)_n$ or SO_2NR^5 , m is 1, 2 or 3; n is 0, 1 or 2; one of R^2 and R^3 represents a halogen, cyano, nitro, amino, hydroxyl, or a group selected from C_1 - C_6 -alkyl optionally substituted by at least one C_3 - C_6 -cycloalkyl, C_3 - C_8 -cycloalkyl, C_1 - C_6 -alkyloxy optionally substituted by at least one C_3 - C_6 -cycloalkyl, C_3 - C_8 -cycloalkyloxy, $S(O)_pC_1$ - C_6 -alkyl or $S(O)_qC_3$ - C_8 -cycloalkyl, each of these groups being optionally substituted by one or more fluorine atoms, and the other of R^2 and R^3 represents a hydrogen or halogen atom or a methyl group; p is 0, 1 or 2; q is 0, 1 or 2; R^4 represents $di(C_{1-2}alkyl)N(CH_2)_t$ where t is 0, 1 or 2 or imidazolyl, or R^4 represents a 3- to 9-membered saturated heterocyclic ring system containing one or two nitrogen atoms, the heterocyclic ring system being optionally substituted by one or more substituents independently selected from fluorine atoms, hydroxyl, C_1 - C_6 -alkyl, acetyl, hydroxy C_1 - C_6 -alkyl, $-NR^6R^7$, $-(CH_2)_rNR^6R^7$, $-CONR^6R^7$ and pyrimidinyl, or R^4 represents a 3- to 8-membered saturated carbocyclic ring system substituted by one or more substituents independently selected from $-NR^6R^7$, $-(CH_2)_rNR^6R^7$, $-CONR^6R^7$ the ring system being optionally further substituted by one or more substituents independently selected from fluorine atoms, hydroxyl and C_1 - C_6 -alkyl; r is 1, 2, 3, 4, 5 or 6; R^5 represents a hydrogen atom or a C_1 - C_6 -alkyl or C_3 - C_8 -cycloalkyl group; and R^6 and R^7 each independently represent a hydrogen atom or a

C₁-C₆-alkyl or C₃-C₈-cycloalkyl group, or R⁶ and R⁷ together with the nitrogen atom to which they are attached form a 3- to 8-membered saturated heterocyclic ring, provided that when R³ represents a cyano group, then X is other than a bond; or a pharmaceutically acceptable salt or solvate thereof.

A compound of general formula:



wherein X represents an oxygen or sulphur atom or a group NH, CH₂, CH₂CH₂ or OCH₂; Y represents a group CH₂ or C=O; R¹ represents a pyridyl or pyrimidinyl group; R² represents a phenyl, pyridyl or pyrimidinyl group, each of which may be optionally substituted by one or more substituents independently selected from a halogen atom or an amino, cyano, hydroxyl, nitro, C₁-C₆-alkyl, halo-C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, (di)C₁-C₆-alkylamino, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-alkylsulphinyl, C₁-C₆-alkylsulphonyl, -NR³SO₂R⁴ or -SO₂NR⁵R⁶ group, or a group -Z-(CH₂)_p-Z-(CH₂)_q-H wherein each Z independently represents a nitrogen or oxygen atom, p is an integer from 2 to 5 and q is 0 or an integer from 1 to 5; R³ and R⁴ each independently represent a hydrogen atom or a C₁-C₆-alkyl group; and R⁵ and R⁶ each independently represent a hydrogen atom or a C₁-C₆-alkyl group, or together with the nitrogen atom to which they are attached form a pyrrolidinyl or piperidinyl group; or a pharmaceutically acceptable salt or solvate thereof.

An additional embodiment of the invention provides a method for treating affective disorders by administering an agent modulating the activity of P2X7R, such as an antagonist of the P2X7R. The term "antagonist" means an agent or drug or a compound that opposes the physiological effects of another. Examples of P2X7R antagonists include, but are not restricted to, adamantane derivatives, isoquinolines and their derivatives, substituted phenyl compounds, piperidine derivatives, piperazine derivatives. P2X7R antagonists are described in the art and include the compounds found in Chen et al., Bioconjugate Chem. 13 (2002), 1100-1111;

WO 99/29660; WO 99/29661; WO 99/296896; WO 00/61569; WO 01/42194; WO 01/44170; WO 01/44213; WO 00/71529; WO 01/46200. P2X7R activity can also be modulated by RNA-based interference mechanisms and methods such as, but not limited to, small interference RNA (siRNA) molecules, and long double-stranded RNA (dsRNA).

In a further embodiment, the invention provides a method for treating affective disorders by administering an agent modulating the activity of P2X7R such as an agonist of the P2X7R. The term "agonist" means an agent or a compound that can interact with a receptor and initiate a physiological or a pharmacological response characteristic of that receptor. Examples of P2X7R agonist include but are not restricted to ATP, ATP-4, and BzATP (2- and 3-O-(4-benzoylbenzoyl)adenosine 5-triphosphate).

Dosage, pharmaceutical preparation and delivery of P2X7R modulating agent for use in accordance with the present invention may be formulated in conventional manner according to methods found in the art, using one or more physiological carriers or excipient, see, for example Ansel et al., "Pharmaceutical Dosage Forms and Drug Delivery Systems", 7th edition, Lippincott Williams & Wilkins Publishers, 1999. Thus, the P2X7R modulating agent and its physiologically acceptable salts and solvates may be formulated for administration by inhalation, insufflation (either through the mouth, or nose), oral, buccal, parenteral, or rectal administration.

For oral administration, the pharmaceutical composition of the P2X7 modulating agent may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutical acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone, hydroxypropyl methylcellulose), fillers (e.g., lactose, microcrystalline cellulose, calcium hydrogen phosphate), lubricants (e.g., magnesium stearate, talc, silica), disintegrants (e.g., potato starch, sodium starch glycolate), or wetting agents (e.g., sodium lauryl sulphate). Liquid preparations for oral administration may take the form of, for example, solutions, syrups, or suspensions, or may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparation

may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol, syrup, cellulose derivatives, hydrogenated edible fats), emulsifying agents (e.g., lecithin, acacia), non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol, fractionated vegetable oils), preservatives (e.g., methyl or propyl-p-hydroxycarbonates, soric acids). The preparations may also contain buffer salts, flavouring, coloring and sweetening agents as deemed appropriate. Preparations for oral administration may be suitably formulated to give controlled release of the agent modulating P2X7R activity.

For administration by inhalation, the agent modulating P2X7R activity for use according to the present invention is conveniently delivered in the form of an aerosol spray presentation from a pressurised pack or a nebulizer, with the use of a suitable propellant (e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In the case of a pressurised aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatine, for use in an inhaler or insufflator may be formulated containing a powder mix of the P2X7R activity modulating agent and a suitable powder base such as lactose or starch.

An agent modulating P2X7R activity may be formulated for parenteral administration by injection, for example, by bolus injection or continuous infusion. Site of injections include intra-venous, intra-peritoneal or sub-cutaneous. Formulations for injection may be presented in unit dosage form (e.g., in phial, in multi-dose container), and with an added preservative. The agent modulating P2X7R activity may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing, or dispersing agents. Alternatively, the agent may be in powder form for constitution with a suitable vehicle (e.g., sterile pyrogen-free water) before use.

An agent modulating P2X7R activity may, if desired, be presented in a pack, or dispenser device which may contain one or more unit dosage forms containing the said agent. The pack may for example comprise metal or plastic foil, such as blister

pack. The pack or dispenser device may be accompanied with instruction for administration.

In a more preferred embodiment the aforementioned methods or uses are envisaged to treat affective disorders selected from the group consisting of major depression, generalized anxiety disorder and bipolar disorder.

In a particularly preferred embodiment said major depression is selected from the group consisting of major depression, dysthymia, atypical depression, premenstrual dysphoric disorder and seasonal affective disorder.

In another particularly preferred embodiment said generalized anxiety disorder is selected from the group consisting of panic disorder, phobias, agoraphobia, social phobia, specific phobia, obsessive-compulsive disorder, post-traumatic stress disorder, separation anxiety disorder, mania, hypomania and cyclothymic disorder.

A still also particularly preferred embodiment is that said bipolar disorder is bipolar disorder type I or bipolar disorder type II.

Additionally, the present invention relates to a kit comprising the nucleic acid molecule, the vector, the host, the polypeptide, the antibody or the aptamer, the primer or pair of primers of the invention or the molecule as identified or characterized in a method herein below of the present invention.

Advantageously, the kit of the present invention further comprises, optionally (a) reaction buffer(s), storage solutions and/or remaining reagents or materials required for the conduct of scientific or diagnostic assays or the like. Furthermore, parts of the kit of the invention can be packaged individually in vials or bottles or in combination in containers or multicontainer units.

The kit of the present invention may be advantageously used, inter alia, for carrying out the method of producing a polypeptide of the invention, the method(s) of identification and/or characterization of molecules specifically interacting with P2X7R ATP-gated ion channels as described herein below and/or it could be employed in a variety of applications referred herein, e.g., as diagnostic kits, as research tools or

therapeutic tools. Additionally, the kit of the invention may contain means for detection suitable for scientific, medical and/or diagnostic purposes. The manufacture of the kits follows preferably standard procedures which are known to the person skilled in the art.

Furthermore, the present invention relates to a method for identifying compounds or mixtures of compounds which are capable of specifically interacting with a polypeptide of the present invention, comprising the steps of (a) contacting a polypeptide of the present invention with a candidate compound or mixture of compounds to be tested; and (b) determining whether said is capable of specifically interacting with said polypeptide. The polypeptide may be provided directly or by expression of a corresponding nucleic acid molecule or vector of the invention, e.g., in vitro or in a suitable host cell.

Additionally, the present invention relates to a method for the characterization of compounds which are capable of altering characteristics of the polypeptides of the present invention, comprising the steps of (a) contacting a polypeptide of the invention with said compound; and (b) determining whether the compound alters a characteristic of said polypeptide.

The term "altering characteristic of the polypeptide of the present invention" means that the functional characteristics to the polypeptides of the present invention in comparison to functional characteristics which they had before being contacted with the compounds identified by the above-described method: as described hereinabove are altered; i.e. changed.

Said identification and/or characterization of which are capable of interacting with or altering characteristics of the polypeptide of this invention, may be, inter alia, achieved by transfecting an appropriate host with a nucleic acid molecule of invention. Said hosts comprise, but are not limited to, HEK 293 cells or are injected into frog oocytes, preferably a *Xenopus* oocyte for functional expression (Goldin, *Methods Enzymol.* 207 (1992), 266). Expressed P2X7R ATP-gated channels can be examined using standard two-electrode voltage clamp techniques (Stuhmer,

Methods Enzymol. 207 (1992), 319; Kohler, Science 273 (1996), 1709). After expression of a P2X7R ATP-gated ion channel as defined herein, membrane currents may be deduced in the absence and/or presence of the molecule to be identified and/or characterized. Methods for the deduction of membrane currents are well known in the art and comprise, e.g., patch clamp methods as described in Hamill, Pfluger's Arch. 391 (1981), 85-100 or two-electrode voltage clamp in oocytes, as described in Methfessel, Pflügers Archive 407 (1986), 577-588. In accordance with the present invention the term "interacting with the polypeptides of the present invention" means that the polypeptides of the present invention interact directly and/or indirectly with compounds identified by the method described above. Furthermore, the present invention relates to a method of screening for molecules which are capable of interacting with the polypeptide of this invention, comprising the steps of (a) contacting a polypeptide of the invention with a molecule; and (b) measuring and/or detecting a response; and (c) comparing said response to a standard response as measured in the absence of said candidate molecule.

The present invention also relates to a method for identifying a compound which is capable of enhancing or reducing the expression of the P2X7R gene comprising the steps of contacting a cell which expresses the P2X7R gene from its natural promoter or a reporter gene driven by the P2X7R promoter and determining whether the expression of the gene is increased or reduced when compared to conditions in which the compound is not present.

Potential candidate molecules or candidate mixtures of molecules may be, inter alia, substances, compounds or compositions which are of chemical or biological origin, which are naturally occurring and/or which are synthetically, recombinantly and/or chemically produced or compounds or compositions described hereinabove. Thus, candidate molecules may be proteins, protein-fragments, peptides, amino acids and/or derivatives thereof or other compounds, such as ions, which bind to and/or interact with wild-type P2X7R ATP-gated ion channels. Such binding and/or interacting candidate compounds may be found employing, inter alia, yeast two-hybrid systems or modified yeast two-hybrid systems as described, for example in Fields, Nature 340 (1989), 245-246; Gyuris, Cell 75 (1993), 791-801; or Zervos, Cell 72 (1993), 223-232.

Furthermore, potential candidate molecules may be contacted with a cell, such as an oocyte or a HEK 293 cell, which expresses a polypeptide of the invention or with a membrane patch comprising a polypeptide of the invention and a corresponding response (inter alia, a dose-response response, a current-response, or single current channel response) may be measured in order to elucidate any effect said candidate molecule causes.

Within the scope of the present invention are also methods for identifying, characterizing and for screening of molecules which are capable of interacting with the P2X7R ATP-gated ion channels according to the invention which comprise so-called high-throughput screening methods and similar approaches which are known in the art (Spencer, Biotechnol. Bioeng. 61 (1998), 61-67; Oldenburg, Annu. Rev. Med. Chem. 33 (1998), 301-311) carried out using 96-well, 384-well, 1536-well (and other) commercially available plates. Further methods to be employed in accordance with the present invention comprise, but are not limited to, homogenous fluorescence readouts in high-throughput screenings (as described, inter alia, in Pope, Drug Discovery Today 4 (1999), 350-362). The method of the present invention for identification, characterization and/or screening of molecules capable of interacting with P2X7R ATP-gated ion channels can, inter alia, employ hosts as defined herein which express the polypeptide of the present invention. Cell-based assays, instrumentation for said assays and/or measurements are well-known in the art and described, inter alia, in Gonzalez, Drug Discovery Today 4 (1999), 431-439 or Ramm, Drug Discovery Today 4 (1999), 401-410. It is also envisaged that the high through put screens described herein are conducted by using, for example cRNA, i.e. synthetic RNA from a cDNA construct) that can be introduced in host cells, such as *Xenopus* oocytes using routine methods in the art. As an example, direct nucleic acid injection can be employed, such as the Eppendorf microinjection system (Micromnipulator 5171 and Transjector 5242). The injected/transformed cells can be analyzed for ion currents about 4 hours later using patch-clamp techniques which are commonly practiced in the art.

Additionally, the present invention relates to a method for the production of a pharmaceutical composition comprising the steps of a method of the invention for identifying, characterizing and/or screening of molecules which are capable of interacting with and/or altering the characteristics of a P2X7R ATP-gated ion channel of the invention and further comprising a step, wherein a derivative of said identified, characterized and/or screened molecule is generated. Such a derivative may be generated by, inter alia, peptidomimetics.

The invention furthermore relates to a method for the production of a pharmaceutical composition comprising the steps of a method of the invention for identifying, characterizing, screening and/or derivatizing of molecules which are capable of interacting with and/or altering the characteristics of a P2X7R ATP-gated ion channel and formulating the molecules identified, characterized, screened and/or derivatized in pharmaceutically acceptable form.

In a more preferred embodiment the present invention relates to a method wherein said molecule(s) comprise(s) (a) neuroprotective, (a) nootropic and/or (a) antiepileptic molecule(s).

Yet another embodiment of the invention is the use of a P2X7R polypeptide, in particular those according to the present invention, to identify biological, chemical, or pharmacological agents that can have an antidepressive effect. The term 'agent' refers to a chemical compound or composition capable of inducing a desired therapeutic or prophylactic effect when properly administered to a subject or cell. For example, the present invention allows the generation of cells expressing P2X7R for the identification and characterization of agents which modulate ionic influx and efflux. For example, HEK293 cells, or other cell lines (e.g., HCN-1A, HCN-2, HIT-T15, RIN-m5F, betaTC3, PC12, HT22, SH-SY5Y, Neuro2A or CA77), can be stably transfected with cDNA encoding the human P2X7R and plated in 12, 96 and 384 well plates. Said cells are cultured in appropriate medium. Examples of such medium are well known in the art, see, for example Freshney, "Culture of Animal Cells: A Manual of Basic Technique, 4th edition, Wiley-Liss Publishing, 2000.

Said cells can then be pre-incubated with said agents for 15 min prior to stimulation with 3 mM ATP for 10 minutes. Reactions are then terminated by rapid aspiration of the extracellular medium in each well. The cells in each well are subsequently extracted overnight with 1 ml 10% HNO₃. Potassium (K⁺) content in the extracts can be determined by atomic absorbance spectrophotometry. Agent function is then measured by the percent inhibition or stimulation of the K⁺ release triggered by 3 mM ATP and compared to K⁺ release in the absence of the agents. P2X7R activity can also be monitored according to the movement of calcium (Ca²⁺; see Denyer et al., Drug Discov. Today 7 (1998), 323-332; González et al., Drug Discov. Today 9 (1999), 431-439; Helmchen and Waters, Eur. J. Pharmacol. 447 (2002), 119-129). Agents can also be verified in the absence of ATP.

P2X7R activity can also be monitored according to secretion of neurotransmitters such as glutamate and GABA. Neurotransmitter levels in treated cells can be quantified by suitable methods, e.g., Enzyme Linked Immunoabsorbent Assay (ELISA), Radio Immuno Assay (RIA), High Performance Liquid Chromatography (HPLC). Using these methods, a large number of compounds can be screened for increase in neurotransmitter (for example, glutamate) secretion. The release of glutamate can be measured for example by Fluorometric glutamate release assays (e.g., Amplex Red Glutamic Acid/Glutamate Oxidase Assay Kit, Molecular Probes) or High-Throughput ElectroPhysiology.

In another embodiment, cells are transfected with nucleic acid constructs encoding a reporter gene regulated by the P2X7R promoter (see above), an increase or decrease in the expression of the reporter gene in response to biological or pharmaceutical agents can be analyzed using methods that detect levels or status of protein or mRNA present in the corresponding cell or detect biological activities of the reporter gene. Suitable reporter molecules or labels, which may be used, include radionucleotides, enzymes, fluorescent, chemiluminescent or chromogenic agents as well as substrates, co-factors, inhibitors, magnetic particles, and the like. Designing such drug screening assays are well known in the art; see Harvey ed., 'Advances in drug discovery techniques', John Wiley and Sons, 1998; Vogel and Vogel eds., 'Drug discovery and evaluation: Pharmaceutical assays', Springer-Verlag Berlin, 1997). For

example, drug screening in animal models, in vitro tests using animal cells, or in vivo tests involving toxicology tests in animals. An in vitro model can be used for screening libraries of compounds in any of a variety of drug screening techniques.

Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 Daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise carbocyclic or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups.

Candidate agents are also found among biomolecules including peptides, amino acids, saccharides, fatty acids, steroids, purines, pyrimidines, nucleic acids and derivatives, structural analogs or combinations thereof. Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs.

Another technique for drug screening, which may be used, provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO 84/03564. In this method, as applied to the proteins of the invention large numbers of different small test

compounds, e.g. aptamers, peptides, low-molecular weight compounds etc., are provided or synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with the proteins or fragments thereof, and washed. Bound proteins are then detected by methods well known in the art. Purified proteins can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support. In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding the protein specifically compete with a test compound for binding the protein. In this manner, the antibodies can be used to detect the presence of any peptide, which shares one or more antigenic determinants with the protein.

The present invention further particularly provides a method, wherein the pharmaceutical composition to be produced further comprises neuroprotective substances, nootrophic substances, brilliant blue, piperidine or derivatives thereof, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate or 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate. It is also envisaged that the pharmaceutical compositions to be produced further comprise antidepressants such as fluoxetine, paroxetine, sertraline, fluvoxamine, mirtazapine, reboxetine, nefazodone or lithium carbonate.

In a preferred embodiment of the present invention, the compounds of the aforementioned methods comprise antagonist(s), partial antagonist(s), partial agonist(s) and/or agonist(s) for an altered ATP-gated ion channel P2X7R.

In accordance with the present invention, the term "antagonist" denotes molecules/substances, which are capable of inhibiting and/or reducing an agonistic effect. The term "antagonist" comprises competitive, non-competitive, functional and chemical antagonists as described, inter alia, in Mutschler, "Arzneimittelwirkungen" (1986), Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart, Germany. The term "partial antagonist" in accordance with the present invention means a molecule/substance that is capable of incompletely blocking the action of agonists through, inter alia, a non-competitive mechanism.

In the context of the present invention, an antagonist is preferably a drug that does not provoke a response itself, but blocks agonist-mediated responses. It is a chemical entity that opposes the receptor-associated responses normally induced by another bioactive agent. For the P2X7R, the antagonists have an IC50 between 10 nanomolar and 300 micromolar.

As "agonist", in accordance with this invention, molecules/substances are denoted which have an affinity as well as an intrinsic activity. Mostly, said intrinsic activity (α) is defined as being proportional to the quotient of the effect, triggered by said agonist (EA) and the effect which can be maximally obtained in a given biological system (Emax): therefore, the intrinsic activity can be defined as

$$\alpha \sim \frac{E_A}{E_{max}}$$

The highest relative intrinsic activity results from EA/Emax=1. Agonists with an intrinsic activity of 1 are full agonists, whereas substances/molecules with an intrinsic activity of >0 and <1 are partial agonists. Partial agonists show a dualistic effect, i.e. they comprise agonistic as well as antagonistic effects.

Preferably, in the context of the present invention, an agonist (or full agonist) is an endogenous substance or a drug that can interact with a receptor and initiate a maximal or complete physiological or a pharmacological response characteristic of that receptor. ATP, the natural ligand for the P2X7R, is an agonist with an EC50 of 300 micromolar while the synthetic P2X7R agonist Bz-ATP has an EC50 of 8 micromolar. Thus, agonists of P2X7R have an EC50 equal or below 300 micromolar. The EC50 is defined as the concentration of agonist that provokes a response half way between the baseline response and maximum response on a dose response curve where the X-axis plots concentration of an agonist and the Y-axis plots ion current. An inverse agonist (also called negative antagonist) is a drug which acts at the same receptor as that of an agonist, yet produces an opposite effect. A partial agonist is an endogenous substance or a drug that also provokes physiological or a pharmacological response but, the maximum response is less than the maximum

response to a full agonist, regardless of the amount of drug applied. In the case of P2X7R, partial agonists have EC50s higher than 300 micromolar.

The person skilled in the art can, therefore, easily employ the compounds and the methods of this invention in order to elucidate the agonistic and/or antagonistic effects and/or characteristics of a compound/molecule/substance to be identified and/or characterized in accordance with any of the above described methods. Preferably, an identified antagonist of the ATP-gated ion channel P2X7R comprising the mutation(s) and/or deletion(s) described hereinabove may be useful to reestablish the properties normally shown by wild-type P2X7R ATP-gated ion channels. An identified agonist of the ATP-gated ion channel P2X7R comprising the mutation(s) and/or deletion(s) described hereinabove may be useful to reestablish the lost functionality of the P2X7R ATP-gated ion channel.

The Figures show:

Figure 1a. Genomic map of the region on the human chromosome 12 associated to bipolar affective disorder. Genes found between markers NBG11 and NBG2 are depicted.

Figure 1b. Graphic illustrating the multipoint analysis using ASPEX on independent sib-pairs.

Figure 1c. Graphic illustrating the multipoint analysis using ASPEX on all sib-pairs

Figure 1d. Graphic illustrating the ASPEX sib_phase by considering only independent sib-pairs

Figure 1e. Graphic illustrating the ASPEX sib_phase by considering all sib-pairs

Figure 2. RT-PCR analysis of the complete coding sequence of P2X7R in different tissues

Figure 3. P2X7R expression in the olfactory bulb, hypothalamus and ependymal cells in the brain of a stress-free mouse. Magnification 100X.

Figure 4. P2X7R expression in the hippocampus/dentate gyrus and subcommisural organ in the brain of a stress-free mouse. Magnification 100X.

Figure 5. Floating behaviour in the forced swim test. Passive stress coping behaviour decreased after long-term treatment with the antidepressant paroxetine (Par28: treated with paroxetine for 28 days, per os). Basal n=8; vehicle n=8; Par28 n=8.

Figure 6. Comparative analysis of P2X7R expression in the olfactory bulb of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 100X.

Figure 7. Comparative analysis of P2X7R expression in the hypothalamus of stress-free, treated-treated and antidepressant-treated mice. Magnification 100X.

Figure 8. Comparative analysis of P2X7R expression in ependymal cells of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 100X.

Figure 9. Comparative analysis of P2X7R expression in the hippocampus of stress-free, vehicle-treated and antidepressant-treated mice. Magnification 25X.

Figure 10. P2X7R expression in the hippocampus of a vehicle treated mouse. Magnification 25X.

Figure 11. P2X7R expression in the hippocampus of a mouse treated with the antidepressant paroxetine. Magnification 25X.

Figure 12. Detailed expression of P2X7R in the dentate gyrus of a mouse treated with the antidepressant paroxetine. Magnification 400x.

Figure 13. Comparative analysis of P2X7R expression and apoptotic cells in the hippocampus of a mouse treated with the antidepressant paroxetine. Magnification 100X.

Figure 14. Floating behaviour in the forced swim test. Passive stress coping behaviour increased after acute intrahippocampal (bilateral, dentate gyrus) of siRNA targeting P2X7R. Vehicle n=7; control RNA n=10; P2X7R siRNA n=9.

Figure 15. Comparative analysis of P2X7R expression in the hippocampus of mice treated with vehicle, control RNA and of siRNA targeting P2X7R. Magnification 100X upper row, 25X lower row.

Figure 16a, b, c, d, e. Three splicing variants caused by polymorphisms in the introns of P2X7R.

A better understanding of the present invention and of its many advantages will be had from the following examples, offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

EXAMPLE 1

Linkage analysis of Bipolar Affective Disorder in a homogeneous human population

41 families of different sizes containing a total of 485 sampled individuals from the region of Saguenay/Lac St-Jean were used in the linkage analysis. Individuals were distributed according to their diagnoses as follows: 105 individuals afflicted with Bipolar Disorder type I (BPI) or schizoaffective disorder bipolar type; 42 individuals diagnosed with Bipolar Disorder type II (BPII); 54 individuals with recurrent major depression; and 57 individuals with single episode major depression. The remaining 227 individual unaffected and normal. For the purpose of the calculation, the following classification was used: individuals diagnosed with either BPI, schizoaffective disorder, bipolar type, BPII and recurrent major depression were considered as affected (n=201); individuals with a single major depression episode were scored as unknown phenotype (n=57); and all others diagnoses as unaffected (n=227).

Blood samples from each individual were collected in 10-ml K3 EDTA Vacutainer tube (Becton-Dickinson) and genomic DNA was isolated by Puregene DNA Isolation kit (Gentra Systems). Blood was poured into 50 ml conical tube and diluted with four volume of Red Blood Cell Lysis Solution. After an incubation of 10 minutes at room temperature, the tube was centrifuged for 10 minutes at 2,000 g and supernatant was removed leaving behind cell pellet and 200-400 µl of the residual liquid. Cells were resuspended by vortexing the tube and 9 ml of Cell Lysis Solution were added with up and down pipetting. 40 µl of RNase A Solution (20 mg/ml) were added and the sample was mixed by inverting the tube several times. Sample was incubated at 37°C for 15 minutes and cooled to room temperature. 3 ml of Protein Precipitation Solution were added to cell lysate. Tube was vigorously vortexed for 30 seconds and

centrifuged at 2,000 g for 10 minutes. Supernatant was poured into a new tube containing 9 ml of 100% isopropanol. Sample was mixed by inverting gently several times. Tube was centrifuged at 2,000 g for 5 minutes. The DNA white pellet was washed with 10 ml 70% ethanol and the tube was centrifuged at 2,000 g for 3 minutes. Ethanol was poured off and pellet allowed to partially air dry. DNA was solubilized in 500 µl of DNA Hydration Solution. Final concentration was adjusted to 300-400 µg/ml.

A fluorescent-based method was used for the genotyping of microsatellite markers. Briefly, the region encompassing each repeated sequence was amplified by PCR using an unlabeled primer and a fluorescent-labeled primer (Applied biosystems inc, CA, USA). The marker-associated dyes and the corresponding PCR product length are listed in table 2. The PCR reaction was performed using 10ng of DNA sample, 0,2 unit of Taq platinum DNA polymerase (Invitrogene, CA, USA), 20 mM Tris-Cl (pH 8.4), 50 mM KCl, 1.5mM MgCl₂, 100µM of dNTP, and 1,5 µM of each primer in a final volume of 7 µl. The samples were incubated at 95°C for 3 minutes to activate the Taq platinum DNA polymerase, then 10 cycles of PCR amplification were performed as follows: 95°C for 15 seconds; 58°C for 15 seconds; 72°C for 30 seconds; after that 15 cycles were performed as follow: 89°C for 15 seconds; 58°C for 15 seconds; 72°C for 30 seconds. Finally, the samples were incubated at 72°C for 30 min. Following the PCR amplification samples were pooled according to their dye-labeled primer and their PCR product length (pool of four samples). Pooled sample were separated on an ABI 3100 DNA analyzer (Applied Biosystems inc, CA, USA). The resulting data were analysed using Genemapper2 (Applied Biosystems inc, CA, USA), and compiled in a 4D database (ACIUS) designed in a Macintosh environment as previously described (Morissette et al., Am. J. Med. Genet. (Neuropsychiatr. Genet.) 88 (1999), 567-587)

Markers used in the following linkage analysis are shown in table 2. Recombination fraction (q) between successive markers was computed according to the analyzed families.

Table 2. Genomic markers used for the linkage analysis

| Locus | Associated dye | Allele length (bp) | Distance (q) | Cumulative distance (cM) | Heterozygosity (%) |
|----------|----------------|--------------------|--------------|--------------------------|--------------------|
| D12S1619 | VIC | 170-210 | 0.0135 | 0.00 | 74.5 |
| NBG11 | VIC | 204-218 | 0.006 | 1.37 | 65.5 |
| D12S1666 | FAM | 241-281 | 0.001 | 1.97 | 66.9 |
| NBG5 | VIC | 253-261 | 0.001 | 2.07 | 38.3 |
| D12S1721 | VIC | 263-299 | 0.005 | 2.17 | 72.1 |
| NBG8 | VIC | 166-188 | 0.011 | 2.67 | 73.3 |
| NBG6 | NED | 182-218 | 0.0115 | 3.79 | 73.9 |
| NBG9 | VIC | 156-180 | 0.0035 | 4.95 | 68.9 |
| NBG10 | FAM | 174-186 | 0.001 | 5.30 | 49.7 |
| NBG12 | NED | 165-207 | 0.009 | 5.40 | 64.2 |
| NBG4 | NED | 171-199 | 0.001 | 6.31 | 66.4 |
| NBG3 | VIC | 182-206 | 0.006 | 6.41 | 64.8 |
| NBG2 | VIC | 171-199 | | 7.01 | 54.2 |

-Haldane's map function was used for cumulative distance in cMorgans.

For bipoint parametric analysis, MOD score analysis were used where parametric LOD score were maximized over genetic models.

The following results were obtained under MOD score analysis for recessive models.

Table 3. MOD score analysis for recessive models

| Locus | Distance (q) | Cumulative distance (cM) | LOD score (q_{\max}) |
|----------|--------------|--------------------------|--------------------------|
| D12S1619 | 0.0135 | 0.00 | 3.46 (0.10) |
| NBG11 | 0.006 | 1.37 | 4.06 (0.04) |
| D12S1666 | 0.001 | 1.97 | 1.22 (0.14) |
| NBG5 | 0.001 | 2.07 | 0.66 (0.16) |
| D12S1721 | 0.005 | 2.17 | 2.82 (0.10) |
| NBG8 | 0.011 | 2.67 | 1.51 (0.00) |
| NBG6 | 0.0115 | 3.79 | 4.77 (0.06) |
| NBG9 | 0.0035 | 4.95 | 0.75 (0.22) |
| NBG10 | 0.001 | 5.30 | 0.74 (0.00) |
| NBG12 | 0.009 | 5.40 | 1.41 (0.16) |
| NBG4 | 0.001 | 6.31 | 3.56 (0.08) |
| NBG3 | 0.006 | 6.41 | 3.96 (0.08) |
| NBG2 | | 7.01 | 2.59 (0.10) |

Model-free LOD score studies using ANALYZE, sib_phase from the ASPEX V1.85 package (David Hinds and Neil Risch 1999; <ftp://lahmed.stanford.edu/pub/aspeX>, see also <http://watson.hgen.pitt.edu/docs/usage.html>) and SIMWALK2 (Sobel and Lange, Am J Hum Genet 58 (1996), 1323-1337) were performed to analyze the allele sharing among affected sib-pairs. The ANALYZE program weights sibships according to their size. The ASPEX sib_phase program uses allele frequencies to reconstruct missing information, and is tailored for data sets where parents are missing, but additional typed children may be used to reconstruct and phase the parents. SimWalk2 is a statistical genetics computer application for haplotype, parametric linkage, non-parametric linkage (NPL), identity by descent (IBD) and mistyping analyses on any size of pedigree. SimWalk2 uses Markov chain Monte Carlo (MCMC) and simulated annealing algorithms to perform these multipoint analyses.

ASPEX sib_phase was used with two computational strategies: First, by using strictly independent sib pairs; secondly, by using all affected sib-pair combinations. ASPEX was performed for bi-point and multipoint calculations.

The bi-point results observed with ANALYZE and ASPEX are shown in Table 4.

Table 4. Bi-point results observed with ANALYZE and ASPEX

| Locus | Distance (q) | Cumulative distance (cM) | Sib-pair from ANALYZE LOD score | sib_phase LOD score indep. sib-pairs | sib_phase LOD score all sib-pairs |
|----------|--------------|--------------------------|---------------------------------|--------------------------------------|-----------------------------------|
| D12S1619 | 0.0135 | 0.00 | 2.31 | 2.55 | 3.14 |
| NBG11 | 0.006 | 1.37 | 2.83 | 2.72 | 3.27 |
| D12S1666 | 0.001 | 1.97 | 1.01 | 2.52 | 3.14 |
| NBG5 | 0.001 | 2.07 | 0.50 | 2.52 | 3.13 |
| D12S1721 | 0.005 | 2.17 | 1.57 | 2.51 | 3.12 |
| NBG8 | 0.011 | 2.67 | 0.51 | 2.24 | 2.75 |
| NBG6 | 0.0115 | 3.79 | 2.55 | 2.11 | 2.64 |
| NBG9 | 0.0035 | 4.95 | 0.49 | 1.65 | 1.97 |
| NBG10 | 0.001 | 5.30 | 0.77 | 1.45 | 2.10 |
| NBG12 | 0.009 | 5.40 | 0.47 | 1.44 | 2.17 |
| NBG4 | 0.001 | 6.31 | 1.21 | 1.29 | 3.07 |
| NBG3 | 0.006 | 6.41 | 1.84 | 1.29 | 3.07 |
| NBG2 | | 7.01 | 1.24 | 1.22 | 3.00 |

SIMWALK2 computed four different statistics based on descent trees. These statistics measure the degree of clustering among the marker alleles descending from the founders.

Statistic A is the number of different founder-alleles contributing alleles to the affected it is most powerful at detecting linkage to a recessive trait. Statistic B is the maximum number of alleles among the affected descended from any one founder-allele and most powerful at detecting linkage to a dominant trait. Statistic C is the 'entropy' of the marker alleles among the affected. Statistic D is the extent of allele sharing among all affected pairs as measured by their IBD kinship coefficient. Statistics C and D are more general statistics indicating whether a few founder-alleles are overly represented among the affected.

Table 5 shows the results observed with SIMWALK2. The authors signal that p-values should be generally conservative. They are expressed as $-\text{Log}(\text{p-values})$. For correspondence purpose, $-\text{Log}(0.05)=1.30$, $-\text{Log}(0.01)=2$, $-\text{Log}(0.001)=3$ etc.

Table 5. SIMWALK2 analysis

| Locus | Distance (q) | Cumul. Distance (cM) | STAT(A) -Log(p-value) | STAT(B) -Log(p-value) | STAT(C) -Log(p-value) | STAT(D) -Log(p-value) |
|----------|--------------|----------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| D12S1619 | 0.0135 | 0.00 | 1.4550 | 0.4103 | 1.1306 | 1.1310 |
| NBG11 | 0.006 | 1.37 | 2.0157 | 1.4375 | 1.5955 | 1.9845 |
| D12S1666 | 0.001 | 1.97 | 2.0236 | 0.9765 | 1.4727 | 1.4614 |
| NBG5 | 0.001 | 2.07 | 1.7596 | 0.8558 | 1.3866 | 1.3602 |
| D12S1721 | 0.005 | 2.17 | 1.6628 | 1.1692 | 1.4235 | 1.6384 |
| NBG8 | 0.011 | 2.67 | 1.5374 | 0.6940 | 1.0623 | 1.1552 |
| NBG6 | 0.0115 | 3.79 | 1.5896 | 0.4452 | 1.0935 | 1.1786 |
| NBG9 | 0.0035 | 4.95 | 1.2677 | 0.3815 | 0.8412 | 0.9133 |
| NBG10 | 0.001 | 5.30 | 1.1117 | 0.3642 | 0.6987 | 0.7554 |
| NBG12 | 0.009 | 5.40 | 1.0809 | 0.3485 | 0.6694 | 0.7179 |
| NBG4 | 0.001 | 6.31 | 1.1024 | 0.4148 | 0.6368 | 0.8544 |
| NBG3 | 0.006 | 6.41 | 1.1040 | 0.4146 | 0.6373 | 0.8559 |
| NBG2 | | 7.01 | 1.0963 | 0.5380 | 0.6587 | 0.9356 |

Multipoint result observed with ASPEX when only independent sib-pairs were used (Figure 1b). The maximum LOD score value was observed at NBG11.

Multipoint result observed with ASPEX when all sib-pairs were considered (Figure 1c). The maximum LOD score value was observed at NBG11 but a second peak appeared at NBG4 and NBG3.

Multipoint and bi-point LOD score values computed by ASPEX were similar. The second peak, observed when all sib-pairs are used, may be explained by the presence of a recombinant affected individual, with many affected sibs, sharing the chromosomal region telomeric to NBG12. This kind of individuals has a large impact on LOD score values when all sib-pairs are used instead of one sib-pair. This situation was observed in two sibships.

Strata analysis was subsequently performed. Although HOMOG did not detect evidence for heterogeneity, a homogeneity test was constructed based on allele sharing found in selected chromosomal regions. Only 20 of the 41 families were used for this analysis since the others were not genotyped in all these regions. For each marker within the selected regions, the proportion of alleles shared IBD by affected sib-pairs was estimated with ASPEX (sib_phase). For each region retained, the proportion of shared alleles was used as variable for a Principal Component Analysis

and the first principal component as an index of linkage. Correlation analysis was done on these indexes to detect heterogeneity (correlation < 0) or epistasis (correlation > 0). Fisher algorithm was used to classify into two groups of families as linked or unlinked to a particular locus. A negative correlation was observed between the chromosome 12 region and the chromosome 15 area ($r=-0.51$; $p=0.023$). Cluster analysis suggested that 11 families out of 20 were linked to chromosome 12. This sub-sample was called the strata.

This strata included 11 families (266 sampled individuals) that include 52 BPI or schizoaffective disorder, bipolar type, 20 BP II and 28 recurrent major depression

The following MOD score values illustrated in Table 6 were obtained under recessive models.

Table 6. MOD scores under recessive models

| Locus | Distance (q) | Cumulative distance (cM) | LOD score (q_{max}) |
|----------|--------------|--------------------------|-------------------------|
| D12S1619 | 0.0135 | 0.00 | 4.03 (0.08) |
| NBG11 | 0.006 | 1.37 | 4.98 (0.00) |
| D12S1666 | 0.001 | 1.97 | 1.49 (0.12) |
| NBG5 | 0.001 | 2.07 | 0.79 (0.14) |
| D12S1721 | 0.005 | 2.17 | 4.23 (0.06) |
| NBG8 | 0.011 | 2.67 | 2.79 (0.00) |
| NBG6 | 0.0115 | 3.79 | 5.06 (0.06) |
| NBG9 | 0.0035 | 4.95 | 1.57 (0.14) |
| NBG10 | 0.001 | 5.30 | 1.73 (0.00) |
| NBG12 | 0.009 | 5.40 | 1.65 (0.12) |
| NBG4 | 0.001 | 6.31 | 4.60 (0.08) |
| NBG3 | 0.006 | 6.41 | 4.84 (0.06) |
| NBG2 | | 7.01 | 2.80 (0.06) |

Model-free LOD score results obtained with ANALYZE and ASPEX applied to the strata are shown in Table 7.

Table 7. Model-free LOD score obtained with ANALYZE and ASPEX

| Locus | Distance (q) | Cumulative distance (cM) | ANALYZE LOD score | sib_phase LOD score independent sib-pairs | sib_phase LOD score all sib-pairs |
|----------|--------------|--------------------------|-------------------|---|-----------------------------------|
| D12S1619 | 0.0135 | 0.00 | 4.54 | 5.29 | 7.65 |
| NBG11 | 0.006 | 1.37 | 4.29 | 5.34 | 7.70 |
| D12S1666 | 0.001 | 1.97 | 2.77 | 5.36 | 7.74 |
| NBG5 | 0.001 | 2.07 | 0.67 | 5.36 | 7.74 |
| D12S1721 | 0.005 | 2.17 | 4.48 | 5.35 | 7.74 |
| NBG8 | 0.011 | 2.67 | 2.97 | 4.87 | 7.00 |
| NBG6 | 0.0115 | 3.79 | 4.05 | 4.59 | 6.76 |
| NBG9 | 0.0035 | 4.95 | 2.03 | 3.72 | 5.41 |
| NBG10 | 0.001 | 5.30 | 2.00 | 3.42 | 5.89 |
| NBG12 | 0.009 | 5.40 | 0.89 | 3.44 | 6.11 |
| NBG4 | 0.001 | 6.31 | 2.84 | 3.71 | 9.00 |
| NBG3 | 0.006 | 6.41 | 3.89 | 3.71 | 9.01 |
| NBG2 | | 7.01 | 1.91 | 3.52 | 8.73 |

Model-free results observed with SIMWALK2 are illustrated in Table 8.

Table 8. Model-free LOD score obtained with SIMWALK2

| Locus | Distance (q) | Cumulative Distance (cM) | STAT(A) -Log(p-value) | STAT(B) -Log(p-value) | STAT(C) -Log(p-value) | STAT(D) -Log(p-value) |
|----------|--------------|--------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| D12S1619 | 0.0135 | 0.00 | 2,5963 | 0,9565 | 3,2156 | 2,3584 |
| NBG11 | 0.006 | 1,37 | 3,0698 | 1,7400 | 3,7747 | 3,0103 |
| D12S1666 | 0.001 | 1,97 | 2,9340 | 1,6546 | 3,5812 | 2,7223 |
| NBG5 | 0.001 | 2,07 | 2,9781 | 1,2722 | 3,6505 | 2,7846 |
| D12S1721 | 0.005 | 2,17 | 2,9680 | 1,2630 | 3,6844 | 2,7752 |
| NBG8 | 0.011 | 2,67 | 3,0954 | 1,0804 | 3,4399 | 2,5654 |
| NBG6 | 0.0115 | 3,79 | 3,1632 | 1,0672 | 3,2670 | 2,5956 |
| NBG9 | 0.0035 | 4,95 | 2,2106 | 1,0137 | 2,7765 | 2,4456 |
| NBG10 | 0.001 | 5,30 | 2,5513 | 1,0251 | 2,7625 | 2,1914 |
| NBG12 | 0.009 | 5,40 | 2,4893 | 0,9868 | 2,6841 | 2,0920 |
| NBG4 | 0.001 | 6,31 | 2,9028 | 1,1312 | 3,4063 | 2,8156 |
| NBG3 | 0.006 | 6,41 | 2,9070 | 1,1326 | 3,4637 | 2,8300 |
| NBG2 | | 7,01 | 2,8430 | 1,1108 | 3,3135 | 2,7978 |

Multipoint results on the strata with ASPEX sib_phase by considering only independent sib-pairs (Figure 1b) or all sib-pairs (Figure 1c) are shown in Figures 1d and 1e. As previously reported a second peak appeared when all sib-pairs were observed.

A confidence interval was calculated. GENEFINDER (Liang et al., Am. J. Hum. Genet. 66 (2000), 1631-1641) was used to estimate the location of the susceptibility gene (say t). The method is based on the IBD (Identity by Descent) sharing of affected sib-pairs for multiple markers. For the purpose of our analysis, pedigrees were divided into sibship. 56 nuclear families and 183 sib-pairs were used. Liang KY, Huang CY, Beaty TH (2000) A unified sampling approach for multipoint analysis of qualitative and quantitative traits in sib pairs. Am J Hum Genet 66:1631-1641

The GENEFINDER results points to localization of a susceptibility gene for affective disorders at 3.19 ± 0.446 cM telomeric to the marker D12S1721 (D12S1721 is approximately located at 136.82 cM on the sex-averaged Marshfield chromosome 12 map).

95% C.I.: [2.32, 4.06];

99% C. I.: [2.03, 4.35]

99.9% C. I.: [1.71, 4.67]

From the strata, 24 nuclei, and 107 sib-pairs were obtained, and the location of the susceptibility gene was estimated at 3.07 ± 0.57 (see map above). The following confidence interval (C.I.) was obtained:

95% C.I.: [1.95, 4.19];

99% C. I.: [1.59, 4.55]

99.9% C. I.: [1.18, 4.96]

An association study using the NBG microsatellite markers was done with CLUMP (Sham & Curtis, Ann. Hum. Genet. 59 (1995), 97-105). Samples were distributed as follow: 83 male/case; 124 female/case; 95 male/control; and 101 female/control. One thousand simulations were used to estimate p-values. The observed results are summarized in Table 9.

Table 9. Association study using the NBG microsatellite

| Locus | Sample | | T1statistic (p-value) | T2statistic (p-value) | T3statistic (p-value) | T4statistic (p-value) |
|--------------|---------------|----------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| | Case | Control | | | | |
| NBG11 | 204 | 129 | 0.226 | 0.562 | 0.410 | 0.421 |
| NBG5 | 206 | 194 | 0.972 | 0.980 | 0.948 | 0.971 |
| NBG8 | 206 | 194 | 0.983 | 1.000 | 0.994 | 0.978 |
| NBG6 | 206 | 194 | 0.147 | 0.074 | 0.759 | 0.485 |
| NBG9 | 206 | 190 | 0.512 | 0.940 | 0.786 | 0.583 |
| NBG10 | 206 | 190 | 0.594 | 0.480 | 0.403 | 0.709 |
| NBG12 | 206 | 190 | 0.002 | 0.019 | 0.003 | 0.117 |

T1 statistic is the usual chi-squared statistic on the raw contingency table

T2 statistic is the usual chi-squared statistic apply on he contingency table obtained after collapsing columns with small expected values together

T3 statistic is the largest chi-squared statistic got by comparing one column of the original table against the total of the others columns

T4 statistic is the largest chi-squared statistic got by comparing any combination of alleles against the rest.

Only the NBG12 marker gave significant association at the 1% level. For the others markers, there was no single alleles that seems to be associated with bipolar disorder. It seems that no founder-alleles was overly represented among the affected. There is no significant result for association of genotypes with the NBG markers.

In conclusion, the parametric and model-free multipoint results suggest to investigate genes located between D12S1619 and D12S1666. Moreover, according to GENEFINDER results, genes situated centromeric to NBG9 should be considered for association and linkage disequilibrium analysis.

EXAMPLE 2

Physical mapping and Mutation analysis of chromosome 12 associating the P2X7R to Bipolar Affective Disorders

The most conservative prediction for the disease-associated region is included between markers NBG11 and NBG2 (see Figure 1a). This region was delimited according to linkage and association analysis described in Example 1, using genethon markers and NBG markers. The approximate length of this region is 5,2 Mb. Two major gaps (between FLJ10701 and FLJ32372, and between FLJ1466 and

MONDOA) were included in this region. At least 73 genes were listed in this area, where 48 are known genes and 25 are unknown but associated to mRNA and/or EST clusters based on the last genome assembly available at UCSC (November 2002). Predicted genes were not listed. However, the estimation of CI 99% (confidence interval) using GENEFINDER has limited the most interesting region between markers D12S1666 and NBG9. This genomic region covers 1,6 Mb and includes at least 28 genes, and has no major gap. Thus, the term fBAD (familial Bipolar Affective Disorders) region to describe the genomic segment between D12S1666 and NBG9. Genes found within this region include CaMKK2, CABP, P2X7, P2X4, PIN, PLA2, G1B, CIT, PXN, Rab35, and APC5. However, given the present art, it would not have been obvious to an ordinary person skilled in the art to select P2X7R as the gene associated with affective diseases. Other genes from the ones listed above would be obvious.

For example, the CaMKK2 gene (also known as Ca^{2+} /Calmodulin-dependent protein kinase kinase beta, or CaMKKb) is a serine/threonine protein kinase involved in Ca^{2+} dependent signalling pathways. CaMKK2 can activate in vitro the downstream kinases CaMKIV and CaMKI, which modulate gene transcription through phosphorylation of transcription factors (e.g., CREB, SRF, MEF2; Corcoran and Means, J. Biol. Chem. 276 (2001), 2975-2978; Soderling, Trends Biochem. Sci. 24 (1999), 232-236). Its role in the Ca^{2+} cascade is not critical. Some studies suggest that CaMKs could be activated without the CaMKKs phosphorylation (Matsushita and Nairn, J. Biol. Chem. 274 (1998), 10086-10093). However, CaMKK phosphorylation step would contribute to amplification of the Ca^{2+} signal since CaMKK is more sensitive to activation by Ca^{2+} /Calmodulin, therefore CaMKK would be an important mediator when the levels of intracellular Ca^{2+} are low (Anderson et al., J. Biol. Chem. 273 (1998), 31880-31889).

CaMKK2 is an obvious target for depression since prior art suggest that cAMP-dependent signaling pathways (mediated by PKA activation) is be affected in brain from patients with Bipolar Affective Disorders (Field et al., J. Neurochem. 73 (1997), 1704-1710; Rahman et al., J. Neurochem. 68 (1997), 297-304; Takahashi et al., J. Neurosci. 19 (1999), 610-618). According to a study using lymphoblastic cell lines,

Bipolar disorder could be related to elevated intracellular calcium levels (Yoon et al., *Mol. Psychiatry* 6 (2001), 678-683). Moreover, some groups found relations between antidepressant drugs and CaMK activation (Budziszewska et al., *Br. J. Pharmacol.* 130 (2000), 1385-1393; Consogno et al., *Neuropsychopharmacology* 24 (2001), 21-30; Mori et al., *Neuropharmacology* 40 (2001), 448-456; Zanotti et al., *Neuropharmacology* 37 (1998), 1081-1089). Furthermore, inhibition of CaMKK by PKA-mediated phosphorylation suggests a close relationship between both pathways (Matsushita et al., *J. Biol. Chem.* 273 (1999), 21473-21481). These observations would suggest to a person skilled in the art that CaMKK2 is the gene responsible for bipolar affective disease.

Another obvious candidate for affective disorders would have been the CABP1 gene which generates four neuronal Ca^{2+} -binding protein by alternative usage of the 9 coding exons, which are L-CABP, S-CABP, calbrain, and caldendrin (Haeseleer et al., *J. Biol. Chem.* 275 (2000), 1247-1260). Their expression is almost totally restricted to brain tissues. A functional study on calbrain reveals its negative effect on Ca^{2+} /Calmodulin-dependent CaMKII activity by competitively interacting with the CaM-binding domain of CaMKII (Yamaguchi et al., *J. Biol. Chem.* 274 (1999), 3610-3616). One would expect similar roles in Ca^{2+} signaling for other CABP1 alternative products. Participation of CABP1 gene in Ca^{2+} -dependent signaling pathways would make it obvious to one skilled in the art to select this gene as a candidate for bipolar affective disorder. However, all CABP1 exons were analyzed for the presence of mutations, and surprisingly only two mutations were detected in noncoding regions.

The PIN gene (Protein inhibitor of NOS (Nitric oxide synthase)) is another obvious candidate responsible for bipolar affective disorder. Nitric oxide (NO) in the brain, may be involved in apoptosis, synaptogenesis, and neuronal development. Because NO cannot be stored in vesicles like other neurotransmitters, its release is regulated by the activity of NOS (Nitric oxide synthase). PIN is a direct inhibitor of NOS by binding and destabilizing the active homodimer complex of NOS (Jaffrey et al., *Science* 274 (1996), 774-777). PIN is highly conserved throughout the evolution and is expressed in many cell types. A recent clinical study evaluating plasma nitrate levels in depressive states suggests that NO production is increased in depression

(Suzuki et al., J. Affect. Disord. 63 (2001), 221-224) and may result from a deficiency in NOS inhibition. Moreover in a mouse model, NO synthase antagonists have been linked to antidepressant properties (Harkin et al., 1999; Karolewicz et al., Eur. J. Pharmacol. 372 (1999), 215-220). Thus, PIN would be an obvious candidate. However, due to the pleiotropic action of NO, a deficiency in PIN function would generate many unrelated disorders throughout the body. Thus, without the information presented in the disclosure herein, a person of ordinary skills in the art would have predicted PIN and not P2X7R as the gene associated with affective disorders.

The human phospholipase A2 group IB (PLA2G1B) catalyses the release of fatty acids from glycerol-3-phosphocholines. Phospholipase A2 genes (PLA2) are expressed in many tissues. Some studies have demonstrated associations between excessive PLA2 activity in brain and affective disorders (Chang et al., Neurochem. Res. 23 (1998), 887-892; Hibbeln et al., Biol. Psychiatry 25 (1989), 945-961). Moreover, other genetic studies have found associations between PLA2G1B gene and bipolar affective disorder (Dawson et al., Psychiatr. Genet. 5 (1995), 177-180). Thus, PLA2G1B represent a likely candidate for affective disorders. However in the present example, only a single silent mutation was found within exon 3 of the PLA2G1B gene.

The human citron kinase gene, Rho-associated protein (CIT) is a 183 kDa protein which associates to the GTPase Rho. CIT shares strong similarity with ROCK and ROK proteins which are other Rho-associated kinases (Madaule et al., Nature 394 (1998), 491-494). Rho GTPases are involved in many processes such as cytoskeletal organization, membrane trafficking, cell growth, and transcriptional activation (Van Aelst and D'Souza-Schorey, Genes Dev. 11 (1997), 2295-2322). Studies on brain variants of Citron-K (without the kinase domain) reveal the association with postsynaptic density proteins (PSD-95), suggesting a role in either synapse organization or function (Zhang et al., J. Neurosci. 19 (1999), 96-108; Furuyashiki et al., J. Neurosci. 19 (1999), 109-118).

The human paxillin (PXN) gene encodes for a 68 kDa protein found in focal adhesions. It is within focal adhesions where adhesion molecules dynamically

interact with the cytoskeleton (Salgia et al., J. Biol. Chem. 270 (1995), 5039-5047). The signaling pathways that regulate these dynamic interactions begin to be elucidated. Many observations suggest that paxillin is involved in transducing signals from growth factor receptors to focal adhesions. The paxillin is expressed in many tissues including brain.

However as set forth below, the gene causative for affective diseases is identified as being the P2X7 receptor (P2X7R).

Mutations were searched in coding sequences and exon-intron boundaries of the above mentioned genes since such mutations are more likely to give a functionally significant Single Nucleotide Polymorphisms (SNP). The starting sample was composed of 16 unrelated affected individuals from the Saguenay/Lac St-Jean region, which gives an 80% power to detect polymorphisms with a frequency of 0.05. To identify polymorphisms, targeted sequences were first amplified by PCR. Then, PCR products are purified on Whatman GF/C membranes (VWR, Montreal, Canada), and quantified using the PicoGreen dsDNA quantitation assay (Molecular probes, Oregon, USA). 4 ng of purified PCR products are sequenced using the DYEnamic ET terminator cycle sequencing kit (Amersham Biosciences, Baie D'Urfé, Canada). The sequencing products are resolved on an ABI PRISM 3730XL DNA analyzer, and an ABI PRISM 3700 DNA analyzer. The PCR products are sequenced in both directions. The SNPs identified in studied genes are listed in Table 10.

Table 10. Mutation analysis between markers D12S1666 and NBG9

| Genes | Positions | Variations | Alleles | Modifications |
|---------|-----------|-------------|---------|---------------|
| Rab35 | Exon06 | RABE06A | 486G-A | Silent Asn162 |
| Rab35 | Intron04 | RABI04A | 51C-T | unknown |
| Rab35 | Intron03 | RABI03A | 33G-A | unknown |
| Rab35 | Intron02 | RABI02B | 85G-A | unknown |
| Rab35 | Intron02 | RABI02A | 76C-G | unknown |
| PXN | Exon11 | PXNE11A | 1527C-T | Silent Thr509 |
| PXN | Exon06 | PXNE06A | 750C-T | Silent Ser250 |
| PXN | Exon02 | PXNE02A | 217G-A | Gly73Ser |
| PLA2G1B | Exon03 | PLA2G1BE03A | 294C-T | Silent Ser98 |
| PIN | 5'UTR01 | PINUTR01A | -49T-G | unknown |
| PIN | 5'UTR01 | PINUTR01B | -80T-C | unknown |
| PIN | Intron02 | PINI02A | 26C-T | unknown |

| PIN | Intron02 | PINI02B | 50C-T | unknown |
|-------|----------|------------|--------------|---------------|
| CaBP | Intron04 | CaBPI04A | 35C-T | unknown |
| CaBP | exon01 | CaBPE01A | -23A-G | unknown |
| OASL | Exon02 | OASLE02A | 213G-T | Silent Gly72 |
| OASL | Exon02 | OASLE02B | 408C-T | Silent Leu136 |
| OASL | Exon05 | OASLE05A | 1042G-A | Val348Met |
| OASL | Exon06 | OASLE06A | 1509G-A | Silent Ser503 |
| P2X7R | 5'UTR | P2XR7UTR5L | 362T-C | unknown |
| P2X7R | 5'UTR | P2XR7UTR5M | 532T-G | unknown |
| P2X7R | 5'UTR | P2XR7UTR5K | 1100A-G | unknown |
| P2X7R | 5'UTR | P2XR7UTR5J | 1122A-G | unknown |
| P2X7R | 5'UTR | P2XR7UTR5I | 1171C-G | unknown |
| P2X7R | 5'UTR | P2XR7UTR5F | 1351T-C | unknown |
| P2X7R | 5'UTR | P2XR7UTR5N | 1702G-A | unknown |
| P2X7R | 5'UTR | P2XR7UTR5G | 1731T-G | unknown |
| P2X7R | 5'UTR | P2XR7UTR5H | 1860C-T | unknown |
| P2X7R | 5'UTR | P2XR7UTR5A | 2162C-A | unknown |
| P2X7R | 5'UTR | P2XR7UTR5B | 2238C-T | unknown |
| P2X7R | 5'UTR | P2XR7UTR5D | 2373A-G | unknown |
| P2X7R | 5'UTR | P2XR7UTR5E | 2569G-A | unknown |
| P2X7R | 5'UTR | P2XR7UTR5C | 2702G-A | unknown |
| P2X7R | Intron01 | P2XR7I01C | 3166G-C | unknown |
| P2X7R | Intron01 | P2XR7I01A | 24778C-T | unknown |
| P2X7R | Intron01 | P2XR7I01B | 24830C-T | unknown |
| P2X7R | Exon02 | P2XR7v02A | 24942T-C | Val76Ala |
| P2X7R | Exon03 | P2XR7E03A | 26188C-T | Arg117Trp |
| P2X7R | Intron03 | P2XR7I03A | 26308A-G | unknown |
| P2X7R | Intron03 | P2XR7I03B | 26422G-A | unknown |
| P2X7R | Intron04 | P2XR7I04A | 32394G-A | unknown |
| P2X7R | Intron04 | P2XR7v05B | 32434T-C | unknown |
| P2X7R | Exon05 | P2XR7E05D | 32493G-A | Gly150Arg |
| P2X7R | Exon05 | P2XR7v05A | 32507C-T | Tyr155His |
| P2X7R | Exon05 | P2XR7E05C | 32783C-T | Silent Cys168 |
| P2X7R | Intron05 | P2XR7I05C | 32783A-C | unknown |
| P2X7R | Intron05 | P2XR7I05D | 35309T-C | unknown |
| P2X7R | Intron05 | P2XR7I05B | 35374C-T | unknown |
| P2X7R | Intron05 | P2XR7I05A | 35378A-C | unknown |
| P2X7R | Exon06 | P2XR7E06A | 35438G-A | Glu186Lys |
| P2X7R | Exon06 | P2XR7E06B | 35454T-C | Leu191Pro |
| P2X7R | Intron06 | P2XR7I06C | 35549T-C | unknown |
| P2X7R | Intron06 | P2XR7I06G | 35641G-C | unknown |
| P2X7R | Intron06 | P2XR7I06D | 35725A-C | unknown |
| P2X7R | Intron06 | P2XR7I06F | 36001T-G | unknown |
| P2X7R | Intron06 | P2XR7I06E | 36064A-T | unknown |
| P2X7R | Intron06 | P2XR7I06A | 36091DelGTTT | unknown |
| P2X7R | Intron06 | P2XR7I06B | 36108C-G | unknown |
| P2X7R | Intron07 | P2XR7I07A | 36374C-T | unknown |
| P2X7R | Intron07 | P2XR7I07B | 36378G-A | unknown |
| P2X7R | Intron07 | P2XR7I07C | 36387T-A | unknown |
| P2X7R | Intron07 | P2XR7I07D | 36398G-C | unknown |
| P2X7R | Intron07 | P2XR7I07E | 37439C-T | unknown |
| P2X7R | Intron07 | P2XR7I07F | 37513T-C | unknown |

| | | | | |
|-------|----------|------------|-----------------------------------|-----------------------|
| P2X7R | Exon08 | P2XR7E08C | 37604C-T | Arg270Cys |
| P2X7R | Exon08 | P2XR7v08A | 37605G-A | Arg270His |
| P2X7R | Exon08 | P2XR7v08B | 37623G-A | Arg276His |
| P2X7R | Exon08 | P2XR7E08D | 37633C-T | Silent Asp279 |
| P2X7R | Intron09 | P2XR7v11A | 47214C-T | unknown |
| P2X7R | Exon11 | P2XR7v11B | 47383G-A | Ala348Thr |
| P2X7R | Exon11 | P2XR7v11C | 47411C-G | Thr357Ser |
| P2X7R | Intron11 | P2XR7I11D | 47563T-C | unknown |
| P2X7R | Intron12 | P2XR7I12A | 54307C-T | unknown |
| P2X7R | Intron12 | P2XR7I12B | 54308G-A | unknown |
| P2X7R | Exon13 | P2XR7v13F | 54399C-T | unknown |
| P2X7R | Exon13 | P2XR7v13A | 54480A-G | Ala433Val |
| P2X7R | Exon13 | P2XR7v13B | 54523C-T | Gln460Arg |
| P2X7R | Exon13 | P2XR7v13G | 54562DelCCCTGAGAG CCACAGGTGCCT | Silent Pro474 |
| P2X7R | Exon13 | P2XR7v13C | 54588A-C | Del of 7aa 488 to 494 |
| P2X7R | Exon13 | P2XR7v13H | 54664C-G | PESHRL |
| P2X7R | Exon13 | P2XR7E13D | 54703G-T | Glu496Ala |
| P2X7R | Exon13 | P2XR7E13J | 54804A-T | Silent His521 |
| P2X7R | Exon13 | P2XR7v13I | 54834G-A | Silen Leu534 |
| P2X7R | Exon13 | P2XR7v13E | 54847G-A | Ile568Asn |
| P2X7R | 3'UTR | P2XR7UTR3A | 55169C-A | Arg578Gln |
| P2X7R | 3'UTR | P2XR7UTR3B | 55170A-C | Silent Pro582 |
| P2X7R | 3'UTR | P2XR7UTR3C | 55171A-C | unknown |
| P2X7R | 3'UTR | P2XR7UTR3D | 55917C-T | unknown |
| P2X7R | 3'UTR | P2XR7UTR3E | 54925G-A | unknown |
| P2X4R | 5'UTR | P2XR4UTR5I | -1956G-A | unknown |
| P2X4R | 5'UTR | P2XR4UTR5H | -1649G-A | unknown |
| P2X4R | 5'UTR | P2XR4UTR5G | -800G-A | unknown |
| P2X4R | 5'UTR | P2XR4UTR5A | -648C-A | unknown |
| P2X4R | 5'UTR | P2XR4UTR5B | -537A-G | unknown |
| P2X4R | 5'UTR | P2XR4UTR5C | -437A-G | unknown |
| P2X4R | 5'UTR | P2XR4UTR5J | -206VNRG | unknown |
| P2X4R | 5'UTR | P2XR4UTR5D | -211C-G | unknown |
| P2X4R | 5'UTR | P2XR4UTR5F | -150VNRGGGGCCCC | unknown |
| P2X4R | 5'UTR | P2XR4UTR5E | -98G-T | unknown |
| P2X4R | Intron01 | P2XR4I01A | 31G-T | unknown |
| P2X4R | Exon02 | P2XR4E02A | 262G-A | Silent mutation Ala87 |
| P2X4R | Intron02 | P2XR4I02A | 4600C-T | unknown |
| P2X4R | Intron03 | P2XR4I03A | 15G-A | unknown |
| P2X4R | Intron03 | P2XR4I03B | 72G-A | unknown |
| P2X4R | Exon04 | P2XR4E04A | 355G-A | unknown |
| P2X4R | Exon04 | P2XR4E04A | 375G-A | Ile119Val |
| P2X4R | Intron04 | P2XR4I04B | 17T-C | Silent Val125 |
| P2X4R | Intron04 | P2XR4I04A | 32G-A | unknown |
| P2X4R | Exon05 | P2XR4E05A | 465T-C | unknown |
| P2X4R | Exon07 | P2XR4E07A | 724A-G | Silent Ser155 |
| P2X4R | Intron08 | P2XR4I08A | DelT | Ser242Gly |
| P2X4R | Exon09 | P2XR4E09A | 944A-G | unknown |
| P2X4R | Intron10 | P2XR4I10A | 11G-T | Tyr315Cys |
| P2X4R | Intron10 | P2XR4I10B | G-C | unknown |
| P2X4R | Intron10 | P2XR4I10C | A-G | unknown |

| | | | | |
|--------|----------|--------------|------------------------|---------------|
| P2X4R | Intron11 | P2XR4I11B | C-G | unknown |
| P2X4R | Intron11 | P2XR4I11C | T-A | unknown |
| P2X4R | Intron11 | P2XR4I11A | 374C-T | unknown |
| CaMKK2 | 3'UTR | CaMKK2UTR3bA | 733C-T | unknown |
| CaMKK2 | 3'UTR | CaMKK2UTR3aB | 390G-A | unknown |
| CaMKK2 | 3'UTR | CaMKK2UTR3aA | 239G-A | unknown |
| CaMKK2 | Intron15 | CaMKK2I15B | 325T-C | unknown |
| CaMKK2 | Intron15 | CaMKK2I15A | 169G-A | unknown |
| CaMKK2 | Intron14 | CaMKK2I14A | 224A-G | unknown |
| CaMKK2 | Intron10 | CaMKK2I10A | 156DelGTGATCCGCCT G | unknown |
| CaMKK2 | intron09 | CaMKK2I09B | 528A-G | unknown |
| CaMKK2 | intron09 | CaMKK2I09A | 521A-G | unknown |
| CaMKK2 | Exon09 | SNP6f18v5 | 1095C-A | Silent Ile365 |
| CaMKK2 | Exon09 | SNP6f18v4 | 1087C-T | Arg363Cys |
| CaMKK2 | Exon05 | CaMKKE05A | 687C-T | Silent Pro229 |
| CaMKK2 | Intron03 | CaMKK2I03A | 10C-T | unknown |
| CaMKK2 | Intron02 | CaMKK2I02A | 39C-T | unknown |
| CaMKK2 | Intron01 | CaMKK2I01B | 2911G-C | unknown |
| CaMKK2 | Intron01 | CaMKK2I01A | 89C-A | unknown |
| CaMKK2 | Exon01 | SNP6f18v2 | 253A-T | Thr85Ser |
| CaMKK2 | Exon01 | SNP6f18v1 | 29G-A | Ser10Asn |
| CaMKK2 | 5'UTR01 | CaMKK2UTR01B | 253T-C | unknown |
| CaMKK2 | 5'UTR01 | CaMKK2UTR01A | 63C-A | unknown |
| APC5 | Intron01 | APC5I01A | 10G-T | unknown |
| APC5 | Intron01 | APC5I01B | 50A-T | unknown |
| APC5 | Intron05 | APC5I05A | 73T-C | unknown |
| APC5 | Intron06 | APC5I06A | 73T-G | unknown |
| APC5 | Exon11 | APC5E11A | 1416C-T | Silent His472 |

Each SNP in genes Rab35, PXN, PLA2G1B, PIN, CaBP, OASL, P2X4R, CaMKK2 and APC5 was designated according to the gene where it was found, and its location in that gene (intronic or exonic regions). Each SNP in the P2X7R gene was designated according to their position on SEQ ID NO: 1. The allele describes the position and the variation observed. In coding regions, the position is relative to the start codon, whereas the intronic SNPs are positioned relative to the beginning of the corresponding intron (when known). Primers used for identifying the SNPs in the P2X7R and the location of each SNPs included in tables 2 and 12 are defined in table 1a and SEQ ID NOs 52 to 111.

Association studies using missense SNPs were performed. Missense SNPs or SNPs that could be closed to the splice sites were used, because it is more likely that

diseases would be associated to an improper function in proteins. Case group was composed by bipolar I individuals, schizoaffective bipolar type (182 subjects) and bipolar II diagnosed persons (31 subjects). Many controls from the Saguenay/Lac-St-Jean region, were sampled from Steinert, Glaucoma and Paget DNA banks. The control individuals were not diagnosed for affective disorders. According to the lifetime risks of bipolar disorders (1%), there is no need to screen controls for psychiatric disorders.

Direct sequencing of PCR products is by far the most accurate method of analysis and is the method of choice in view of our sequencing platform capacity. PCR products were analyzed by direct sequencing as described above. After sequencing analysis, individuals are automatically typed for the corresponding SNP using a home-developed program, GENO.pl. The results of SNP genotyping are compiled in a 4D database.

The association hypothesis was tested with CLUMP (Sham & Curtis 1995, Ann. Hum. Genet. 59:97-105). One thousand simulations were used to estimate p-values. Results are illustrated in table 11. The T1 statistic, which is the usual chi-squared statistic on the raw contingency table, was used to test for allelic association. Moreover, the largest chi-squared statistic got by comparing one column of the original table against the total of the other columns, called T3 statistic, was added to the previous one to test for potential genotype association since T1 statistic results may be biased when the contingency table contains cells with low values.

Table 11. Association hypothesis using CLUMP

| gene | SNPs | Effective | | Allele Analysis p-value (T1) | Genotype Analysis | |
|--------|-----------|-----------|----------|---------------------------------------|-------------------|-----------------|
| | | Cases | Controls | | p-value (T2) | p-value (T3) |
| P2X7R | P2XR7v11B | 208 | 211 | 0.795 | 0.036 | 0.028 |
| | P2XR7v13A | 212 | 214 | 0.344 | 0.250 | 0.186 |
| | P2XR7v13E | 212 | 211 | 0.780 | 0.017 | 0.017 |
| CAMKK2 | SNP6f18v5 | 206 | 135 | 1.00 | 1.00 | 1.00 |
| | SNP6f18v4 | 206 | 135 | 0.816 | 0.962 | 0.841 |
| | SNP6f18v2 | 205 | 135 | 0.057 | 0.110 | 0.095 |
| | SNP6f18v1 | 206 | 135 | 0.512 | 0.532 | 0.385 |

The association studies using SNPs in P2X7, P2X4, and CaMKK2 reveal associations significant at level of about 5% or less. Three genotype associations in P2X7 were observed. However, SNPs P2XR7v11B and P2XR7v13E are closely linked together based on a contingency table. There is also an allele association at level of 5,7% for SNP6f18v2 in CaMKK2. The information associated to each relevant SNP can be found in Tables 10 and 12.

EXAMPLE 3

Polymorphisms found in the P2X7R in individuals suffering from depression

Association studies using SNPs in the P2X7R gene was performed in a case/control sample (535 individuals) from a German population. The case group was composed of 36 individuals diagnosed with bipolar type I or type II, and 279 individuals diagnosed with unipolar disorders (i.e. depression) representing 133 affected males and 182 affected females. Among controls, we count The remaining 220 control individuals were normal (i.e. diagnosed as non depressive), and comprising 81 males, 182 females and 14 of unknown gender. The same sexual distribution was noted in both groups.

SNPs were identified in this sample by using a subgroup of 24 affected individuals. Similar SNPs in the P2X7R gene detected in the German population were similar if not identical to the SNPs seen in the Saguenay/Lac-St-Jean population (see table

12). Other rare missense SNPs were also noted in the German population, such as Arg117Trp (P2XR7E03A), Glu186Lys (P2XR7E06A), Leu191Pro (P2XR7E06B), Ile568Asn (P2XR7E13J). These amino acids are quite conserved between ortholog P2X7 genes. It is possible that the Ile568Asn (P2XR7E13J) mutation may be involved in the surface expression of P2X7.

Table 12. Comparison between polymorphisms in the Saguenay/Lac-St-Jean population and the German population in the human P2X7R gene

| Associated exons or introns | Variation (SNP or others) | Allele | Position* | Modification | Frequency (Canada) | Frequency (Germany) |
|-----------------------------|---------------------------|--------|-----------|---------------|--------------------|---------------------|
| 5'UTR | P2XR7UTR5L | T-C | 362 | unknown | | |
| 5'UTR | P2XR7UTR5M | T-G | 532 | unknown | 0,13 | 0,08 |
| 5'UTR | P2XR7UTR5K | A-G | 1100 | unknown | 0,16 | 0,1 |
| 5'UTR | P2XR7UTR5J | A-G | 1122 | unknown | 0,13 | 0,13 |
| 5'UTR | P2XR7UTR5I | C-G | 1171 | unknown | 0,13 | 0,13 |
| 5'UTR | P2XR7UTR5F | T-C | 1351 | unknown | 0,06 | 0,02 |
| 5'UTR | P2XR7UTR5N | G-A | 1702 | unknown | 0,3 | 0,12 |
| 5'UTR | P2XR7UTR5G | T-G | 1731 | unknown | - | 0,02 |
| 5'UTR | P2XR7UTR5H | C-T | 1860 | unknown | 0,17 | 0,15 |
| 5'UTR | P2XR7UTR5A | C-A | 2162 | unknown | 0,07 | 0,15 |
| 5'UTR | P2XR7UTR5B | C-T | 2238 | unknown | 0,07 | 0,12 |
| 5'UTR | P2XR7UTR5D | A-G | 2373 | unknown | 0,3 | 0,27 |
| 5'UTR | P2XR7UTR5E | G-A | 2569 | unknown | 0,07 | 0,12 |
| 5'UTR | P2XR7UTR5C | G-A | 2702 | unknown | 0,1 | 0,02 |
| Intron01 | P2XR7I01C | G-C | 3166 | unknown | 0,31 | 0,27 |
| Intron01 | P2XR7I01A | C-T | 24778 | unknown | 0,03 | - |
| Intron01 | P2XR7I01B | C-T | 24830 | unknown | 0,03 | - |
| Exon02 | P2XR7v02A | T-C | 24942 | Val76Ala | 0,03 | RARE |
| Exon03 | P2XR7E03A | C-T | 26188 | Arg117Trp | 0,06 | 0,08 |
| Intron03 | P2XR7I03A | A-G | 26308 | unknown | - | RARE |
| Intron03 | P2XR7I03B | G-A | 26422 | unknown | 0,7 | 0,44 |
| Intron04 | P2XR7I04A | G-A | 32394 | unknown | 0,18 | 0,12 |
| Intron04 | P2XR7v05B | T-C | 32434 | unknown | 0,03 | 0,01 |
| Exon05 | P2XR7E05D | G-A | 32493 | Gly150Arg | 0,33 | 0,29 |
| Exon05 | P2XR7E05E | G-A | 32506 | Silent Val154 | RARE | 0,02 |
| Exon05 | P2XR7v05A | C-T | 32507 | Tyr155His | - | RARE |
| Exon05 | P2XR7E05C | C-T | 32548 | Silent Cys168 | 0,33 | 0,44 |
| Intron05 | P2XR7I05C | A-C | 32783 | unknown | RARE | 0,02 |
| Intron05 | P2XR7I05D | T-C | 35309 | unknown | 0,25 | - |
| Intron05 | P2XR7I05B | C-T | 35374 | unknown | ND | 0,35 |
| | | | | | 0,7 | 0,67 |

| | | | | | | |
|----------|------------|----------------------------------|-------------|---------------------------------------|-------|------|
| Intron05 | P2XR7I05A | A-C | 35378 | unknown | 0,7 | 0,65 |
| Exon06 | P2XR7E06A | G-A | 35438 | Glu186Lys | - | 0,02 |
| Exon06 | P2XR7E06B | T-C | 35454 | Leu191Pro | - | 0,02 |
| Intron06 | P2XR7I06C | T-C | 35549 | unknown | 0,04 | 0,08 |
| Intron06 | P2XR7I06G | G-C | 35641 | unknown | - | 0,02 |
| Intron06 | P2XR7I06D | A-C | 35725 | unknown | 0,21 | 0,27 |
| Intron06 | P2XR7I06F | T-G | 36001 | unknown | 0,17 | 0,3 |
| Intron06 | P2XR7I06E | A-T | 36064 | unknown | 0,11 | 0,1 |
| Intron06 | P2XR7I06A | DelGTTT | 36091-36094 | unknown | 0,14 | 0,3 |
| Intron06 | P2XR7I06B | C-G | 36108 | unknown | 0,14 | 0,29 |
| Intron07 | P2XR7I07A | C-T | 36374 | unknown | 0,07 | - |
| Intron07 | P2XR7I07B | G-A | 36378 | unknown | 0,21 | 0,28 |
| Intron07 | P2XR7I07C | T-A | 36387 | unknown | 0,21 | 0,28 |
| Intron07 | P2XR7I07D | G-C | 36398 | unknown | 0,42 | 0,4 |
| Intron07 | P2XR7I07E | C-T | 37439 | unknown | 0,41 | - |
| Intron07 | P2XR7I07F | T-C | 37513 | unknown | - | RARE |
| Exon08 | P2XR7E08C | C-T | 37604 | Arg270Cys | RARE | - |
| Exon08 | P2XR7v08A | G-A | 37605 | Arg270His | 0,46 | 0,24 |
| Exon08 | P2XR7v08B | G-A | 37623 | Arg276His | 0,03 | 0,02 |
| Exon08 | P2XR7E08D | C-T | 37633 | Silent Asp279 | RARE | - |
| Intron09 | P2XR7v11A | C-T | 47214 | unknown | 0,08 | 0,03 |
| Exon11 | P2XR7v11B | G-A | 47383 | Ala348Thr | 0,5 | 0,44 |
| Exon11 | P2XR7v11C | C-G | 47411 | Thr357Ser | 0,08 | 0,07 |
| Intron11 | P2XR7I11D | T-C | 47563 | unknown | 0,43 | 0,44 |
| Intron12 | P2XR7I12A | C-T | 54307 | unknown | 0,32 | - |
| Intron12 | P2XR7I12B | G-A | 54308 | unknown | 0,03 | - |
| Exon13 | P2XR7v13F | C-T | 54399 | Ala433Val | 0,13 | - |
| Exon13 | P2XR7v13A | A-G | 54480 | Gln460Arg | 0,13 | 0,17 |
| Exon13 | P2XR7v13B | C-T | 54523 | Silent Pro474 | 0,1 | 0,07 |
| Exon13 | P2XR7v13G | DelCCCTGAGA GCCACAGG TGCCT | 54562-54582 | Del of 7aa 488 to 494 (PESHRCL) | RARE | - |
| Exon13 | P2XR7v13C | A-C | 54588 | Glu496Ala | 0,13 | 0,06 |
| Exon13 | P2XR7v13H | C-G | 54664 | His521Gln | 0,03 | - |
| Exon13 | P2XR7E13D | G-T | 54703 | Silent Leu534 | 0,1 | 0,02 |
| Exon13 | P2XR7E13J | A-T | 54804 | Ile568Asn | - | 0,01 |
| Exon13 | P2XR7v13I | G-A | 54834 | Arg578Gln | - | RARE |
| Exon13 | P2XR7v13E | G-A | 54847 | Silent Pro582 | 0,4 | 0,45 |
| 3'UTR | P2XR7UTR3A | C-A | 55169 | unknown | 0,48 | 0,37 |
| 3'UTR | P2XR7UTR3B | A-C | 55170 | unknown | 0,09 | 0,1 |
| 3'UTR | P2XR7UTR3C | A-C | 55171 | unknown | 0,05 | 0,06 |
| 3'UTR | P2XR7UTR3D | C-T | 55917 | unknown | 0,001 | - |
| 3'UTR | P2XR7UTR3E | G-A | 54925 | unknown | - | 0,01 |

The position and numbering of the polymorphism corresponds to the human P2X7R gene as defined in SEQ ID NO: 1. To identify the genomic organization of the P2X7R gene, BAC clones were firstly organized using known polymorphic markers, sequence tag sites (STSs), BAC-end sequences and expressed sequence tags (ESTs). Unorientated and unordered DNA regions were reassembled into a sequences using Phrap and reordered the pieces using P2X7R exons as scaffolds. No complete gene organization for P2X7R has been done. There is only a partial gene structure from exon6 to 13, NT_037809. Therefore, this genomic sequence encompassing the P2X7R gene as depicted in SEQ ID NO: 1 could contain some sequence errors, specifically in intronic regions. Primers used for SNP amplification and sequencing are shown in Table 1a and depicted in SEQ ID NOs: 52 to 111.

Statistical analysis was performed according to the CLUMP method (Sham & Curtis 1995, Ann. Hum. Genet. 59:97-105). Table 13 resumes the allelic and genotypic association studies for SNPs in P2X7 gene.

Table 13. Allelic and genotypic association studies using CLUMP

| Locus | Allele Frequencies* | Effective | | Allele Analysis p-value (T1) | Genotype Analysis | |
|------------|----------------------|-----------|----------|------------------------------|-------------------|--------------|
| | | Cases | Controls | | p-value (T1) | p-value (T3) |
| P2XR7v02A | 2(0.09); 4(0.91) | 313 | 218 | 0.501 | 0.729 | 0.591 |
| P2XR7I04A | 1(0.04); 3(0.96) | 314 | 220 | 0.604 | 0.433 | 0.348 |
| P2XR7v05B | 2(0.69); 4(0.31) | 314 | 220 | 0.133 | 0.270 | 0.325 |
| P2XR7E05D | 1(0.03); 3(0.97) | 314 | 220 | 0.842 | 0.827 | 0.827 |
| P2XR7E05E | 1(0.006); 3(0.994)** | 314 | 220 | 0.048 | 0.045 | 0.045 |
| P2XR7v05A | 2(0.60); 4(0.40) | 314 | 220 | 0.038 | 0.144 | 0.219 |
| P2XR7E05C | 2(0.98); 4(0.02) | 314 | 220 | 1.000 | 1.000 | 1.000 |
| P2XR7I07F | 2(0.002); 4(0.98) | 315 | 219 | 1.000 | 1.000 | 1.000 |
| P2XR7v08A | 1(0.23); 3(0.77) | 315 | 219 | 0.454 | 0.673 | 0.634 |
| P2XR7v08B | 1(0.02); 3(0.98) | 315 | 219 | 0.636 | 0.638 | 0.638 |
| P2XR7v11A | 2(0.95); 4(0.05) | 311 | 218 | 0.348 | 0.391 | 0.436 |
| P2XR7v11B | 1(0.45); 3(0.55) | 312 | 218 | 0.605 | 0.803 | 0.790 |
| P2XR7v11C | 2(0.93); 3(0.07) | 312 | 218 | 0.793 | 0.256 | 0.924 |
| P2XR7I11D | 2(0.45); 4(0.55) | 312 | 219 | 0.665 | 0.735 | 0.740 |
| P2XR7v13A | 1(0.87); 3(0.13) | 305 | 215 | 0.017 | <0.001 | <0.001 |
| P2XR7v13B | 2(0.93); 4(0.07) | 305 | 216 | 1.000 | 0.228 | 0.677 |
| P2XR7v13C | 1(0.91); 2(0.09) | 305 | 216 | 0.151 | 0.006 | 0.008 |
| P2XR7E13D | 3(0.94); 4(0.06) | 315 | 219 | 0.402 | 0.429 | 0.474 |
| P2XR7E13J | 1(0.01); 4(0.99) | 315 | 219 | 0.618 | 0.603 | 0.603 |
| P2XR7E13I | 1(0.004); 3(0.996) | 315 | 219 | 0.999 | 1.000 | 1.000 |
| P2XR7v13E | 1(0.46); 3(0.54) | 314 | 219 | 0.699 | 0.866 | 0.845 |
| P2XR7UTR3E | 1(0.02); 3(0.98) | 315 | 219 | 0.147 | 0.161 | 0.161 |

* The column Allele Frequencies presents the allele for each SNP (A=1, C=2, G=3, T=4) and their respective frequency.

** For this SNP we observed a zero cell in both (allele and genotype) 2X2 contingency tables. p-value= 0.045 was observed exact Fisher test.

For the SNP analysis, the Hardy-Weinberg (HW) equilibrium was controlled in the control samples. The Hardy-Weinberg principle (HWP) maybe stated as follow: In a large, randomly mating population, in which there is no migration, or selection against a particular genotype and the mutation rate remains constant, the proportions of the various genotypes will remain unchanged from one generation to another. Take a two allele system with alleles A and a. If the proportion of A in the population is represented as p and the proportion of a as q, then p plus q represent the sum total of alleles at this locus, that is $p+q=1$. The HWP is useful to evaluate some population problems like marital assortment, Inbreeding, population stratification, admixture, decreased viability of a particular genotype. The SNP P2XR7v13A did not respect the Hardy-Weinberg equilibrium.

The association hypothesis was also tested using an allele positivity table known to be suitable for the detection of susceptibility alleles showing a dominant mode of inheritance (Ohashi and Tokunaga, J. Hum. Genet. 44 (1999), 246-248; Ohashi et al., Ann. Hum. Genet. 65 (2001), 197-206). Similar results were obtained using this method as those obtained using the allele frequency tables, with the exception of P2XR7v05A where the p-values were 0.253. Thus, P2XR7v05A presented a less significant association in this analysis. This difference can be attributed to the mode of inheritance.

The proportion of unipolar individuals in analysis of the German population is quite important since the American Psychiatric Association (Diagnostic and Statistical Manual of Mental Disorders-4th Edition Text Revision (DMS-IV-TR), American Psychiatric Press, 2000) has reported an increase in susceptibility for unipolar disorders in female groups. To determine whether the sexual variable could influence the association analysis, additional association studies were performed by controlling the sexual parameter. Normal individuals in the German population without gender information were omitted from the study. Then, a logistic regression model was

derived by including the sex as factor. In order to obtain a model that is as stable as possible, the regression model was minimised by using the difference between log-likelihood's for models with or without interaction (Hosmer, and Lemeshow, "Applied logistic regression", John Wiley and Sons, 1989). The strategy used for handling the zero cells from contingency tables was to eliminate associated category completely. Calculations were done with SAS v8.0 SAS is a statistical software package that allows the user to manipulate and analyze data in many different ways. Because of its capabilities, this software package is used in many disciplines, including medical sciences, biological sciences, and social sciences.

The introduction of a sexual parameter did not perturb the association already observed in previous analysis. Moreover, this analysis model revealed additional results: a potential allele association with P2XR7v05B ($p=0.064$), and a genotypic association for P2XR7v08A ($p=0.042$) was observed.

Association studies using pooled samples was performed by merging individuals from the samples of the Saguenay/Lac St-Jean with those of the German population. Results are illustrated in table 14. The aim of this analysis is to highlight common features between both populations. However, according to differences between both samples (mainly the phenotype of affected individuals i.e. bipolar disorder in the Saguenay/Lac St-Jean samples, versus mostly unipolar disorder in the German population) some parameters were controlled, including sex and ethnicity. The modelling strategy for logistic regressions was described above.

Table 14. Association studies using pooled samples from both populations

| Locus | Allele analysis | | Genotype analysis | |
|--------------|------------------------|------------------------|--------------------------|------------------------|
| | p-value for SNP | p-value for sex | p-value for SNP | p-value for sex |
| P2XR7v02A | 0.8254 | 0.0085 | 0.8650 | 0.4531 |
| P2XR7v05B | 0.1751 | 0.3714 | 0.2034 | 0.5110 |
| P2XR7v05A | 0.3808 | 0.0266 | 0.0885 | 0.1392 |
| P2XR7v08A | 0.0452 | 0.0041 | 0.1021 | 0.3452 |
| P2XR7v08B | 0.3471 | 0.0040 | 0.3413 | 0.3617 |
| P2XR7v11A | 0.3559 | 0.0136 | 0.5888 | 0.4404 |
| P2XR7v11B | 0.5902 | 0.0093 | 0.3897 | 0.4302 |
| P2XR7v11C | 0.3731 | 0.0094 | 0.7648 | 0.4615 |

| | <i>Allele analysis</i> | | <i>Genotype analysis</i> | |
|-----------|------------------------|--------|--------------------------|--------|
| P2XR7v13A | 0.0047 | 0.0209 | <0.0001 | 0.4814 |
| P2XR7v13B | 0.5129 | 0.2352 | 0.9584 | 0.4092 |
| P2XR7v13C | 0.2466 | 0.0284 | 0.2225 | 0.4228 |
| P2XR7v13E | 0.8168 | 0.0159 | 0.3713 | 0.4990 |

An allelic and genotypic association was observed for the P2XR7v13A locus ($p=0.0047$) which was stronger than in the separate analyses. A significant allelic association was also noted for the P2XR7v08A locus ($p=0.0452$). In addition, the present analysis also demonstrate the potential relationship between SNP P2XR7v05A and the origin with a p -value= 0.0515 (not shown in the table) which is in agreement with previous association analysis done in both samples separately (see Table 13).

The haplotype analysis was performed using the German population. The PHASE program (Stephens et al., Am. J. Hum. Genet. 68 (2001), 978-989) was used to estimate SNPs haplotypes within exons of the P2X7R gene. Haplotypes were created for each exon having more than one associated SNP (see Table 15 for exon-associated SNPs). Case groups varied from 218-220 individuals, whereas control groups varied between 312-316 individuals. Association hypothesis was tested with the CLUMP method since many haplotypes were created for each exon. T1 and T3 statistic tests performed as described above. T2 and T4 statistics were also calculated owing to the presence of small effective cells in the contingency tables. T2 statistic is the usual chi-squared statistic applied on the contingency table obtained after collapsing columns with small expected values. T4 statistic is the largest chi-squared statistic obtained by comparing one column of the original table against the total of the other columns. One thousand simulations were used to estimate p -values. The resulting data was analyzed with the logistic regression model (describe above) using SAS V8.0 in order to consider the sexual parameter (for these tests the sample was reduced by 14 normal individuals). However, this analysis method is limited by the reliability of reconstructed haplotypes.

Table 15. Exon-associated SNPs

| <i>Exons</i> | <i>Associated SNPs</i> |
|--------------|---|
| 5 | P2XR7E05D P2XR7E05E P2XR7v05A P2XR7E05C |
| 8 | P2XR7v08A P2XR7v08B |
| 11 | P2XRv11B P2XRv11C |
| 13 | P2XR7v13A P2XR7v13B P2XR7v13C P2XR7E13D P2XR7E13J P2XR7v13I P2XR7v13E |

Table 16. Genotypic association with haplotypes in exon 13 of P2X7R

| <i>Exon (haplotype)</i> | <i>Allele analysis</i> | | | <i>Genotype analysis</i> | | |
|-----------------------------|--|--------------|----------------|---|--------------|----------------|
| | Clump* | p-value(sex) | p-value(haplo) | Clump | p-value(sex) | p-value(haplo) |
| 5(5) | T1:0.032 T2:0.068 T3:0.054 T4:0.059 | 0.3133 | 0.1947 | T1:0.193 T2:0.159 T3:0.099 T4:0.304 | 0.460 | 0.5355 |
| 8(3) | T1:0.551 T2:0.585 T3:0.646 T4:0.646 | 0.3813 | 0.3064 | T1:0.812 T2:0.689 T3:0.644 T4:0.756 | 0.5428 | 0.6652 |
| 11(3) | T1:0.750 T2:0.786 T3:0.726 T4:0.726 | 0.0886 | 0.7396 | T1:0.625 T2:0.919 T3:0.929 T4:0.921 | 0.2305 | 0.9494 |
| 13(15**) | T1:0.088 T2:0.079 T3:0.147 T4:0.072 | 0.1871 | 0.1264 | T1:0.001 T2:0.002 T3:0.057 T4:<0.001 | 0.4610 | 0.019 |

*T1 test should not be considered because of contingency tables with zero cells.
 **Among these 15 haplotypes, we observed 8 haplotypes where case cells have less than 3 individuals.

Table 16 illustrates a genotypic association with haplotypes in exon 13 of the P2X7R genes. Interestingly, many haplotypes for the exon 13 were observed. The

differences between statistics in exon 13 (T3 less significant) can be explained by the involvement of more than one genotype of haplotypes in the disease. A potential allelic association was also noted with haplotypes in exon 5 of the P2X7R gene.

EXAMPLE 4

P2X7R gene structure and mRNA expression and transcript sequence

A 1700 bp nucleotide sequence corresponding to the human P2X7R promoter was analyzed by using MatInspector V2.2 and Transfac 4.0 algorithms. This analysis showed that the P2X7R gene does not contain a standard TATA box, but has SP1 sites that can make up for transcriptional initiation. Besides the SP1 sequences, there are binding sites for the transcription factors GATA, Oct and Ikarus. These sites are thought to provide tissue specificity. Interestingly, the P2X7R promoter has binding sites that suggest responsiveness to different cytokines such as AP-1, NFAT and CEBPB.

P2X7R possesses 13 exons and 12 introns (Buell et al., Receptors Channels 5 (1998), 347), providing a basis for alternative splicing that would yield in theory different transcripts and produce different isoforms with possible different functions. No alternatively spliced variant was clearly identified. However, experiments of EST clustering allowed the description of three splicing variants. One is defined by the lack of the exon 5. This P2X7v02 variant corresponds to the clone IMAGE: 3628076 isolated from brain-derived cell lines. The P2x7v02 lacking the exon 5 produces a frame shift, thus generating a shorter polypeptide. The second splicing variant, P2X7v03, is characterized by the presence of the short intron 10 into the mRNA. This variant is supported by two high quality sequences, the cDNA clone BRAMY2008977 (AC number: AK090866) from human amygdala and the EST clone dbEST:7339877 derived from an unknown human tumor. The last variant, P2X7v04, is defined by the lack of the first exon that suggests an alternative promoter usage closed to the exon 2. A high quality EST clone dbEST:4782844 derived from a head and neck tumor supports this variant. These variants are shown in Figures 16a to 16e.

P2X7 variants.

| | | | | | | | | | | | | | | | | | |
|---------|--------|--------------------------------------|-----|-------|-------|-------|------|-------|------|-------|------|------|-----|-----|-----|-----|-----|
| P2X7v01 | MPACCS | SDV | FQY | ETNK | VTRI | QSM | NYGT | IKW | FFHV | IIFS | VVCF | FALV | SDK | LYQ | RKE | PVI | SS |
| P2X7v04 | MPPVD | -----AFPCLPFS-----FALVSDKLYQRKEPVISS | | | | | | | | | | | | | | | |
| P2X7v02 | MPACCS | SDV | FQY | ETNK | VTRI | QSM | NYGT | IKW | FFHV | IIFS | VVCF | FALV | SDK | LYQ | RKE | PVI | SS |
| P2X7v03 | MPACCS | SDV | FQY | ETNK | VTRI | QSM | NYGT | IKW | FFHV | IIFS | VVCF | FALV | SDK | LYQ | RKE | PVI | SS |
| | 1 | | 10 | | 20 | | 30 | | 40 | | 50 | | | | | | |
| P2X7v01 | VHTK | VKG | IAE | VKEE | IVENG | VKKL | VHS | VFD | TADY | TFPL | QNS | FFVM | TN | FLK | TEG | QE | QRL |
| P2X7v04 | VHTK | VKG | IAE | VKEE | IVENG | VKKL | VHS | VFD | TADY | TFPL | QNS | FFVM | TN | FLK | TEG | QE | QRL |
| P2X7v02 | VHTK | VKG | IAE | VKEE | IVENG | VKKL | VHS | VFD | TADY | TFPL | QNS | FFVM | TN | FLK | TEG | QE | QRL |
| P2X7v03 | VHTK | VKG | IAE | VKEE | IVENG | VKKL | VHS | VFD | TADY | TFPL | QNS | FFVM | TN | FLK | TEG | QE | QRL |
| | 61 | | 70 | | 80 | | 90 | | 100 | | 110 | | | | | | |
| P2X7v01 | EYP | TRR | TL | CSS | DRG | CKK | GW | MDP | QSK | GI | QT | GRC | VV | HEG | NQ | KT | CEV |
| P2X7v04 | EYP | TRR | TL | CSS | DRG | CKK | GW | MDP | QSK | GI | QT | GRC | VV | HEG | NQ | KT | CEV |
| P2X7v02 | EYP | TRR | TL | CSS | DRG | CKK | GW | MDP | QSK | GI | QT | GRC | VV | HEG | NQ | KT | CEV |
| P2X7v03 | EYP | TRR | TL | CSS | DRG | CKK | GW | MDP | QSK | GI | QT | GRC | VV | HEG | NQ | KT | CEV |
| | 121 | | 130 | | 140 | | 150 | | 160 | | 170 | | | | | | |
| P2X7v01 | LLNS | AE | NFT | VL | IK | NN | ID | FP | GH | NY | TT | RN | IL | PGL | NIT | CT | FH |
| P2X7v04 | LLNS | AE | NFT | VL | IK | NN | ID | FP | GH | NY | TT | RN | IL | PGL | NIT | CT | FH |
| P2X7v02 | LLNS | AE | NFT | VL | IK | NN | ID | FP | GH | NY | TT | RN | IL | PGL | NIT | CT | FH |
| P2X7v03 | LLNS | AE | NFT | VL | IK | NN | ID | FP | GH | NY | TT | RN | IL | PGL | NIT | CT | FH |
| | 181 | | 190 | | 200 | | 210 | | 220 | | 230 | | | | | | |
| P2X7v01 | NFSD | VAI | QGG | IM | GIE | IY | WDC | NL | DR | WF | HH | CH | PK | YS | FR | RL | DD |
| P2X7v04 | NFSD | VAI | QGG | IM | GIE | IY | WDC | NL | DR | WF | HH | CH | PK | YS | FR | RL | DD |
| P2X7v02 | NFSD | VAI | QGG | IM | GIE | IY | WDC | NL | DR | WF | HH | CH | PK | YS | FR | RL | DD |
| P2X7v03 | NFSD | VAI | QGG | IM | GIE | IY | WDC | NL | DR | WF | HH | CH | PK | YS | FR | RL | DD |
| | 241 | | 250 | | 260 | | 270 | | 280 | | 290 | | | | | | |
| P2X7v01 | ENN | VE | KRT | LI | KV | GI | RF | DI | LV | FG | TG | GK | FD | II | QL | VV | YG |
| P2X7v04 | ENN | VE | KRT | LI | KV | GI | RF | DI | LV | FG | TG | GK | FD | II | QL | VV | YG |
| P2X7v02 | ENN | VE | KRT | LI | KV | GI | RF | DI | LV | FG | TG | GK | FD | II | QL | VV | YG |
| P2X7v03 | ENN | VE | KRT | LI | KV | GI | RF | DI | LV | FG | TG | GK | FD | II | QL | VV | YG |
| | 301 | | 310 | | 320 | | 330 | | 340 | | 350 | | | | | | |
| P2X7v01 | NCCR | SHI | YP | WCK | CC | QPC | CV | VNE | YYY | RKK | CES | IVE | PK | PT | LKY | VS | FV |
| P2X7v04 | NCCR | SHI | YP | WCK | CC | QPC | CV | VNE | YYY | RKK | CES | IVE | PK | PT | LKY | VS | FV |
| P2X7v02 | NCCR | SHI | YP | WCK | CC | QPC | CV | VNE | YYY | RKK | CES | IVE | PK | PT | LKY | VS | FV |
| P2X7v03 | NCCR | SHI | YP | WCK | CC | QPC | CV | VNE | YYY | RKK | CES | IVE | PK | PT | LKY | VS | FV |
| | EGSD | ----- | | | | | | | | | | | | | | | |
| | 361 | | 370 | | 380 | | 390 | | 400 | | 410 | | | | | | |
| P2X7v01 | LQDV | KG | QEV | PR | PAM | DF | TD | LS | RL | PL | AL | HD | TP | PI | PG | QEE | IQL |
| P2X7v04 | LQDV | KG | QEV | PR | PAM | DF | TD | LS | RL | PL | AL | HD | TP | PI | PG | QEE | IQL |
| P2X7v02 | LQDV | KG | QEV | PR | PAM | DF | TD | LS | RL | PL | AL | HD | TP | PI | PG | QEE | IQL |
| P2X7v03 | LQDV | KG | QEV | PR | PAM | DF | TD | LS | RL | PL | AL | HD | TP | PI | PG | QEE | IQL |
| | 421 | | 430 | | 440 | | 450 | | 460 | | 470 | | | | | | |
| P2X7v01 | SCLP | SQ | LP | ESH | RCL | EEL | CCR | KK | PG | AC | IT | TSEL | FR | KL | VLS | SR | HV |
| P2X7v04 | SCLP | SQ | LP | ESH | RCL | EEL | CCR | KK | PG | AC | IT | TSEL | FR | KL | VLS | SR | HV |
| P2X7v02 | SCLP | SQ | LP | ESH | RCL | EEL | CCR | KK | PG | AC | IT | TSEL | FR | KL | VLS | SR | HV |
| P2X7v03 | SCLP | SQ | LP | ESH | RCL | EEL | CCR | KK | PG | AC | IT | TSEL | FR | KL | VLS | SR | HV |
| | 481 | | 490 | | 500 | | 510 | | 520 | | 530 | | | | | | |
| P2X7v01 | TNSR | LR | HC | AY | RC | YA | TW | RF | GS | QD | MA | DF | AI | PS | CC | RR | IR |
| P2X7v04 | TNSR | LR | HC | AY | RC | YA | TW | RF | GS | QD | MA | DF | AI | PS | CC | RR | IR |
| P2X7v02 | TNSR | LR | HC | AY | RC | YA | TW | RF | GS | QD | MA | DF | AI | PS | CC | RR | IR |
| P2X7v03 | TNSR | LR | HC | AY | RC | YA | TW | RF | GS | QD | MA | DF | AI | PS | CC | RR | IR |
| | 541 | | 550 | | 560 | | 570 | | 580 | | 590 | | | | | | |

Therefore the transcriptional and translational start sequences of the human P2X7R were analyzed using Blast, Genescan and HMMgene computer software. This analysis indicated that P2X7R possesses with high probability only one translation start site. Most P2X7R expression sequence tags (ESTs; Unique cluster Hs. 193470) having a reliable 5' end showed identical transcriptional start site. None of the ESTs showed any indication of alternative splicing. Therefore, *in silico* analysis suggests that there is a low probability to find different transcripts produced by alternative splicing or alternative promoter usage.

The above mentioned *in silico* data were confirmed by RT-PCR analysis spanning the whole predicted human P2X7R coding sequence using 14 and 19 bases (5'-ATGCCGGCTTGCTG-3'; 5'-GTAGGGATACTTGAAGCCA-3') oligonucleotides corresponding to the beginning and end of the coding sequence, respectively. Total RNA from whole brain, different dissected brain areas, thymus, spleen and kidney were isolated and analyzed for P2X7R expression. RT-PCR reactions were performed using the C. Therm One Step polymerase system (Roche Applied Science) and a protocol for touch down PCR with hot start. Briefly, Reverse Transcription was performed at 52°C according to the manufacturer's conditions. PCR reactions were executed with an annealing temperatures of 64°C for the first five cycles and of 54°C for the next 30 cycles.

A single specific band of the size of 1785 bp corresponding to the complete coding sequence of P2X7R was detected. P2X7R mRNA was detected in the whole brain, hippocampus, cerebellum, leukocytes and thymus but not in cerebral cortex, hypothalamus, spleen and kidney (Figure 2). All PCR products were cloned using the pGEM-T-Easy plasmid (Promega), selected in Top-10 bacteria (Invitrogen) by blue-white selection and tested by EcoRI digestion. Clones having fragments of the expected size were amplified and purified for sequencing. The sequence confirmed the identity of the 1785 bp clones as the complete coding sequence of wild-type P2X7R. Therefore, in all the tissues tested, wild-type P2X7R is expressed as a single transcript which includes the complete coding sequence. The presence of tissue specific isoforms is unlikely. These studies provide useful information about the

P2X7R mRNA expression and transcript processing. This information can be used to synthesize riboprobes for in situ hybridization, Northern and Southern blot as well as engineering cells for the overexpression of P2X7R.

EXAMPLE 5

P2X7R expression in the mouse brain

The expression of P2X7R was further studied by immunohistochemistry of serial sections of complete mouse brains using a polyclonal antibody directed against an internal peptide of P2X7R (Santa Cruz Biotechnology). The brains from stress-free mice were shock frozen, cut into 16 μ m slices and fixed with paraformaldehyde for 5 minutes. The sections were blocked for 30 minutes at room temperature with 1:10 horse serum. All antibodies were diluted in TBST buffer (Tris-buffered saline with 0.05% Tween-20). The first antibody was used in a dilution 1:200 and incubated overnight. All washes were performed with TBST buffer. As a secondary antibody, an anti-goat IgG biotinylated (Vector Laboratories) was used and detection was performed using the streptavidin-biotin-horse-radish peroxidase complex system (Vector Laboratories) in combination with diaminobenzidine. Slides were counterstained with toluidine blue using standard procedures. The same procedure in the absence of the primary antibody was performed as a negative control. As a positive control to test the Preservation of the tissue was verified with an antibody specific for the protein Patched1 (Santa Cruz Biotechnology). Patched1 was used as positive control since it stains all relevant brain structures and is not affected by stress or antidepressants. Very specific staining pattern was detected, consistent with the specific subcellular localization of P2X7R in brain cells. Negative controls were completely devoid of signal. Positive control with Patched1 showed identical signal intensity and distribution in all samples, indicating that all tissues were equally well preserved and processed.

Proceeding from frontal to caudal, P2X7R protein was observed in the glomerular layer of the olfactory bulb at low levels (Figure 3). P2X7R was also present at very low levels in a restricted area of the periventricular hypothalamic nucleus (figure 3).

Ependymal cells surrounding the lateral ventricles also showed a faint staining (Figure 3). A stronger signal was detected in restricted areas of the hypothalamus, where the signal was present in single cells of the polymorph layer, the lacunosum moleculare and the oriens layer (Figure 4). In more posterior areas of the hippocampus, the signal was present in the molecular layer, stratum radiatum and near the CA3. In a further caudal position, P2X7R was expressed in the subcomisural organ (Figure 4). Therefore, the basal P2X7R expression in the brain of stress-free mice is restricted to areas that had been previously associated with depression, stress, learning and memory.

EXAMPLE 6

P2X7R is modulated in mice treated with an antidepressant

Further validation of role of P2X7R in affective disorders was performed by examining its expression pattern in response to stress and treatment with antidepressant drugs. A treatment schedule which has been proven to produce antidepressant effects on the behavioural level was administered to mice which were characterized as antidepressant-responsive by using a variety of behavioural paradigms suitable to detect anxiolytic and antidepressant effects of classical antidepressants like the selective serotonin reuptake inhibitor paroxetine. Paroxetine was delivered by gavage to naive male mice over a time period of 28 days at a dosage of 10 mg/kg bodyweight twice per day. In parallel, a control group of mice was given vehicle solution (i.e. without paroxetine) using the same treatment regiment while a second control group of mice was left undisturbed and stress-free (i.e. untreated) during the same period of the experiments. At the end of the long-term treatment, part of the mice of each experimental group were tested in the dark/light box (test of anxiety behaviour) and in the Porsolt's forced swim test (test of depressive-like behaviour) to confirm the effectiveness of the treatment (Figure 5). Passive stress coping behaviour decreased after long-term treatment with the antidepressant paroxetine. The other part of the experimental groups (i.e. mice without test experience) were decapitated, brains rapidly removed and frozen at -80°C until usage.

The expression of P2X7R in the brains of mice under stress-free conditions, and mice under mild stress produced by the vehicle application, and mice under paroxetine treatment was evaluated using three different brains from each group. Serial slides from each group of animals were analyzed in parallel by immunohistochemistry using the same materials in order to produce completely comparable results. No significant change in P2X7R expression in the olfactory bulb was seen in response to stress or to paroxetine treatment (Figure 6). However, in the periventricular nucleus of the hypothalamus, paroxetine produced a slight inhibition of P2X7R expression (Figure 7). No significant change was observed in the ependymal cells from different brain areas (Figure 8). The most dramatic changes were observed in the hippocampus, where P2X7R was strongly inhibited by stressful handling whereas paroxetine treatment produced a marked stimulation above basal levels (Figures 9, 10 and 11). This effect was observed all along the hippocampus but was more evident in the polymorph layer near the dentate gyrus. In the subcommissural organ, P2X7R expression remained unchanged by the different treatments. Therefore, P2X7R expression is strongly regulated in two specific brain areas involved in depression and stress. Other brain areas, which showed low levels of P2X7R and are not directly involved in depression, did not show changes.

In the samples from mice treated with paroxetine and showing a strong P2X7R expression, it was possible to analyze the distribution of P2X7R in more detail (Figures 10 and 11). The P2X7R protein was not only present in cell bodies but also was clearly detected in projections innervating the granular layer of the dentate gyrus (Figure 12). This subcellular localization of P2X7R is consistent with a role in neurotransmitter release and long term potentiation.

Since some reports (Muria et al., Biochem. J. 288 (1992) ,897-901; Ferrari et al., FEBS Lett. 447 (1999), 71-75) suggest that chronic and high dose stimulation of P2X7R may cause apoptosis in some cell types, the hippocampus of the above described animals were analyzed for the co-localization of apoptotic cells and P2X7R expressing cells, in consecutive sections, using TUNNEL staining and immunohistochemistry. In correlative sections, only few apoptotic cells were detected and they were present along the granular layers of the hippocampus where no

P2X7R expression was observed (Figure 13). No significant differences in the numbers of apoptotic cells were observed between the different treatment conditions. Therefore, the location and number of apoptotic cells did not correlate with the location and number of cells expressing P2X7R and rules out an involvement of P2X7R in the induction of apoptosis in the hippocampus.

Thus, P2X7R expression is considerably restricted to specific brain areas involved in depression. Moreover, P2X7R expression is inhibited by stress and strongly stimulated by antidepressant treatment in these specific areas. Therefore, P2X7R fulfils all criteria required for the actions of antidepressants according to the highest standards in the field of depression research. In addition, these results suggest that modulation of function of P2X7R is associated with chronic stress, which serves as a model for several aspects of affective disorders.

EXAMPLE 7

The behavioural effect of P2X7R inhibition in mice

To demonstrate that P2X7R inhibition acts as a causative agent for affective disorders, P2X7R function was specifically inhibited in distinct regions of the brain without affecting any other brain function. This was achieved by delivering double stranded small interference RNA molecules (siRNA) into restricted areas of the brain.

According to the observed expression pattern of P2X7R in the hippocampus (Figures 9, 10, and 11) and the known involvement of the hippocampus in depression, the dentate gyrus (hippocampus) was selected as target region for siRNA application. Male, naive mice were bilaterally implanted with a guide cannulae (23 gauge, length 8 mm) by means of a stereotactic instrument. The coordinates, in relation to bregma, were -2.0 mm posterior, ± 1.0 mm lateral, and -1.0 mm ventral. Following a recovery period of 5 days, the mice were divided into three experimental groups: vehicle (veh), control double stranded RNA (control), and P2X7R specific double stranded siRNA (siRNA). Sequences used for P2X7R siRNA are 5'-GUGGGUCUUGCAACAUGAUCTT-3' and 5'-GAUCAUGUGCAAGACCCACTT-3'. Both sequences were annealed and injected together as a double stranded

RNA. On day 6 after surgery, mice were slightly anaesthetized with Isofluran and injections of siRNA were carried out. The concentration of the control and siRNA was 0.1 nmol/ μ l, and a volume of 1 μ l per side was infused using specifically adapted injection systems (30 gauge, length 9 mm). The anaesthesia for the infusion was of short duration and the mice were awake immediately or few seconds after the manipulation.

Once delivered into the brain the siRNA molecules specific for P2X7R were taken up by brain cells and specifically induce the degradation of the complementary P2X7R mRNA with high efficiency. As a result, P2X7R function was specifically inhibited for a short period without affecting any other brain function. In this regard, injection of vehicle or control siRNA did not result in any obvious changes in normal behaviour, i.e., food and water intake, or motor behaviour in the home cage.

The effects of P2X7R inhibition on depressive-like behaviour was assessed 24 hours and 48 hours after infusion of siRNA, control or vehicle according to the standard test paradigm, the Porsolt's forced swim test (Porsolt et al., Arch. Int. Pharmacodyn. 229 (1977), 327-336; Porsolt, Rev. Neurosci. 11 (2000), 53-58). The parameter used to evaluate depressive-like behaviour is the time the animal is floating in the water, a behaviour which is associated with behavioural despair as the animal does not make any effort to actively cope with the stressful situation. Compared to vehicle application, no influence of control double stranded RNA (5'-CAACUUCAUCUUCUACGCGTT-3') on floating behaviour (passive stress coping) was detected. In contrast, compared to controls, mice infused with P2X7R specific siRNA showed a significant increase in passive behaviour, which is construed as depressive-like behaviour. This interpretation becomes moreover evident when the effects of antidepressants on passive stress coping behaviour in the forced swim test are visualized (Figure 14). Passive stress coping behaviour increased after acute intrahippocampal (bilateral, dentate gyrus) of siRNA targeting P2X7R. The Porsolt's forced swim test is a standard test used to assess the effectiveness of antidepressants and it has been proven by many studies that the test is selectively sensitive for these effects, given that the right animal model is used. The paradigm has been widely used to test pharmaceutical compounds and to validate animal

models of depression, which show an increase in passive behaviour as do the mice where P2X7R has been inhibited (siRNA).

At the end of the experiment, the mice were sacrificed and the brains were examined to confirm the location and efficiency of the siRNA injections. For this purpose the brains were cut into sections and the slides were stained by immunohistochemistry using the above mentioned protocols. Brains from mice injected with the specific double stranded siRNA, with control double stranded RNA and with vehicle were examined in parallel. Under these conditions, the specific siRNA directed against P2X7R injected near the dentate gyrus induced on average an 80% inhibition of P2X7R protein expression as compared to the samples from mice injected with vehicle or with control double stranded RNA. Both the number of cells expressing P2X7R as well as the intensity of the expression were strongly reduced (Figure 15). The injections with siRNA did not produce any sign of local inflammation or infiltration at the hippocampus. Thus, P2X7R expression is specifically and locally inhibited by siRNA application in vivo. This inhibition produced behavioural changes indicating a causative role for P2X7R in affective disorders. These results in combination with those mentioned above support and confirm the observation of mutations in P2X7R being associated with affective diseases in humans and that modulation of P2X7R activity has antidepressive effects.

New EP-patent application
NeuroNova
Our Ref.: H1566EP

Claims

1. A nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of:
 - (a) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 5'UTR region corresponding to positions 362, 532, 1100, 1122, 1171 or 1702 of the genomic sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;
 - (b) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of the ATP-gated ion channel P2X7R, wherein in the exon as indicated in column "Exon" of the following Table A the amino acid residue as indicated in column "Amino acid residue" of Table A corresponding to the position as indicated in column "Position in wild-type" of Table A of the wild-type ATP-gated ion channel P2X7R amino acid sequence as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue

Table A

| Exon | Amino acid residue | Position in wild-type |
|---------|--------------------|-----------------------|
| exon 3 | R (Arg) | 117 |
| exon 5 | G (Gly) | 150 |
| exon 6 | E (Glu) | 186 |
| exon 6 | L (Leu) | 191 |
| exon 8 | R (Arg) | 270 |
| exon 13 | I (Ile) | 568 |
| exon 13 | R (Arg) | 578 |

- (c) a nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in exon 5 or 8 corresponding to position 32548 or position 37633 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1,

wherein at said position said nucleotide is replaced by another nucleotide

- (d) a nucleic acid sequence encoding a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted;
- (e) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R, wherein in the intron as indicated in column "Intron" of the following Table B the nucleotide as indicated in column "Replaced nucleotide" of Table B corresponding to the position as indicated in column "Position in wild-type" of Table B of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as depicted in SEQ ID NO: 1 is replaced by another nucleotide

Table B

| Intron | Replaced nucleotide | Position in wild-type |
|---------------|----------------------------|------------------------------|
| intron 1 | G | 3166 |
| intron 1 | C | 24778 |
| intron 1 | C | 24830 |
| intron 3 | A | 26308 |
| intron 3 | G | 26422 |
| intron 4 | G | 32394 |
| intron 4 | T | 32434 |
| intron 5 | A | 32783 |
| intron 6 | G | 35641 |
| intron 6 | A | 35725 |
| intron 6 | T | 36001 |
| intron 7 | G | 36378 |
| intron 7 | T | 36387 |
| intron 7 | G | 36398 |
| intron 9 | C | 47214 |
| intron 11 | T | 47563 |
| intron 12 | C | 54307 |
| intron 12 | G | 54308 |

- (f) a genomic nucleotide sequence encoding an ATP-gated ion channel P2X7R and which contains a mutation in the 3'UTR region corresponding to position 54925, 55169, 55170, 55171 or 55917 of the wild-type ATP-gated ion channel P2X7R nucleotide sequence as

depicted in SEQ ID NO: 1, wherein at said position said nucleotide is replaced by another nucleotide;

- (g) a nucleotide sequence comprising at least 20 or 21 nucleotides and comprising the mutations or deletions as defined in any one of (a) to (f);
 - (h) a nucleic acid sequence comprising a nucleotide sequence as shown in any one of SEQ ID NOs: 13 to 51;
 - (i) a nucleic acid sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NOs: 5 to 12;
 - (j) a nucleotide sequence which hybridizes to a nucleotide sequence defined in any one of (a) to (g) or to the nucleotide sequence of (h) and having a mutation as defined in any one of (a) to (f); and
 - (k) a nucleic acid sequence being degenerate as a result of the genetic code to the nucleic acid sequence as defined in (j).
2. The nucleic acid molecule of claim 1 derived from mouse, rat or human.
 3. The nucleic acid molecule of claim 1 or 2 which is DNA, RNA, PNA or phosphorothioates.
 4. A vector comprising the nucleic acid molecule of any one of claims 1 to 3.
 5. The vector of claim 4 which is an expression vector, a gene targeting vector and/or a gene transfer vector.
 6. A host transformed with a vector of claim 4 or 5 or transformed with the nucleic acid molecule of any one of claims 1 to 3.
 7. The host of claim 6 which is a mammalian cell, an amphibian cell, a fish, an insect cell, a fungal cell, a plant cell or a bacterial cell.
 8. The host of claim 7, wherein said mammalian cell is selected from the group consisting of CHO cells, HEK293 cells, COS-7 cells or PC12 cells.

9. The host of claim 7, wherein said amphibian cell is an oocyte, preferably a *Xenopus* oocyte.
10. The host of claim 9, wherein said oocyte is a frog oocyte.
11. The host of claim 6 which is a non-human transgenic organism.
12. The host of claim 11, wherein said non-human organism is a mammal, amphibian, a fish, an insect, a fungus or a plant.
13. A method for producing the polypeptide encoded by a nucleic acid molecule of claims 1(b) or 1(d) comprising culturing/raising the host of any one of claims 6 to 11 and isolating the produced polypeptide.
14. A polypeptide encoded by the nucleic acid molecule of claim 1(b) or 1(d) or produced by the method of claim 13.
15. An antibody specifically directed to the polypeptide of claim 14, wherein said antibody specifically reacts with an epitope generated and/or formed by the mutation in the ATP-gated ion channel P2X7R selected from the group consisting of:
 - (i) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein the R (Arg), G (Gly), E (Glu), L (Leu), R (Arg), I (Ile) or R (Arg) residue corresponding to position 117, 150, 186, 191, 270, 568 or 578 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 is replaced by another amino acid residue; and
 - (ii) an epitope specifically presented by a polypeptide which has an amino acid sequence of an ATP-gated ion channel P2X7R, wherein amino acids corresponding to positions 488 to 494 of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 3 or 4 are deleted.

16. The antibody of claim 15 which is a monoclonal antibody.
17. An aptamer specifically binding to a nucleic acid molecule of any one of claims 1 to 3 or to the polypeptide of claim 14.
18. A primer or pair of primers capable of specifically amplifying a nucleic acid molecule as defined in any one of claims 1 to 3.
19. The primer or pair of primers of claim 18, which is selected from the group consisting of SEQ ID NOs.: 52 to 111.
20. A composition comprising the nucleic acid molecule of any one of claims 1 to 3, the vector of claim 4 or 5, the polypeptide of claim 14, the antibody of claim 15 or 16, the aptamer of claim 17 and/or the primer or pair of primers of claim 18 or 19.
21. The composition of claim 20 which is a diagnostic composition.
22. The diagnostic composition of claim 21, optionally further comprising suitable means for detection.
23. Use of the nucleic acid molecule of any one of claims 1 to 3, the vector of claim 4 or 5, the polypeptide of claim 14, the antibody of claim 15 or 16, the aptamer of claim 17 and/or the primer or pair of primers of claim 18 or 19 for the preparation of a diagnostic composition for the detection of an affective disorder.
24. A method of diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2XR7 protein expressed in the cells of said individual is non-functional, shows an altered ATP-gating in comparison to

the wild-type P2XR7 protein or is over- or under-expressed in comparison to the P2XR7 protein level an unaffected individual.

25. A method for diagnosing an affective disorder or a susceptibility to an affective disorder comprising the step of determining in a sample obtained from an individual whether the P2X7R gene sequence or encoded protein thereof comprises a mutation in comparison to the wild-type P2X7R sequence.
26. The method of claim 25, wherein said mutation is a mutation as defined in claim 1 and/or a nucleotide replacement or deletion selected from the following Table C indicating in column "Region of P2X7R" the region of the P2X7R genomic nucleotide sequence in which the replacement or deletion occurs, in column "Nucleotide" of Table C the nucleotide which is replaced by another nucleotide or the nucleotides which are deleted and in column "Position in wild-type" of Table C the corresponding position in the nucleotide sequence of the wild-type ATP-gated ion channel P2X7R as depicted in SEQ ID NO: 1

Table C

| Region of P2X7R | Nucleotide | Position in wild-type |
|-----------------|------------|-----------------------|
| 5'UTR | T | 362 |
| 5'UTR | T | 532 |
| 5'UTR | A | 1100 |
| 5'UTR | A | 1122 |
| 5'UTR | C | 1171 |
| 5'UTR | T | 1351 |
| 5'UTR | G | 1702 |
| 5'UTR | T | 1731 |
| 5'UTR | C | 1860 |
| 5'UTR | C | 2162 |
| 5'UTR | C | 2238 |
| 5'UTR | A | 2373 |
| 5'UTR | G | 2569 |
| 5'UTR | G | 2702 |
| intron 1 | G | 3166 |
| intron 1 | C | 24778 |
| intron 1 | C | 24830 |
| exon 2 | T | 24942 |

| | | |
|-----------|--------------------------------------|----------------|
| exon 3 | C | 26188 |
| exon 3 | A | 26308 |
| exon 3 | G | 26422 |
| intron 4 | G | 32394 |
| intron 4 | T | 32434 |
| exon 5 | G | 32493 |
| exon 5 | G | 32506 |
| exon 5 | C | 32507 |
| exon 5 | C | 32548 |
| intron 5 | A | 32783 |
| intron 5 | T | 35309 |
| intron 5 | C | 35374 |
| intron 5 | A | 35378 |
| exon 6 | G | 35438 |
| exon 6 | T | 35454 |
| intron 6 | T | 35549 |
| intron 6 | G | 35641 |
| intron 6 | A | 35725 |
| intron 6 | T | 36001 |
| intron 6 | A | 36064 |
| intron 6 | deletion of GTTT | 36091 to 36094 |
| intron 6 | C | 36108 |
| intron 7 | C | 36374 |
| intron 7 | G | 36378 |
| intron 7 | T | 36387 |
| intron 7 | G | 36398 |
| intron 7 | C | 37439 |
| intron 7 | T | 37513 |
| exon 8 | C | 37604 |
| exon 8 | G | 37605 |
| exon 8 | G | 37623 |
| exon 8 | C | 37633 |
| intron 9 | C | 47214 |
| exon 11 | G | 47383 |
| exon 11 | C | 47411 |
| intron 11 | T | 47563 |
| intron 12 | C | 54307 |
| intron 12 | G | 54308 |
| exon 13 | C | 54399 |
| exon 13 | A | 54480 |
| exon 13 | C | 54523 |
| exon 13 | deletion of CCCTGAGAGCCACAGGTGCCT | 54562 to 54582 |
| exon 13 | A | 54588 |
| exon 13 | C | 54664 |
| exon 13 | G | 54703 |
| exon 13 | A | 54804 |

| | | |
|---------|---|-------|
| exon 13 | G | 54834 |
| exon 13 | G | 54847 |
| 3'UTR | G | 54925 |
| 3'UTR | C | 55169 |
| 3'UTR | A | 55170 |
| 3'UTR | A | 55171 |
| 3'UTR | C | 55917 |

27. The method of claim 26, wherein the occurrence of the mutation in the ATP-gated ion channel P2X7R gene is determined by PCR or immunological methods.
28. The composition of claim 20 which is a pharmaceutical composition.
29. The pharmaceutical composition of claim 28, optionally further comprising a pharmaceutically acceptable carrier.
30. A method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic acid molecule as defined in any one of claims 1 to 3 or a therapeutically effective amount of the polypeptide of claim 14 to a subject suffering from said disorder.
31. Use of the nucleic acid molecule of any one of claims 1 to 3, the vector of claim 4 or 5, the polypeptide of claim 14, the antibody of claim 15 or 16 and/or the aptamer of claim 17 for the preparation of a pharmaceutical composition for the treatment of an affective disorder.
32. A pharmaceutical composition comprising a nucleic acid molecule comprising a nucleotide sequence which encodes a functional ATP-gated ion channel P2X7R and which is selected from the group consisting of:
 - (a) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence as depicted in SEQ ID NO: 3 or 4;
 - (b) a nucleotide sequence comprising the nucleotide sequence as depicted in SEQ ID NO: 1 or 2;

- (c) a nucleotide sequence which hybridizes to the nucleotide sequence of (a) or (b); and
 - (d) a nucleotide sequence which is degenerated as a result of the genetic code to the nucleotide sequence of (c).
33. A pharmaceutical composition comprising a compound the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells or comprising a nucleic acid molecule the expression of which in cells or the administration of which to cells leads to a reduction of the expression of a nucleic acid encoding an ATP-gated ion channel P2X7R in the cells.
34. The pharmaceutical composition of claim 33 which is an antisense nucleic acid, a ribozyme, a co-suppressive nucleic acid, iRNA or siRNA.
35. A method of treating an affective disorder comprising administering a therapeutically effective amount of the nucleic acid molecule as defined in any one of claims 32 to 34 or a therapeutically effective amount of the polypeptide encoded thereof to a subject suffering from said disorder.
36. Use of the nucleic acid molecule as defined in any one of claims 32 to 34 for the preparation of a pharmaceutical composition for treating an affective disorder.
37. Use of a modulator of P2X7R activity for the preparation of a pharmaceutical composition for treating an affective disorder.
38. The use of claim 37, wherein said modulator is selected from the group consisting of piperidine and piperazine derivatives, adamantane derivatives, substituted phenyl compounds, brilliant blue, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate and 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate.

39. The use of any one of claims 23, 30 or 36 to 38 or the method of any one of claims 24 to 27 or 35, wherein said affective disorder is selected from the group consisting of major depression, generalized anxiety disorder and bipolar disorder.
40. The use or the method of claim 39, wherein said major depression is selected from the group consisting of major depression, dysthymia, atypical depression, premenstrual dysphoric disorder and seasonal affective disorder.
41. The use or the method of claim 39, wherein said generalized anxiety disorder is selected from the group consisting of panic disorder, phobias, agoraphobia, social phobia, specific phobia, obsessive-compulsive disorder, post-traumatic stress disorder, separation anxiety disorder, mania, hypomania and cyclothymic disorder.
42. The use or the method of claim 39, wherein said bipolar disorder is bipolar disorder type I or bipolar disorder type II.
43. A kit comprising the nucleic acid molecule of any one of claims 1 to 3, a vector of claim 4 or 5, a host of any one of claims 6 to 11, a polypeptide of claim 14, an antibody of claim 15 or 16, an aptamer of claim 17 and/or a primer or pair of primers of claim 18 or 19.
44. A method for identifying compounds which are capable of specifically interacting with the polypeptide of claim 14, comprising the steps of
 - (a) contacting a polypeptide of claim 14 with a compound or a candidate mixture of compounds to be tested; and
 - (b) determining whether said compound or a candidate mixture of compounds is capable of specifically interacting with said polypeptide.

45. A method for the characterization of compounds which are capable of altering characteristics of the polypeptide of claim 14, comprising the steps of
- (a) contacting a polypeptide of claim 14 with a compound or a candidate mixture; and
 - (b) determining whether the compound or a candidate mixture alters a characteristic of the polypeptide of claim 14.
46. A method of screening for compounds which are capable of interacting with the polypeptide of claim 14, comprising the steps of
- (a) contacting a polypeptide of claim 14 with a compound or a candidate mixture of compounds;
 - (b) measuring and/or detecting a response; and
 - (c) comparing said response to a standard response as measured in the absence of said candidate molecule.
47. A method for the production of a pharmaceutical composition comprising the steps of the method of any one of claims 41 to 45 and comprising a further step, wherein a derivative of said identified, characterized and/or screened molecule is generated.
48. A method for the production of a pharmaceutical composition comprising the steps of the method of any one of claims 44 to 47 and formulating the molecules identified, characterized, screened and/or derivatized in pharmaceutically acceptable form.
49. The method of claim 48, wherein the pharmaceutical composition to be produced further comprises neuroprotective substances, nootropic substances, brilliant blue, piperidine or piperazine derivatives thereof, adamantane derivatives, substituted phenyl compounds, oxidized ATP, 2-O-(4-benzoylbenzoyl)adenosine-5-triphosphate or 3-O-(4-benzoylbenzoyl)adenosine-5-triphosphate.

50. The method of any one of claims 44 to 49, wherein said compound(s) or candidate mixture(s) of compounds comprise(s) antagonist(s), partial antagonist(s), partial agonist(s) and/or agonist(s) for an altered ATP-gated ion channel P2X7R.
51. A method for diagnosing an affective disorder of an individual comprising:
- (a) isolating DNA from cells obtained from an individual;
 - (b) determining all or part of the nucleotide composition of the P2X7R gene; and
 - (c) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism(s) mutation or allelic variation.
52. A method for diagnosing an affective disorder of an individual comprising:
- (a) isolating RNA from cells obtained from an individual;
 - (b) converting said RNA into cDNA;
 - (c) determining all or part of the nucleotide composition of the P2X7R gene; and
 - (d) analyzing said nucleotide composition of P2X7R for the presence of one or more polymorphism(s), mutation or allelic variation.
53. A method for diagnosing an affective disorder of an individual comprising:
- (a) isolating RNA or protein from cells obtained from an individual ;
 - (b) determining the levels of P2X7R RNA or protein; and
 - (c) comparing the levels of P2X7R RNA or protein with the corresponding levels from a normal individual not afflicted with an affective disorder.

Abstract

The present invention relates to nucleic acid molecules, preferably genomic sequences, encoding an ATP-gated ion channel P2X7R which contain a mutation in the 5'UTR or 3'UTR regions, a mutation in exon 3, 5, 6, 8 or 13 or in introns 1, 3, 4, 5, 6, 7, 9, 11 or 12 or a deletion in exon 13, which allow to diagnose affective disorders. The invention further relates to polypeptides encoded by said nucleic acid molecules vectors and host cells comprising said nucleic acid molecules as well as to methods for producing polypeptides encoded by said nucleic acid molecules. The present invention also provides antibodies specifically directed to polypeptides encoded by said nucleic acid molecules and aptamers specifically binding said nucleic acid molecules.

Additionally, primers for selectively amplifying said nucleic acid molecules are provided in the present invention as well as kits, compositions, particularly pharmaceutical and diagnostic compositions comprising said nucleic acid molecules, vectors, polypeptides, aptamers, antibodies and/or primers. Moreover, the present invention relates to methods for diagnosing affective disorders associated with a non-functional P2X7R protein, an altered ATP-gating of the P2X7R protein, an over- or underexpression of the P2X7R protein or associated with the presence of any one of the aforementioned nucleic acid molecules or polypeptides encoded thereby. Additionally, the present invention relates to uses and methods for treating affective disorders employing a functional or non-functional ATP-gated ion-channel P2X7R.

The present invention also relates to uses of modulators of P2X7R activity for treating affective diseases.

Furthermore, the present invention also relates to methods for identifying and characterizing compounds which are capable of specifically interacting with or altering the characteristics of the polypeptides of the present invention as well as to methods for the production of pharmaceutical compositions.

5,2Mb

1,6Mb




Figure 1a




Figure 1b




Figure 1c




Figure 1d




Figure 1e




Figure 5

P2X7v01 MPACCSVDVFQYETNKNVTRIQSMNYGTIKWFFHVIIIFSVCFALVSDKLYQRKEPVISS
P2X7v04 MPPVD-----AFCLPFS---FALVSDKLYQRKEPVISS
P2X7v02 MPACCSVDVFQYETNKNVTRIQSMNYGTIKWFFHVIIIFSVCFALVSDKLYQRKEPVISS
P2X7v03 MPACCSVDVFQYETNKNVTRIQSMNYGTIKWFFHVIIIFSVCFALVSDKLYQRKEPVISS
1.....10.....20.....30.....40.....50

P2X7v01 VHTKVKGIAEVKEEIVENGVKKLVHSVFDTADYTFPLQGNSEFFVMTNFKTEGQEQRLCP
P2X7v04 VHTKVKGIAEVKEEIVENGVKKLVHSVFDTADYTFPLQGNSEFFVMTNFKTEGQEQRLCP
P2X7v02 VHTKVKGIAEVKEEIVENGVKKLVHSVFDTADYTFPLQGNSEFFVMTNFKTEGQEQRLCP
P2X7v03 VHTKVKGIAEVKEEIVENGVKKLVHSVFDTADYTFPLQGNSEFFVMTNFKTEGQEQRLCP
61.....70.....80.....90.....100.....110

Figure 16a

```

P2X7v01 EYPTRRTLCSDRGCKKGWMDPQSKGIQTGRVCVVHEGNQKTCEVSAWCPIEAVEEAPRPA
P2X7v04 EYPTRRTLCSDRGCKKGWMDPQSKGIQTGRVCVVHEGNQKTCEVSAWCPIEAVEEAPRPA
P2X7v02 EYPTRRTLCSDRGCKKGWMDPQSKGLLS-----
P2X7v03 EYPTRRTLCSDRGCKKGWMDPQSKGIQTGRVCVVHEGNQKTCEVSAWCPIEAVEEAPRPA
          121.....130.....140.....150.....160.....170

P2X7v01 LLNSAENFTVLIKNNIDFPGHNYTTRNILPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
P2X7v04 LLNSAENFTVLIKNNIDFPGHNYTTRNILPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
P2X7v02 -----
P2X7v03 LLNSAENFTVLIKNNIDFPGHNYTTRNILPGLNITCTFHKTQNPQCPIFRLGDI FRETGD
          181.....190.....200.....210.....220.....230

```

Figure 16b


```

P2X7v01 NFSDVAIQGGIMGIEIYWDCNLD RWFHHCHPKYSERRLDDKTTNVSLYPGYNFRYAKYYK
P2X7v04 NFSDVAIQGGIMGIEIYWDCNLD RWFHHCHPKYSERRLDDKTTNVSLYPGYNFRYAKYYK
P2X7v02 -----
P2X7v03 NFSDVAIQGGIMGIEIYWDCNLD RWFHHCHPKYSERRLDDKTTNVSLYPGYNFRYAKYYK
      241.....250.....260.....270.....280.....290

P2X7v01 ENNVEKRTLKVFGIREFDILVEGTGGKFDIIQLVVYIGSTLSYFGLAAVFIDFLIDTYSS
P2X7v04 ENNVEKRTLKVFGIREFDILVEGTGGKFDIIQLVVYIGSTLSYFGLAAVFIDFLIDTYSS
P2X7v02 -----
P2X7v03 ENNVEKRTLKVFGIREFDILVEGTGGKFDIIQLVVYIGSTLSYFGLVRDLSFHALGKWEG
      301.....310.....320.....330.....340.....350

```

Figure 16c

| | | |
|---------|---|--|
| P2X7v01 | NCCRSHIYPWCKCCQPCVVNEYYRKKCESIVEPKPTLKYVSFVDESHIRMVNQQLLGRS | |
| P2X7v04 | NCCRSHIYPWCKCCQPCVVNEYYRKKCESIVEPKPTLKYVSFVDESHIRMVNQQLLGRS | |
| P2X7v02 | ----- | |
| P2X7v03 | EGSD----- | |
| | 361.....370.....380.....390.....400.....410 | |
| P2X7v01 | LQDVKGQEVPRPAMDFDLSRLPLALHDTPPIPGQPEEIQLLRKEATPRSRDSPVWCQCG | |
| P2X7v04 | LQDVKGQEVPRPAMDFDLSRLPLALHDTPPIPGQPEEIQLLRKEATPRSRDSPVWCQCG | |
| P2X7v02 | ----- | |
| P2X7v03 | ----- | |
| | 421.....430.....440.....450.....460.....470 | |

Figure 16d

| | |
|---------|--|
| P2X7v01 | SCLPSQLPESHRCLEELCCRKKPGACITTTSELFRLVLSRHVLQFLLLYQEPLLDVDS |
| P2X7v04 | SCLPSQLPESHRCLEELCCRKKPGACITTTSELFRLVLSRHVLQFLLLYQEPLLDVDS |
| P2X7v02 | ----- |
| P2X7v03 | ----- |
| | 481.....490.....500.....510.....520.....530 |
| P2X7v01 | TNSRLRHCAIRCYATWRFSGQDMADEFAILPSCCRWRIRKEFPKSEGQYSGFKSPY |
| P2X7v04 | TNSRLRHCAIRCYATWRFSGQDMADEFAILPSCCRWRIRKEFPKSEGQYSGFKSPY |
| P2X7v02 | ----- |
| P2X7v03 | ----- |
| | 541.....550.....560.....570.....580.....590 |

Figure 16e

SEQUENCE LISTING

<110> NeuroNova AG
<120> A Method for Diagnosing and Treating Affective Disorders
<130> XXX
<160> 111
<170> PatentIn version 3.1
<210> 1
<211> 56580
<212> DNA
<213> Homo sapiens

<220>
<221> exon1
<222> (3000)..(3124)
<223>

<220>
<221> exon2
<222> (24841)..(25009)
<223>

<220>
<221> exon3
<222> (26134)..(26202)
<223>

<220>
<221> exon4
<222> (30958)..(31030)
<223>

<220>
<221> exon5
<222> (32481)..(32577)
<223>

<220>
<221> exon6
<222> (35416)..(35496)
<223>

<220>
<221> exon7
<222> (36113)..(36242)
<223>

<220>
<221> exon8
<222> (37541)..(37677)
<223>

<220>
<221> exon9
<222> (45470)..(45560)
<223>

<220>
<221> exon10
<222> (47229)..(47295)
<223>

<220>
<221> exon11
<222> (47380)..(47529)

<223>

<220>

<221> exon12

<222> (50438)..(50539)

<223>

<220>

<221> exon13

<222> (54392)..(54889)

<223>

<400> 1

tgtgtccca gctactcagg aggtgtggt agggggatca ctggagccca gaagtttgaa 60
 gctgcagtga gctatgattg cgccactgca ctccagcctg gacgacagag caagaccag 120
 tctctaaaat aaattaatta aattaatttt taaaaagaa agataagata aagagtcata 180
 gaagtacaaa tgagaaacca ggagaacatg ggaccagga aagtcattga aaccaaggga 240
 aaatactggt ccaagatgca gaaagtagcc agtgatgttg aatgcatcca agagatcaag 300
 taacataagg aataaaatat atccaccag tctggcaact gagaagtctt tggtaacctc 360
 attgagaaca gaatccatgg agtggaggag gtagcagtga tgaaatagcg gacaagggct 420
 gaaggagtgg agaggggaga gcatgaggga ccagtgcag agtgcagcag aaagacaaca 480
 aacaggccgg gtgtagtggc tcacgcctgt aatccaaca ctttgtagg ctaaggcggg 540
 cagatcgctt gtgaccagga gtcgagacc aacctagcca acatggtgaa accccatctc 600
 tactaaaaac acaaaaatca gctgagcgtg cgcatggctg taattgcaga tactcatgag 660
 gctgaggcac aagaatcgct tgaaccaggg aggcagggggt tgcagtgacc caaagtcttg 720
 ccactgcact acagcctggg caacaggggg agactccatc tcaaaaaaac aaaaaaaaaa 780
 aagaaagaaa gaaaagacaa agaagaaaag aaagaaagaa agaaagaaag aaagaaagaa 840
 agaaagaaag aaagaaagaa agaagaaaag agaaagaagg aaagaaagac aagaaaatag 900
 atgtctttga gtggtagcaa gagcaacact gaagtgatt gcctattctg gacatttctt 960
 ataatggag tcgtaaaaga tgtatccttt tgtgtctggc ctctttact taagcgtagt 1020
 gtttcaagg ttcattcatg ttggagcatg ggtcagtgtt tcattccttt ttatggctga 1080
 ataattatcc ctgtatgga ctttcgtgct ttgtgtatc catcatcag ctgatggaca 1140
 ttgacttgt ctgcctgttc actgccatat ctactattcc tgggtctctt gggaatgagt 1200
 gaatgaatag cataaacaga gctgtccaag gtcacagagc tggtaatgag tgaagccagg 1260
 atcgggaagcc aggccattag tcccctagag cctatgttct aagcatcagg ctttacctgt 1320
 gaatctcttc ttttacaga tgaagatgac tgtatcactc agattcccgg caggaaagca 1380
 atggcactact caagtggggg aactaatgat ggaaccattt acaagggtgt ggacagagtt 1440
 aagaaaaagc aataggagat agtgagcttc ctggggctgg taagagtggg gagcccttac 1500
 cactcccagg actaaaggag ggagtggtgc ccagaagccc tgcctatatg caactagaaa 1560
 gggcagggcc agggagtac gtccatctc actgctctcc agtctcctga actggaagcc 1620
 agaaggtgag gggaaccctg atgcagtttg tatgtgtgag aaagtacaat tagtttagac 1680
 tgaaaaactg aaaatctacc cggccactta gcaggctgga ataacagaaa tggatcaagc 1740

cagctgtaaa gataacaggg aacaataatt ctctgtagct gtaaagtgat aatacaacce 1800
 tgcacttttg agtgactgct gaaacattgt cctttaaaat cagagacctt cagaaacttc 1860
 gctgtttgaa attacatgac taagactgaa atattccaat ttgcctgga agatttaagt 1920
 catcttgaca cagagaagca gcctcaattt acaactcagg agcagagcct cagataaaga 1980
 tttctggac acatttgaca tgtatcttag ctatgttgct tcctaggaaa cagggccctg 2040
 ggtcctcttt gcaatccaga ctgaagtga ctgctttgta caaacctgtt ttgctttgag 2100
 tccatcaaaa catgacttca ttgatttt atctcaactc cactttcctc ggaatcctat 2160
 actaaattgc tgtttcctt tgtttggtga tgtgcgtagc tcttcgggtg ggtggtgtcc 2220
 ctactgaat aggtaacaa acctaacttt gttggactgc cactgtgtcc ctggtgatct 2280
 ttggctgatt ggtctaggtc atagatcgac ctgccggggt gcagaggagg gtggagagta 2340
 actcagaggg tcaagcatga aagatctggc agaaaaataa agccctcca cccccaccac 2400
 ccctaccctt gcaaacttga ttccccccac caactgcaga ccagagtatt ataaggggag 2460
 gtggaagaag agggggagat ctcatctac ccagagctcc tatacatcag gggctgaata 2520
 aaggggttga gaaatgaatg aatcaatctc tgagtggggc ttcaggcagt gaaagatct 2580
 cagtcctttt ctgaggcata atggaagctc ccagtcttgt gacatttga aggctgcccc 2640
 tttctccaa gagacatgag accaaaaaag tgaaaggaaa ggggggaaaa gggagaattc 2700
 tgaaatgcc catctctga acaccatctt tgttaggca tctgggggag gccagctggg 2760
 gtgaggtcat ctgccagcca ggcccgtagg acttggcgct tctgtttat cacagccaca 2820
 tgtggggcca ctgccagggc ccgcccacac tctgcagtca ttggaggagc ttgaagtaa 2880
 agactcctgc taaaaaccag tacgtttcat ttgcagtta ctgggagggg gcttctgtg 2940
 gccctgtcag gaagagtaga gctctggtcc agctccgcgc agggaggagg gctgtcacca 3000
 tgccggcctg ctgcagctgc agtgatgtt tccagtatga gacgaacaaa gtcactcgga 3060
 tccagagcat gaattatggc accattaagt ggttctcca cgtgatcatc ttttctacg 3120
 ttggtaagt gggatctggg gaggaccag atctctgcag tggccgacag cacagaaagc 3180
 cccagcgggc agcttcaggt gcacattctg aatctcatat ggttttcgaa tctgagacgt 3240
 gctctcacag ccagctgggc gggaggagg aagcagcagc aggcaagagg aaacggtgcc 3300
 aggtgcagc agagagaagc cacaggacaa gcgggattcc tttctgtct acttcaggcc 3360
 cgccagggcg cgcagggcag ggcgtgcctg gggaaggtag gaaagcgag ggcaacacc 3420
 tggatccca gggaggaggc gaggatctca gggcacgcci ggtgatcatg ctggcatctg 3480
 agtcaccatg ctggggagga ataggaccag gcttgaaaat gtgtataac tttagtcct 3540
 caccaacgtc aggaaggccc tgcttttgg tttttttc ttctaaaaga aacttactga 3600
 gatataatt ataccata caattgacct atttaaaggg taccatttaa tgattttcag 3660
 attattcca gagtttgca accatccca aatcaattt tagaatatt taatcgactc 3720
 aaaaggaatc ccacactcct tcaccatcat ttccaact gtctctctc ctccaccc 3780
 atcaatttcc tttctgctc tatggattg ccgattctgg acatttcata taaatggaat 3840

cacataatat gtggctcttt gtggcactta gcgtgttttc aatgctcatc catgtttag 3900
 catgtttag tacttcattc aatttttt tttaaagaga cggggcttca ctatttgc 3960
 caggctggtc tcaaaactcct ggactcaagt gatctgectg cctcggcctc ccaaagtgc 4020
 aggattacag gcgtgagcca ttgcacccgg ctgatacttc attcctttt acggctgagt 4080
 agtactccat tgcatggata gaccactttt ttctatccat tcatccattg atggacattg 4140
 ggggtgtttc tctttttgg ctatcatgaa taatgccaca tgaacatttg tgtacaaggt 4200
 ttatgtgga tatatatctt cctttcttc gaatatgtac ctaagagtaa aaattgtag 4260
 gtcatatgtt aactatgttt cacctttggg gggaatgtgg agctgaattt cacagcagct 4320
 gcagttttt acattcctat cagcagagta tgagggatcc aatttctcca catcctcacc 4380
 aacgcttgtt atcgtctgtc ttttgggtt ttgttgtgt cattttgtt ttgtcttga 4440
 gatgaagctt tgctctgttg cccaggctgg agtgagtggt cgcaatcttg gctcactgca 4500
 acctccacct ccccggttca agcgattctc ctgcctcagc ctcatgagta gctgggatta 4560
 cagggtgtgc tcaccactcc tcactaattt ttgtatttt agtagagatg aggtttcgcc 4620
 atgtaggcca ggctgtgttc aaactcctga cctcaagtga tccgcccacc ttggcctccc 4680
 aaagtgtgtg gattacaggc atgagccgcc acgcccggct gattgtctgt ctttttatt 4740
 atagccatgc tagtgggtgt gaagtgttag ttcatgttg ttgtgattg aatttcctg 4800
 atgggtgagt cctcttattc tctgtctga ttgataatga tgatgaaggc aatttgtatc 4860
 tatagagtgg cagtgtagtt tactaagagt tagggtaact tatttcatag tactggctat 4920
 gtcttctggg ccaagtcatt aacttctctg agcctcagtt tctgcatctg ttcatagggt 4980
 tgtggcaatt aaccaaaaaa aaaaggcatg aacagccctt atcatgatga ctgacatagg 5040
 ataagagctc cataactagt atctatttt aaaaataatc ttttaagtc tgggagtgtt 5100
 ggctcacacc tgtaatccca acactttggg aggccgaggc ggggtggatca cgaagtcagg 5160
 agtttgagac cagcctggcc aatatgttga aaccccatct ctactaaaaa taaaaaatt 5220
 agtggggagt ggtgtgtcac acctgtaac ccagctacta gggaggctga ggcaggagaa 5280
 tcgctgaac ccggaggcgg aggtgtcagt gagccgagat caagccactg cactccagcc 5340
 tgggtgacag agcaagactc catctcaaaa taataataat agtaataatt ttttgatta 5400
 tataatagta tatatgtata taaaatacat gtatgtattt ttatctatat cctctgctct 5460
 gaccctcaaa gtaaccacgt ccaagttcag gatttgaat ctggaaacgt ggattcaaaa 5520
 atccttcacc tctttgagcc ttggtttcat catctgtaa atggggagaa ttgttgatg 5580
 gaatattaaa tgaactaata aatgcaaagc tgtttgagaa atatatggca tatagtaatc 5640
 cctgattaag tgtagttctt tattattaat aatgtatta ttaggattat tattattcga 5700
 ttcatatgtt tactgttcaa caaatattga atgataaaca tatatgctgg gtccggcatg 5760
 gtggcccatg cctgtaattc cagcactttg ggaggccaag gcgggcaggt cacttgaggt 5820
 caagagtgtg agaccagcct ggccaacgtg gtggaaactc cgtctgtgct aaaaatacaa 5880
 aaattagccg ggcattgttg tgggtgcctg taatcccagc tactcgggag gctgagacag 5940

gagaatcact tgaaccagg aggtggaggt tgcagtgagc caagattgca ccactgcact 6000
 ccagcctgag ccacagagca agactctgtc tcaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 6060
 tatatatata tatatatata tatatatata tagtattttt agtagagatg gggttttgcc 6120
 atctcttata ttttttata tttttatatt ttttatatat ttatatatat ttatatgtat 6180
 ttatatatt tatatatatg tgcagaaatc taaaataaga catatcactt aaaccagcta 6240
 tcacaggacc aatggtaaag agcataatta gtaaatgccc ccttctgcat taatacgtag 6300
 attgtatgtc acaatttcat gtagcagggtg atgggggttc ctgacattct tgggccacagc 6360
 atacaactct tttatttctt ttttagagac aggggtctcat tgtgttgccc taggctggag 6420
 tgcagtgtca ttatcacagc tcgccacagc ctcaacttct caggctcaag tgatcctccc 6480
 gcctcatcct cccaagtagc tgggaccata ggacacagcc atgatgtctg actaattttt 6540
 tttttttt tgtggagatg ggggtctccc tatgttgctc aggctagtct caaactcctg 6600
 gactcaagt atcctcctgc ttggcctcc caaagtgtg ggattatagg catgagccac 6660
 ttgtctggc ctacaattcc ttttttct ttttttct tttttttt tttttagt 6720
 gagtctcact ctgtcaccta ggctgtatg cagtgggtgt atctcagctc actgcaacct 6780
 atacctccca ggcccaagca atcctccac ctcaacctcc cgagtgtg ggaccacagg 6840
 tgtacaccac cagcccccagc taatttttg ttttttgg agagacaggg ttccaccata 6900
 gttgtccagg ctggcttga atcctgagc tcaagcaatc caccgcctt ggctcccaa 6960
 agtgctggga ttacagggtg gggtcaccat gccagccct caatttctc agattattgt 7020
 ataataataa aatttctaaa aacagattac tatgagaaca cctgtaatg ataaatatt 7080
 tttttcaca tggtgaaatc atacttaca tgtatttaac tccctcaaat tcagtatac 7140
 atgttccatg gggacatgga gagagagaga acaatataaa taatgcaatg attcagcaac 7200
 cgggttcgct ggagtgggtg tgagatggga tactggcatc tgggttccc tccgtcaatc 7260
 tctgttccc cgtgttttt agcgagcatc ttgctgtct acttattaat ctaggatcat 7320
 attttcagc cacatgaccc ctaaaataag tcaactcctt caccttcacc tggagcaaaa 7380
 ctgtaaaaac agcaacttgt tctagatctg gaggcaaaat tggttccagc agaccactg 7440
 ggaagtggga ggtgtttgt atgctttgt gacaatttaa gtgttataa ggcaatttg 7500
 atgagccatg atttcatct taagtccaa cttagaaga taaccttcat ataaaagttt 7560
 ctctacagt tttattttaa aatataaaat ggaagaaaa actatcacca gctttgctt 7620
 tttctctg gtaatggac tatctattaa cgtctcacta aatggccaaa tggagtctca 7680
 cagtacacag ttgggaaat gctgccttaa gtgtctgt gaagccttaa gttcgtgtaa 7740
 ccatgactta gccatctgtg aatgggtag acctatggg cataggaatt ggggaatgta 7800
 tttgcatca agtaggactg gtgtggacca caggtggaaa tccagacagg ctgagcaat 7860
 gttcaggata cagcagcaag tgacagggtc tggccagctg tccctactgg ggctgcaagg 7920
 actaagagcc aggataagga gacaggccct cggttattga aatgggata caaagcaggg 7980
 actggttatg gcggaaatta accaagaggc tgttcagctt ctctattct cacaaccaat 8040

tcagatctgg attcaaaggc agggatgcag ctacaagagg agtttgtgtt ggaggtctgt 8100
 gtagatagga tgctaagcag gtctgtgtca gccaggatcg ttgggcagt aagtacaga 8160
 aacccaactt acgttaacca caaagcctca aaggggaatt tattggatca tgtaactggc 8220
 cagttgaaga gatggtgctg tcctcggacg tggatgaaga taagagtca aactgtgtca 8280
 tcaggacgga tccgtctccc tacatcctgg ctccatttgc ttctatttgg ctctcttctt 8340
 gaacaactct gtatgtacaa gatatcctca cattctcttc tttaacaac ctacagaggag 8400
 caaacaagtc ttctctgata gctctgctaa gaaagtccca ggaagactc tgattgatca 8460
 gatgtgagtt acagatccat cctgaacca atcactgigg ccagggccaa ggtgtcatcc 8520
 cattggccag cctgggtcat gtgctggctc ccgtagcagg aagtagggag aagacgaggg 8580
 agccgtgtga gccctacgtg aaccacatcc agtggatttt ccttaggaag gaagattctg 8640
 tgaccagacc aaatggaaaa ggaagtgggg agggtaggca gacaaaaatc ggtgatctat 8700
 tacaaggctt ttcaaaaagc cagaaattag ggatttctaa gtgtacctct ctatgcctgg 8760
 ctgtctcag ccctgcagct gggctgtgga gactccatca gcaattgctg accccctatt 8820
 ctacctcgg ccccttcctg agtcccacca ggagccctgc tctcatctg ggacatgctt 8880
 ctgggccc at ggaggttgcc agtccactt ggtactttac tactatttgg ttctgtgac 8940
 caccaagctg ttgcttaaac caagccgcct atggtggcac cactttggag aattctttt 9000
 ttttttga gacaggttct cactctggtt accgagactg gagtgcagtg gtatgatctt 9060
 ggctcactgc agcctcgacc tcccgggttc aggtgatttt cccacctcag cctccccagt 9120
 agctgggact attggtgcat gccactacac ccggctcatt ttgtatttt tagtagagac 9180
 ggggtttgc catgttgcc gggctgtgtc tgaactcctg acctcaggtg atctgcccc 9240
 ctgggcctgt caaagtgtg aaattacagg cgtgagccac cgcacccggc tggataattc 9300
 tttagattt attccaagt gggtgagaag ccattgggaag attgtaagca aaggaataaa 9360
 agttggtgc attctcttg gaggatttcc ctctccctg gagaccagtc ctacagaag 9420
 cttttcaat ggccacagaa gactaaggaa gattgagaga ggcacaggaa acaattccac 9480
 tcagagctgc tgcttcgcca ctgcccagtc agcctccact ggccccaaca cgagcctatg 9540
 ctgtcatcc tgggcccagg ctggctgtcg ctgtggagat ggtgctgctg tgctagtgtg 9600
 tggctgttaa aatggagctc ctggctggac acctgtttcc cagtgggtgg ccacaaacct 9660
 gtgagtctgg gcctcctcca gctgagtaga agtgagccca gcacagaggc cggggaggcc 9720
 ccagacgcca ggaggagcaa atccaaatcc aaccaagca tggccctct gacctgacc 9780
 atcttttt tttttttt tttagagaca gagtctcct ctgtcaccca gactggagtg 9840
 cagtggcacg atctggctc attgaaccc ccacctctg gttcatgctg attctctgc 9900
 ctacgctcc ctataggga ctacaggcat gcaccactac accagataa tgtctgtatt 9960
 tttagtag agtgggttct gctatgttg ctaggctgt ctgaactcc tggcctcaag 10020
 cgatccatcc gtcttggtt cccaaaatgc tgagattaca ggtgtgagcc actgcgcca 10080
 gcctgcccct gactgccttc agaaaagtag aagcaggagg tctgccat tctcatcca 10140

tccttgacac ctgggtgatg tgctagagct tgaggggaag gtcagaggta acccagttcg 10200
 ggaagtgacc ccgtaggagg agaagaatcc agccacgtga cacacagtac gaagagaaca 10260
 gtaaatgcaa agacctggaa acgaggaaca gcaaggacag catgtcccag aacctgggga 10320
 gagaggaggg agcagccgga gctgagggca gctcagggga agagttgcta ttgattttc 10380
 agtgcattggg aagccactga aaagttttat gcagagaagt aaaaccatct cactcatgtt 10440
 tctagaagcc ttatgggcca ctgtgcagag catagagtgt ggtgggacaa gaatgacggt 10500
 gaggaaccag tgagaagggt cctggagtca ccagatgcga ggagatagag gcttgggcta 10560
 gaatgggctg gcggagctgg agagagggtg acaaagctga gatgtgttt ggaggcagaa 10620
 atgatatatt tgccaggagg tgaagatgg ctctagattc ctgagggaacg atgatgcctt 10680
 ttaccaacct gaagaagacc aaggagaggc ttgaggacta acacggaact gtgaaggagg 10740
 ccaggcaaa gttggagactg ccgtgagctg tgatcacgcc actgcaccct cgcctgggtg 10800
 acagactgag accctgtctt aaaatatata tgtaataaaa aataataaat aaagggccag 10860
 gcgtgtgtgg tcaacgcctg taatccaac actgggaggc tgaagcaggt ggatcacctg 10920
 aggtcagaag ttcaagacca gcctgtgcaa catggcgaaa tccatctat aataaaaata 10980
 caaaaattag ccagacatgg tggcacatgc ctgtaatccc agctacttgg gaggctgagg 11040
 caggagaatc acttgaacct gggaggcaga ggtgcagtg agcctaaac acaccactgc 11100
 actccagcct gggcaacaga gtgagacccc ataaataaat aaataaatac ataaataaag 11160
 gatatcaggc aagcaattaa aacacctgcc tactaccatc tatccgcatc cttttttca 11220
 tgtctaacac caaaacacca aaatagctgg aaacatatca tgtaaaaga taagcatgat 11280
 tctttcaat gaatggctgt ggacttatga aaagctgcct ccgcgtctga tttctgtt 11340
 tcaaggggta atgatggagg ttctctcct gcacctgcag gcctccgggg atgctgcccc 11400
 agtgcagcct cccaccctgg gctggagcct ccagccccg ctgcatgggt agcacagtgt 11460
 tgggaagagg aaggagcagt ttgtctggct tttcacagg cccacagctg gccggtacag 11520
 tggcagagaa ggcgaggctg aggtggggct ggggaagcca gcaggggcct gatcatggag 11580
 cgccttgagc tgcatccac aggtactgag aaagcatgga gagctgtccc ccagcccca 11640
 gcactttatt agaaaaatatt gaaacataac agaaaacatg aagggtttta tgtcatatt 11700
 ttcattttag agagatcatt ctgctgaaa tataacacct actgagtact gctgaggact 11760
 taccatgtgc cgggttctgt tcttagcatg tgacatgtat tggcctattt gattctccta 11820
 cataatcctg gaagattggt tgtgggagtg ggtgatcatg acctgaacag cctaaccatg 11880
 tttagggaac actcigtgtt atgcgttcca ccacacaatt catgtcacag ctgaatgtcc 11940
 cagtttcct tgcagctagg acacaacggg agaccaagc ttccaaca tatgccccag 12000
 gcaacacttt gatacacaag gaataacact ggatgctggg tgtgcacaga attcaaattc 12060
 aggaatggag ggaagtactg ccccttgacc catacaatag gggatcccct tccttgacc 12120
 ccatggctta gagggcccca ctctgtccct ctgcagactg tgagctgccc tatggaatga 12180
 ggggtgtctg aggctgtatt aatttctatt tgttacacag cagtggaaac taatacagtc 12240

ccacaggcct aaatcaaggc gtcagcaggc tgagttcctg cctggagact ctggggaaaa 12300
 ttacitcca aactcattt tgtgttggc aaaattcaat cccttgaggc tgcaggactg 12360
 ggcttgcct ctcactggct gtcagcctgg ggccactctc agattctgaa gtctccac 12420
 attcctgcc acatggctct ctccatctcc aagcctgcta tgacacaacg catgttctc 12480
 atgttcaaa tctctgactt ctctccggg accagaagaa gaatactctc ttctttaag 12540
 gggctcatgt gattgggtca ggtaactct atcatatcat ggagtctcac cgtctgccc 12600
 aggtgggtct tgaacacctg ggctcaagtg atctctctgc gttggcctcc caaagtctg 12660
 ggattacagg cgggaactgc tgtgccagc caaggccagc atcttcagac ctctctgccc 12720
 tgttttaca tggcttctc ctcttctgt gtgtgtcaaa tctctctg cctctctat 12780
 aaggacgtgt gtaatgacac tcagggccca cctggatgac acagggtcat ctgccatct 12840
 caaatcgtt aactttggc aggtgcagtg gctcatgcct gtaatctcag cactttggga 12900
 ggctgagaca ggtggatcac ttgaggtcag gaggtaaga ccagcctggc cagcatggtg 12960
 aaacctgtc tctactaaaa atacaaaagt tagctgggca cgcacctgta gtcccagcta 13020
 ctcaggagac tgaggcaaaa gaatctctag aacctgggag gcagagactg cagttagcca 13080
 agattatgcc actgcactc cagcctggg gacagagtga gacttcatct caaaaaaaaa 13140
 aaaaaaaga tgactgggca cgggtggctca catctgtaat cccagcattt gggaggccca 13200
 ggtgggcaga tcacctgggt tcaggagtic attcaagacc agcctggcca acatggtgaa 13260
 agtctctct tactaaaaa aaaaaaata caaaaattac ccagatgtgg tggcagggtc 13320
 ctgtaatccc agctacttg caggccgagg caggagaatc acttaaacct gggaggcaga 13380
 ggggtcagtg agctgagatc actctgtgc actccagcct ggacgacaga gcgaaactcc 13440
 atctcagaaa aacaaattaa ccaggcatgg tctgctgctc ctgtagtccc agctactggg 13500
 aaggctgagg tgggaggatt gcttgatctt gggagggtga ggctgcagtg agctgtgtt 13560
 gtgccactgc actccagcct gggcgacaga gtgagaccct atctgaaaaa aaaaaaatcg 13620
 gccaggcgcg gtggctcacg tctgtaatcc cagaactctg ggagactgag gtgggtggat 13680
 cacctgaggt caggagttc agaccagcct gaccaacagg gtgaaacccc gtcttacta 13740
 aaaatacaaa aattagctgg gcatggtggc aggcacctat ataatccag ctactcggga 13800
 ggctgaggca ggagaatgc ttgaactcag gaggcagagg ttgcagttag ccgaggtcat 13860
 ggcatgac tccagcctgg tggcagagt agactccgtc tcaaaaaaa gaaagtctct 13920
 tttaact ttttgaggt ataactaca tatcagaaa tcaccggtt taaatataca 13980
 tttaacgac ttttagtaa ttcccaagt tgtgaacca tcatcacaat ccagtttag 14040
 aacattcat cagccagta agagccctca caccattaa cagtcactcc ccactccac 14100
 ttctctgg tgggtgtca acccaggtaa ccatgaatct attctctgcc tctggagatt 14160
 tgtctttt ggacaactt tctcttaaa ctacagagag tgagttctgt tgttagaac 14220
 tagaaacct gcccaataga atactaatat cctctgtt tgcaaggag ttagagtctc 14280
 gctctgtgc ccaggctgga gtgcagtggt atgatctgc ctacttga cctctacct 14340

ccggattcaa gcgattctcc tacctcagct tcctgagtag ctgggattac agacgtgcac 14400
 caacacacct ggctaatttt tgtactttta gtagagatgg ggtctcacca tgttgccaa 14460
 gctggcttg aactcctgac tcaaatgag ccaccgcct tggcctccca aagttctggg 14520
 attacaggca tgagccactg caccagccc taatatcctc ttattacaaa agaagaaaca 14580
 atgactcaga aagattaagt cacttaccca aggtcatgta gtcagtaagc aatagagctg 14640
 agactgaacc tcactccaaa gtctatgctc tcactaaagt attgcagtgt agagagtaga 14700
 ggggacaggg tgagactgga ggaagggagg gcatttagga gactattggg atgtccagat 14760
 ctgagagctg ttggcgacct ggactgggga ggtgatagag gcagaggaga gaagtagatg 14820
 aatttgagca tattaggaag tgaacagta ggatttgga actgaacggg tgagtggaca 14880
 agaaggagag gatgagtaaa agagtgcacc atgcttctgg cttgggcaac tgatgggtac 14940
 tggaataact tgctgggata ggaatgcag gaaaaacagc caatttgggg tggaagatga 15000
 ggtcaggact aaaggcattg agtttaggt gtctatggaa tgtacaaatg gagatgttga 15060
 atggatgggt ggatatacga atctgcagct gagatgtagg ttctatatc accagcatat 15120
 agacggcaac cagaaccctg gaagaagggt caaccgacca gcaagagaag atgtagatga 15180
 gaaagcaaaa gggcctagaa cagccccaa aggagtgtca accttaagg cgtgggagcg 15240
 cagaagtcac caaagattc tgaggagtag ccacggcgat ccagtgaag tcaggcaca 15300
 gtggactcct gcaagctaag ggaaggaggc gagggaagtg gggcatggc acagtcagt 15360
 tcaactgctg ctgagagttc caggaagggt aggtccatgg tgatctatt ggatttagtg 15420
 acaacgacat tgtgtttac ttggcaagag aggtttaga gaagggtgg ggtgggagc 15480
 cagactagag tgggtagagg agtgagtga tcaagatgcg gaagtggatg tgatgactcg 15540
 cctttctt ctctttct tctttttt tttttgaga tgcagcttc ctctgtgcc 15600
 caggctggag tgcgatggtg tgatctccac tcaccacaac ctctgcctcc tgggtcaag 15660
 tgattctct gcctcagcct tccaagtagc tgggattaca ggcattgtcc accacaccg 15720
 cctaatttt tgtatttta gtagagacgg ggttcacca tgttgccag gctggtctca 15780
 aactcctgac ctctgatcc acctgcctca gcctccaaa gtgctgggat tacaggcgtg 15840
 agccaccgta cccggcctgg tgactcactc ttcaaggga ctcagctata aagaagagag 15900
 gcatggagtg gtggtaggat ggattaagct ggggagggga ggcatttta agatgagaaa 15960
 tgatgtttaa atgcagatgg aaatgggaca ggagagaaag ggaagttgat gggataaatg 16020
 aaaattatca tcatcatct cctcaccatc accagcagca accacaagga aagctcgttc 16080
 tgctctggga ctgcctgtt agacatttct tccttatact gagccacaat ctatctccct 16140
 gtaactttcg cttttggta gatttctgcc ttgtgtccct ctctactct tcaaatggct 16200
 gcacgtgct actatgtctg tctcttct caaactcaca gccccatcc tacgggcct 16260
 gtgtccag cctggtgacc cgacctcta ctgtccagc caagttcaca tgaccatct 16320
 ctatttgc aagcctct tttttttt tttttgggag gagtcttact 16380
 ctgtccagg ctggagggca tggcacaatc tcagtcact gcaacctctg cctctggtt 16440

tcaagtgatt ctccctgcctc agcctcctga atgctgggat tacaggcata catgccatg 16500
 cctggtaat tttgtattt itagtagaga cggggtttca ccatgttggc caggctggc 16560
 tgaactcct gacctccagt gatccccctg cctcagcttc ccaaagtgt gggattacag 16620
 cgtgagccac tgagccgggc caagcccctc tgaagaggca gtgtgcacac tctgtcttg 16680
 ctgatagaag catgaaatgg ttaagttt tgtgatttg gaaaatgaac tcagctataa 16740
 aatgtgcaca cccagaagt gtcctatgga cttaatgtct taggggtgag aagaagcttg 16800
 tgcagggatt cctggcaaca tgaatgcggg atggacagac agtgagcaca gtgtgagga 16860
 gcacagcctg agtgcctc cctgagatca ggaacaggac aaggatgccc gcttcaccg 16920
 ctgctgttcg acattgtact ggaagtcta gccagagcaa ttaggcaaga aaaagcaata 16980
 aaaggcatcc agattgtaa ggatgaagta aaactatct tagtcacagg tggcacaatc 17040
 ccatataggg agaatccaa agaattcaca agaaaactgt tagagcaaat aaactaattc 17100
 agtcaagtig caaggaacaa gatcaaccac aaaaattaat tgtgtgggtt tttttacac 17160
 cagcaatgaa cattgaaaa ggaattcga aaaagtatt caticactgt agcatttaag 17220
 aaagctgagg ctgtgaagct acatggctgt gtttgagct ggtgtgcta ttaccatgt 17280
 gacctggac aagcttctca ctccatcat gacaccgtt atcagctat taagtggggg 17340
 taataatagt accttctta tagagtggg tttgtttt tgtttgtt tttgtgggg 17400
 gagagtaagg attgcaagag ttaatatag ttgcaaaaat ggacttagaa gagtgggttg 17460
 cacatggtaa ggactatatt agggttacca aaaaatgtt ataattggga aaatgaaca 17520
 aaggttccag taaatatcca tccagaggc tggttgaca attatcaca tgtaccttt 17580
 agctgaacc atatggaact gcctatatt tatagctatg gcctacaaa tgcaaatgg 17640
 gcagttttt ttgtgtgtg ttgtgttg ttgtgtgt tgtctgaga cagtctcagt 17700
 ctgtcaccag gctgaagtc agtggcagca tctggctca ctgcaacctc tgcctccag 17760
 gttaagcga ttctctgcc tcagcctccc aagtagctgg gattacaggt gtgcaccacc 17820
 acgcccagct aatttttga ttttagtag agatgggggt tccaccatgt tggccaggcg 17880
 ggtctgaac tcttgccctc aattaatct cctgcctcgg cctccagag ttctgggatt 17940
 acatgagtga gccaccgagc ccagccaaa tgggccgtt catatgttca attaaacac 18000
 taagatactc tttagccact gaaaatgact gtgtatcata gatctatct taccacagat 18060
 ctatcttac tgtcatagaa agaactaata acatattatt gaacaaaaat aacaagttac 18120
 aaaacagctt gcagccacgt ttgtaacatc agtatatcta ttgtgtgc taggtgtgga 18180
 cataaagaga tatctagaag acatcccat tgcctatggg aatgacttg gctgggatct 18240
 ttttatita tttattatt ttttatita tttattatt tttttgaga cagagtctca 18300
 ctctgcacc caggctggag tgcaatggca tgatcttggc tcaactgcaac ctccgctcc 18360
 caggttcaaa cgattctccc acctcagcct cccaagtagc tggaattata ggcacccatc 18420
 accgcaccta gctaatttt gccttttag tagaaatggg gtttgcctat gttgccagg 18480
 ctggtctcaa actctgac tcaaatatc cactgcctc agcctccaa agtgcctgaa 18540

ttacaggcgt gagtcaccgc acccagctgt gatcttaatt tatattttt cttttaacc 18600
 ctgtatttt accctgtttg actgacttta ggagtcigac ccccaaaact gcaaaagaat 18660
 aaattagtgt tgttgaagc cactaagttt gtggaattt gtagagcaa caaccagaaa 18720
 ctaatacaat cgggattaaa ctggccctgg ggggtgctggc tctccccac ttcttctca 18780
 gagcatctct tgggtcctcg ccccttctg agccccacc atggccgggt ggaccagcga 18840
 tggacctgat cggattctcc tcttgggatt ttaaataaga gatgcagagg ctgcacgcca 18900
 agcttctctg tgccgcactg caggcacagt cctggcgcta ctgttctta ggggtccatg 18960
 aagatgttg ttggcctggc ttagggtga gctaagagat acaagggata cactttggag 19020
 gctcagccct ttgatgtac cttgcgctg gaagcccagg ggctgtaaga cggagaggaa 19080
 ctttctgaa agggagggtcc tccctgcaga gctgaagag caatctaacc atgctggcat 19140
 ggggcccac ccacctgta tgccagtga agtctgtgac agtcactcat attcagagca 19200
 tgtggattga gactctgaag aacgacttta tgggtggcac ctcatctcc atctgggtgg 19260
 tccagagcaa gcagcctcag ggtcccatgg gcccaacctc accctccatc cttatttccc 19320
 tgccactcca gggctctcc tcagccagac ttgcctgtga gtgcccctg atggctgtat 19380
 tagtctgtc tcacgtgct aataagaca taccgaagac caggtaattt ataaagaaa 19440
 gaggtttaat gggcccacag ttcccatgg ctgggaaggc ctcaaatca tggcagaggg 19500
 cgaaaggcat gtcttacatg gtggcaggca aaaagtgaat gacaaccaag caaaagaaga 19560
 aacctcttat aaaaccatca gatcccatga gacttactca ctaccagag aacagtatgg 19620
 gggaaactgc ccccatgatt caattatct ccaccaggtc cctcccaca cacatgaaa 19680
 ttatgggagc tataattcaa gatgaaattt ggatggggac acagccaaac catatcagt 19740
 gccctggctt ctctccatt agctctgtc tacttccaa aatcttcca gctttgtct 19800
 ttttagaga ctggagctc actgtgttc ccaggctgt ctgaactcc tgggtcaag 19860
 ggatcctct acctcggtct aacaagtag tgccaatata ggtgcacacc accacacccc 19920
 cacaacttac ccactctcc cggtcagtg aaatccctag gtccttaca ctcccgttag 19980
 ctccaggtag cagcggctt gccttctca gaatcataa cattatagcc tcttagagca 20040
 aaaaggatc ggggatggg aaggctggga gtctctctt cattcaggga ttaacctac 20100
 cacctggagg ggtgacatg ctacacttag tcaccaagca aggtcgggag agtatttctg 20160
 ggatggatc tagcacttc ttccaaagg gaggatacta caaactcact ttcatgatca 20220
 cagtatccc cccgacaggt aaagtgtgca aactcgctt cattccctt ttgtcttga 20280
 cgagctctg aagcatttta attcttaaa ttacatgtt catgatgaca cagagacag 20340
 ttctgttt aggagagggc ctgattccat taaggctgat cgatcatatg tgcacattc 20400
 tgggataaat atgagagatc agaattgagc ttccaaatg gtgggtcaca tgacccagt 20460
 ttgtccagg agaacgtgc aacatccagc atgagaaagc gctgcatgca ccatgtatgt 20520
 gttgggatgc tgggtgaaa tgtattaatg agcaagtgtt atcatattg tctatggaat 20580
 tctattaat agctatgtct cgtatgtct gattgggtct caaatcggg tatcttacc 20640

caacatcttt ctgaaaatc ctctcttta aataaatgtt ggccgtgtgc actggcttat 20700
 gcctgtaac ccaacacitt gggaggccaa ggcgggtgga tcacctgagg tcaggagttc 20760
 aagaccagcc tgaccaacat ggagaaaccc catctctact aaaaatacaa aaattagctg 20820
 ggcacgatgg tggatgcctg taatcccagc tacttgggag gctgaggcag gagaactact 20880
 tgaacctggg aggcggaggt tgcaagtaagc tgagatcgcg ccattgcact ccagccttgg 20940
 caacagagtg agactccgic tcaataagt aataaataaa tgtatagaa atactaaagt 21000
 gtattgtgca taatataaca cccacgtacc accatccaac ttaagaaata aagcattact 21060
 aatccagttt tgctccctgc aaactcatcc ttcacccat tctcttctt tttgctttat 21120
 aaacatcctg cagccaggca cagtggctgc acctgcagcc tgagctactt gggaggctga 21180
 gatgggagga tcacttgagc ccaggagttt gtggctgtcg tgtgctgatt gcacctgtga 21240
 aatagccact gtgtccagc atgggcaaca tagcaagact ccatctctaa atgaaatcct 21300
 tttttttt ttcttttg agacagagtc tcactctatc acccaggctg gagtgcagtg 21360
 gcgccatctc ggctactgc aacctccgic tccagggttc ctgcctcagc ctccggagta 21420
 gctgggacta caggcacatg ccaccacaac tggctaattt ttgtatttt agtagagacg 21480
 gggtttggcc aggttgggtt cgaaccagg acctcagatg atccacctgc ctggcctcc 21540
 caagattac aggcattgag caccatgccc agcctaaaag aaatcctgaa ttgacattt 21600
 ttaattccct tgtttgttt aatatttcta cagcacatat atgtctacat atataaaaa 21660
 tataatttt tattgggggtg aaattcacat aacgtaaaat taaccactgt aaacatgaac 21720
 aattcagtg catttaggac attcatagta ctgtgcaacc accacctcta tctagctcca 21780
 gaacatttc ataccccaa aaggacactc catgcccccc aggagtcact cccatttct 21840
 tcccactctg tcccctgcaa cgaccaatct actctctgic tctgtggatt cacctattct 21900
 ggatattica tataaatgga atcatacat atgtggcctt ttgtactgg tctctttgac 21960
 ttactgggt gtgggtggcag gagctacca cctacttggg aggtgaggc aggagaatcg 22020
 ctgaactcc agccataaat tatattgtt tcaaggttca tcaatattac tcgccaggc 22080
 tggagtgcag taatgtgacc ttggccccct gcaacctcca cctccccgt tcaaaggatt 22140
 ctctgcctc accctccga gtagttgga ttacaggcgc ccccccccc caccggcta 22200
 atttttga ttttagtag agagagggtt tcaccatgtt ggtcatgctg gtcctgaact 22260
 cctaattca agtgatccac ccacttggc ctcccaaat gctgagatta caggtgtgag 22320
 acactatgc aggcctagtc aattcattt cactgctgca aagcatctca ttttaaaaa 22380
 ttattttt attatttt ttgacaggg tctactctg ttgccaggc tggagtacag 22440
 gggtgcaatc acagcttact atagcctga cctccgggc ccaagcaatc ctccactgt 22500
 ggccctcaga gtagtgga ctacagcct gtgccaccac acacagctaa tttgttct 22560
 ttgttttg tttgtttg tttgtttt tgagtggag ctgtgctt gttgccagg 22620
 ctggagtga gtggcgcat ctacgtcac cacaacctc atctccagg ttcaagcat 22680
 tctctgct cagcctccg aatagggtgg attacaggct cagccaccg taccagta 22740

atttttatt ttagtagaga tgaggtttca ccatattggc caggatggcc tcaaactcct 22800
 gacctcaggt gatccatccg ccttggcatc ccaaagtctt gggattacag gcgtagacca 22860
 ctgtgcctgg ctcaaagcat ctattgtat gaatccaaca attttttat ccattcgttg 22920
 tttagtaca ttaggttat ttcctgcttt cctctttaca agcaatgctg cagaaacatt 22980
 ctgtccata gctcctccgg cacatgtaca agagcttctc tagggcaca atgtaagagt 23040
 ttattgctg ggccatgggg aatgctcttc tcaaatgta ctgctattg tcaaatctga 23100
 gctatcttt caggagccta ctactgggtc tcagccctgg ctatgcatta gaatcactg 23160
 ggaactttaa aaaaatatgt gggatctgg ataactcccc cgacccgtgg aatcctagtc 23220
 tccagggatg ggactgggga ctgaaagct ccctgggtga ttacacagcc aaggtggaga 23280
 acagctggga ctgagcttg gccactctg tctcagcttc ccaaggtag ctcatagagg 23340
 gcagccatca tcttagactt gctaataca ctacagcagct gagcagatag aagggactaa 23400
 ataaataaag tctgaatgag taaataaagc atttaaaaca aaccctttag ggctgggtgt 23460
 ggtggctcac acctgtaac ccagcacttt gggaggccaa ggtgggtgga tcccttgagg 23520
 ttgggcattc aagaccagcc tgaccaacat ggcgaaacct tgtcttact aaaaataca 23580
 aaattagcaa agtggtggg tacggcctg taatcccagc tactcggaag gctgaggcag 23640
 gagaatcact tgaacccggc aggcagaagt tgcagtgagc tgagatggag tcaactgact 23700
 ccagccggg ctacagagca agactccatc tcaaaaaaa cataataaa caaaccttt 23760
 aggtgggtg cgggtggcga tgcggtaat cctggtaact atctctact gttgtacga 23820
 ttaaggaga tgggtcatgc agagtactg gcacatatga agtgctcaat aaagagtaac 23880
 tctgcagcc tgggtgggtg gccacacct gtaaccacaa catgttgga ggctcaacat 23940
 gttggcagga ggattgctg aggccagggt ttgagacca gcctgggcaa cataagaga 24000
 ccccgctc tataaaaaat aataaaaact agctgagcat ggtgatacat gcctgtagt 24060
 ccagctactc aggaggctga ggtgggagga tcaactgag ctggggagt ggaggctga 24120
 gtgagctacg atcacgccac tgcctccag cctgggcaac agagcaaggc ccttctctg 24180
 gaataaaaa aaaagagagg ctgggcgag tggctcacac ctgtaatccc agctacttg 24240
 gaggtgagg caggagaatc actgaacct gggagggtga agttgcagt agccaagatc 24300
 gcaccattgc actccagtct gggcaacaag agcaaaactc tgttcaaaa aaaaaaaag 24360
 agagagagag agagtactg ccattttcaa tgtattgat aaagcagggt ggtgtttctg 24420
 aaaatgcacc agatgcagcc ttagactgga aggtgctgat gtgtactga gcctccagac 24480
 aaagctggcg aaccaccgg ggcgagact tatgactga gaagctgtcc tatctatgag 24540
 ggtccatgtt agggaatatt aaagaatagg attgggggca gtactttcc ttagcaagt 24600
 gctagctatg tgcagatc tgggcaagg agtctgcaga cgtctctcat ttcactca 24660
 caatccatga ggcaggatg actattcttc ccattatcca gagagggaaa acaagtcaca 24720
 cggaagcaag tcacgcagca gagctaggat tgaacagaa gtgcctgcat cctcaacgc 24780
 ctgcaccca acccgctgtg ctatgcctcc cgttgatgct ttccatgtc tgccattag 24840

cttgtcttg gtgagtaca agctgtacca gcggaaagag cctgtcatca gttctgtgca 24900
 caccaagggtg aaggggatag cagagggtgaa agaggagatc gtggagaatg gagtgaagaa 24960
 gttgggtgcac agtgtcttg acaccgcaga ctacacctic ctttgcagg tgagcacctc 25020
 gtagcattct cccaggctcg tcgtgggtca ccgtcgccag ggcctagctc cttcccta 25080
 ggatctacag cctcactcca gaaaaaacgc tggctctatt ttaaagctc tgaacatc 25140
 ccaactgaga aaacctaaaa atgcataaac tgggtgagat ccaggagatg attcttctgct 25200
 ttatcaagtc ctacagggtt tctcaaaata gcctcatgct ctgggtgcat gaaaacagga 25260
 ttgattcac tcgcgttga ttaccgcga gctttttaga cgcacagcca ccaagccagc 25320
 aaggctctgcc gagattcaga aggacagaga actttctacc aaagagcacc agcccttcta 25380
 gattgtgtag catttcatca acaaaaatga ctccaggaga ccattcttgg gtacatttt 25440
 ccacaaagaa aggatgtaaa tgacaaaaag aagaaaggca tcggtcactg agagaaggcg 25500
 tgtgacttct agatttcaa gcagggaaaa tgaagcaga attgccaaaa aagagaaatt 25560
 aaagtagggt tgaggagtcc aaaaaacaga taagggggcg agagggacaa aggggcacgt 25620
 ttgcggcttg gggggctaaa atgcttga aagacccatg aagcccaagc tgctgttgt 25680
 ttctaacagt gggtcattaa aatcctgtga ttgctcaggt ggtgggatgc ctagtgccc 25740
 ttcatataa gtctgggaaa atctgaacag tttgtgatg aaggctgctc ccctacctic 25800
 gcctcccag gtctctgag ttcaattta atgagattt tactgcgtaa aaaaaaaaaa 25860
 aaacaaaaac aaaaacatga aaaggctgct ttgagactg catttgtaaa tgactttca 25920
 acccattcaa acgctcttc actctcctc agctcagcag ggctgctcgt ccagcttga 25980
 tattaagccc ttggcatatt ccaagttgcc cacagatcct gatttctaga agcttagaaa 26040
 agtggagagg ttgcccagc aagctggatt attataatta agtagttctc tttcaaagg 26100
 ccttgcatit tctagctc tccttctcca cagggaact ctttctctg gatgacaaac 26160
 ttctcaaaa cagaaggcca agagcagcgg ttgtgtccc aggtaggag gggacctgga 26220
 gtgtgggtc aggtcttaag agttcttgg ggagggtgcaa gtcggaagaa gcagaaatgc 26280
 ggacctggg gtgtattga gccacaaat atctactgag cacctgcctc tttgtggg 26340
 tagggctggg ctgagtggag gaaggagag aacacatggc agtgtctc catggccacc 26400
 agatcttct ctgctgctc cggtctctg ttctagcact agggacctgt gcagacggca 26460
 accctgctt ttctatctt ggctctctc gcagcttatg acaatgggtga ttctggcat 26520
 ttgcacaaca ttacattt tacttgaggc tcacaatgac ttaggagaa aaatggcatt 26580
 ttacaaacca gaacacagat tcagagatgt tgtgactcc cccagggtca cacagcttgc 26640
 tgcaggcaga gccagggtta ggacctgagt gtcctcactc acagcttctc ctaccatggt 26700
 tcacagtga ctgtctgag ggctggtt ggaaaaccct ggcacacagc tctctgctc 26760
 attatccac ttcttagat ctacagctc ccatcacgt catgtgcccc catgactccc 26820
 tgccaccacc ccagacaatc ttccaact tggcttctct ctctcaaaag agccgctct 26880
 ttctggaaga gttctctcc tatcaatca tattgtttt attctgggt aggcacgggt 26940

gctcacacct ataatctcag cactttggga ggccaaggca ggaggatcac tggagcccag 27000
 gaggtaaga ccagcctggg caacatagtg agacctgtc tctaaaaa aaacaaaaca 27060
 aaaactgtat ttgtcctatt cttacctca ccatcctgtc ctctcctct aaaccagctt 27120
 ttcttctaag ttatctata ccagcgtgtt actgctaggc attacctgta actcccaacc 27180
 tccagccctc ccctctcagg taacgtaact gtgctgtacc agggaccatt tccaacactg 27240
 accgttgtag cagcgtggt gatacccagc cagatcctcc actgatgagc aggaccagcc 27300
 ctgccggtgg agttttgtc acggggccac tctggcaaca aggtctagct tctgagtgt 27360
 ccaggacctc tgggcaaaag tgctgtctt gactacagtc ttgccaatca ttgtacca 27420
 aattcagaac caagataaca ttcccttggg gtctatcttg ataactcctc gtctaaaagg 27480
 agagctttct acattgttt tgctctcctc acccctaac tctcctgtc gctcagatga 27540
 aatctctgac agccaggcca gccaagccct ccctctgtc tccctccca gcagtgtcc 27600
 gtctccaatc cagttgggg gcctgttcc cttttgtcc atccttggg catctctaca 27660
 ccacactcta cctcatgta cacataaaga cacttttga cttattgaga ctggaagaa 27720
 aaagagttag gatagcccag agattaagag cctgggctca gaagctagac cgctaaaggt 27780
 tcaaaactta aactcttaa ctctgtgacc tcaggcaggt gactcttcac ctctctgtc 27840
 caagtgtgt gcatctgtaa aatggggctg agaacaaaac ctactgttc ggattattga 27900
 aacaaatgag tgaatacctg tgacacgctt aaaacaaggc atgttacatt ataagcactc 27960
 aatgaatgac agttgttatg attcaaatga catgttact ctcatagagg atgtcactca 28020
 gagtactgg aacccaacag gtccataga agcacttcca gccagagag cacatgtgac 28080
 tcttgggat tttaactat gctccaggct ggttgggtt ggtctaattg ggcatttgc 28140
 tctctctgc ccatcaacac ccctctatct cgggtgtct cggtaaatat gaccaagttc 28200
 cagttacaca ccaacatgaa tagatattg ctccatttat ctgtggctgc aaaacaaact 28260
 accctgatca tgggtgggtt aacacaagca tttattaac tctcacagt cagtgggtga 28320
 ctgagttcag ctggttggtt ctctgtctg tctactagg gatctctca ggagctgcag 28380
 tcagatggca gctgggggtg gcatccctt gagcctcaat tggcaccag ggatagctag 28440
 acatctctc ctctctggga acccctaaga gttaccctct ctcaaggtc tgtccgcat 28500
 gtctctccag cagggtagtc aaacttcta atagtggctc aggttctca aaacaaaaca 28560
 aaaaaataga agctccagg ccttctaag gcttaggcct ggaactggca acagcattcc 28620
 ttctgtctg tctattggct tgcagtcac agcacaggct cagcttcaga aagtagggga 28680
 aacactccac ctctcaatgg agaaagtac caagaattg tggccatctt taatccatgc 28740
 aacaatcaca aaagagaaga ttaaaacaaa actaatgtc tgtttttaa aaaaaaatt 28800
 agaattcctt ttggagagca aaataaaact gatttctct ctctctgag attcatctcc 28860
 agtgataat gaaaatgtaa ctttccatga gccacaatga atgatcacac attcattcct 28920
 tccatgacta tcaagaccta ctattgcca ggcactggg ctgcaagat atgaccttgg 28980
 ctctgtattc aaagggttca taatctaac tggggagtca caaggggagg agaccaacgg 29040

catgccatgt gacaaggcag cctgggaaat aaagagatca tgagagagtt tgaaaactac 29100
 aagaacatt cattgagcat gtactatgtg ctaaaccctt tctttcata actcactta 29160
 tcttcacaac aaccaggcaa gacaattgtg tggctgggca tgatggctca cgcctataat 29220
 cccagcactt tgggaggtcg aggcaggcag atcatgaggt caggagtca agaccagcct 29280
 ggccaagatg gggaacacc gtcttacta aaaatacaaa aattagccag gcatgggtgt 29340
 gcgtgcctgt aatcccagct actcaggagg ctgaggcagg agaattgcct gaactcagga 29400
 ggtggaggtt acagttagcc aagatcacac cattgcactc tagcctggtc gacagagcaa 29460
 gactctgtct ccaaaaaaaaa aaaaaagaca aatctgttca ccctgctttt 29520
 acagatgaac agtgagctgg ccaaggcttc tcagctagaa aatgggtcag ctactgtga 29580
 accagatata cctgggtcca agacctaggt ccttgacctg aatcactgta gaagccacaa 29640
 attgcacaga tgacaaagag cagaggccat tticacacct atgtcatgtt ttggtctccc 29700
 caagactggc catgaagcct gtcctctaga aaacaggaag agaaatatgc ataaaccagg 29760
 gcagctcatt tctctctcc gtgtctgtc cttcagccct gagaccagc tggctcatt 29820
 tcttgggcc aggcgaggtg cctggccaat gtctctcagt tccatgtgtc ctgctgagtc 29880
 tccgcacctg ttgccctgc tcagtcctt cagccacagg gccacttga ccaagccacc 29940
 gtcccccat cccagccagc atccctgaaa gtgatcccta gcttccaga cctctgctt 30000
 tcccatttt atttattgt attgggatat aattcacata ccataaaccc caccatttta 30060
 aagtgtacat acggttcagt ggttttagt ataatcacag agttgtgcaa ccatcaccac 30120
 cgtctaattc cagaatatt tcttctct ttttcttt ttttcttt tcttcttt 30180
 ttttcttt ttttcttt ccttcttt ttttcttt taggtggagt ttgtctgt 30240
 tcaccaggc tggagtgcag tgacgaatc tcagctcact gcaacctca cctcccaggt 30300
 tcaagtgaat ctctgcctc agcccccta gtagctggga ttacaggcgc acaccaccac 30360
 atctgataat tttgtatt ttagtagaga cgggggttca ccatgttgag caggctggtc 30420
 tcagactcct gacctcaggt gatctgccc ccttggcctc ccaaagtgt gggattacag 30480
 gcgtgataag ccacggcgcc cagccccag aacattttca tcacctaaa aggaacccc 30540
 aaatccagta gcagtcact cccattctc ccttcccctg tccctggcca cagtctact 30600
 tctgtctcta tagatgccta ttctggacat ttctataaa tagaattgta tatggtgtg 30660
 cctttgtgt ctgtcttt tcaactcagca tcatgttctc caggccatc catgtttag 30720
 cctgtgtcat tgcctatcc ttcttatggc taaataagat tctgtgtatg aatgtaccac 30780
 attttattg tccattatc cgtcagtggc cacttgcag gtttccatt tttggcgat 30840
 tctgagtagt gctgctataa gcattcgtgt gcacattctg gtggatatcg aatcattct 30900
 ccacatctta gtaacacag tcacttact cccactctgt catcttcta tctgcagtat 30960
 cccaccgca ggacgtctg ttctctgac cgagggtgta aaaagggatg gatggaccg 31020
 cagagcaaag gtacctctg tttttttc cgagacccta ggggtggatg gtctggcatc 31080
 ttggtgacat ttgtatgcc caggctaggt cttcagctc tgctctcagc tgcctcttc 31140

caccatcacc aagccatagg cgagtctgcc catgcttcgg ctctgtcccc agcagaccag 31200
 ctgctgactg taaacatgac tccagtttc cagtgagaga agaagctcct aaaaacctag 31260
 caggttcagg attctaactg gtgaaaaatt cacatggcct atagcatcat ctgagtattc 31320
 taaactttcc ccctgaattt cctcaaaggt tgaggacat gaactttac ccccaggga 31380
 cctggcagca ataccatata taacctgcag aattttttt gttttttt ttattttt 31440
 ttttaacat ttttgcaact gttttttt gatttgatt ttgattttat ttatatctaa 31500
 gtgcagtgtc attgcgatac ctgcagaatt tctttatctc acattttaac taaaaaggc 31560
 acagggcagc gagcgagag gctgggtgcct gtaatccag cactttggga gggtaggca 31620
 gatggatgct tgaggtcagg ggttcgagaa cagcctggaa aacatggtga aaccccgctc 31680
 ctactaaaaa taaaaaatc agccagacat ggtggcacac gcttataatc ccagctactt 31740
 gggaggctga gacgtgagaa tcactgaac ctggaaggca gaggttcag tgagccaaga 31800
 tcatgccact gcactccagc atgggtgaca gagcgagacc ctttaaaaa aaaaaaaaaa 31860
 aggcacaggg caattttaaa aatactgcaa atagtaaaaa aaaaaaatc agtggttata 31920
 atgcaaacac acacaaaaag gcatatgcc attactgcat tctactccat actgtatgtg 31980
 tatttgatt agtataaaag ttatttaac attgctcact atttaattaa ttctcccttg 32040
 gaaactgatt aatcatcctg gcactccagg aagatgtgcc atgctgattt catggcttg 32100
 cacatcctgg gcaggctgtg tacccttga gggactgtg cccctttgag aggccatgtt 32160
 ctagtccatt tatactaagt gagagcatac acctgttccg ctcccctcat gggcacctt 32220
 tcttataaag aaacaaaaga gccagcagaa tccacagtct ttctgtgtc tctctgatc 32280
 ttattgtt ttgctgtt gccttgcctt gtgttcgtt tggttaggat gggcttgatg 32340
 gaagctgaag ctgcgtgggt tggaaagcct ggtcaaagcc tagtctctc cccgggttga 32400
 gttaatgatg tccctcctgg agaagctct ctctgcagtt cttcacatc tgtgttcta 32460
 cgatgcttg acccctatag gaattcagac cggaagggtg tagtgcatg aagggaacca 32520
 gaagacctgt gaagtctctg cctgtgtccc catcgaggca gtggaagagg cccccgggtg 32580
 agtcgcatgg ggagacagac acagtggccc tcagcgcca ccagatgagg ccttgcgag 32640
 gctgcttggg ccttcccctc tcagcacagc cctgcaaagt cctgggtcct accggcttg 32700
 ggacccctgc gctctggatg cactgcttg cacaactag tatctctgg agggccatgg 32760
 tggttgtaa actgtgttaa caatcctga ccaactgga aatagctact accctgagca 32820
 tcttgggtg tccctggccc ctctctccc ccagatctc cagggtaccc ccagacccc 32880
 tctgtagtg ccacagcagg atccctctg actgtcagt gtccatactg agtgatcaag 32940
 gatagaaagg aaggaggag atggaaggga aggacgaagc gaggaagag aagggaagg 33000
 ggaggaaaa gcaaaagggg tgagggtaaa agggggggga aaggaagtt tctcagatta 33060
 aatgcttaca atgacataca gatttggtg tccctgtat tgatgctcg cttaataca 33120
 caaagtcaca atgttaaatc tcagaagca caagggtga tgatttcag cagagaatag 33180
 ttagaaagac ctggattcaa ttctaactc taacaccatt ttgctgtgtg tcttggga 33240

aatggcttaa cctctctgag ttacagtgtc ctacactgta aaagcagaat aataattca 33300
 ccaacttcat agggctgttg taaggattaa atgagatgat actgtacag ttattgtaag 33360
 gtaagcccca tgcctgctg gcttacacac acacacacac acacacacac acacgcacac 33420
 acgcacacac acacacacac acacaatcta cccctagaag tgtgggtggt ctagaccagc 33480
 actgtccaat tgaactgat gcagtgtgg aaatttctgt atctgtgctg tccaataggg 33540
 cagctactag gtacatgttg ctattgagta catgaaatgc gactactgaa ttttgaaag 33600
 agatgataga ttagatagat aaagatagat agatagataa atagataata gatagataga 33660
 caggtagata gatagataga tagatagata gatagataga tagatagata gatagagttt 33720
 tgctatgttg cccaggctgg tttgaactc ctgggctcaa gcgactctcc tgccttgccc 33780
 tcccaaagtg ctgggggtac aggtttgagc cattgctccc agcctgaatt ttaattaaa 33840
 ttaaattha aatagccaca catgtctagt ggctaccata ttggacagcg cagttctaga 33900
 ccgatgtgat tcaggatcat tccctcagca tcgtggggca aagagaaaac tgccccaagc 33960
 tggcctgtag aaggctcagg cgaagggttc ccaatgccgg gatggggggg gcgctcagca 34020
 gcacacccc ttatgattct caatcgctaa tagctccact cagggtcatt tctcggtcag 34080
 gggcatttct ttgggaatca cccagctctg ggagatacag cagcctccac tcaggtagtc 34140
 cttgttcaag acaagcggcc ctgactgac tgcagtttca gtccagctc tgctatcaac 34200
 tcactcatta aataaactgc atctccagtg tgctgcctc tgggctggat ttgacgtga 34260
 cctgggcaag caactcctg aacttcagtt tctcatatat tatatgaatt agctaagatg 34320
 gticgtttaa tcattcattc aacacatcca tcaccagta gtaggtgta gatatttatt 34380
 tcatacgtaa ctacgcataa gagactttgc taagttttag gtaaaataca agtcccagat 34440
 acggagcaag tctcaaccac tttacatacc tgaatgtgta attacatcac tgaagaggt 34500
 gccacagtaa atgccactgg gtctgtgtt agtccattct cacacaaaga actacctagc 34560
 cagggtcggg ggctcagcc tgaatccca acattttggc aggtgagggc aggcggatca 34620
 ctgagggtca ggagttgat accagcctgg caaacatggt gaaaccccat ctctattaaa 34680
 aatgcataa attagccagg tgtgggtgca cagcctgta atcccggcta ctggggaggc 34740
 tgaggcagga ggtcgttg aacccgggag gtggagggtg cagtacccg agatcgcgcc 34800
 actgcactcc agcctgggtg acagagttag actccatctc agaaaaaat aaaaataaaa 34860
 ataaagaact acctgagacc aaatacttta cgaaaaaaa gaggtttaat tgactcacag 34920
 ctccacaggc ttaacaggaa gcctcaggag acttacaatc atggcagaag gcgaagggga 34980
 agcaaacaca tcttaccatg atggagcagg aggcgaggtt cgggggatgt gccgcacact 35040
 ttaaatgat cagatctcgt gagaactcac tcactatcac gagaacagca aggaggaagt 35100
 ccgccccat gattcagta cctcccacca ggccctct ctggcacatg gggattacaa 35160
 ttaagatga gatttgggtg gggacacaga gccaaacct atcagatctc aagaaggag 35220
 aaattctct tggaggagct ggaggggctt tgtggagagt ttcagaatgc ttgccact 35280
 aggttgctg taicatttc tctcatgta tccaaagac caagccaaga aaccagaagc 35340

ctctgtgtccc actggcccat gggctccctc ggtccccacc gtcactaatg gccattttgc 35400
 atgtctctct cccaggcctg ctctcttgaa cagtgccgaa aacttactg tgctcatcaa 35460
 gaacaatac gacttccccg gccacaacta caccacgtaa gtgcccaggc tgcctggctg 35520
 tcttagttat ctactgtga gtaataaatt atcccaaacc tcagaagcct gaaacaacaa 35580
 acgcctattg tctcccacgg ttctgtggg tcaggaatct gggaatgact ttgtgcgtg 35640
 gtctggctc aaggtctgtc aggtgttagc caagctgtca accagggtg cagtcattc 35700
 taggcttgac tggggctgga gaacacttt ccaagctctc acacagtgc tcgtgggaga 35760
 gctcagttcc tcaccagtg aacctgccc tagaccactt gagtatcctt ggtatatgtt 35820
 ggctggcttc tcccagagca agtgaccaa gagagacaga gcaagcaacc aagagtataa 35880
 ccaagatgga agccacagtc ttgggggga gacccaaca ctctgcat atgccattgg 35940
 tcacacagat caacctggt ccagtgtgag aggcactgc ccagggttc caggaggcag 36000
 tgatcattg gggctttat ggaacctc caccactg gtcactcct gggaagaga 36060
 cagatctgt tcaatcag atgtttgtt gttgcttt aattatgcac aggagaaaca 36120
 tctgccagg tttaacatc actgtacct tcacaagac tcagaatcca cagtgtcca 36180
 tttccgact aggagacatc ttccgagaaa caggcgataa ttttcagat gtggcaattc 36240
 aggttggtg tgctttgac actgggatgt ggggctgtgt gtctaggat ggaggatgc 36300
 aaacagccaa gaggccgggc cactgggtct tcataatgt gctcacatt actgagcatt 36360
 tagtaatcc acccgctcg ctaaggtct tacctacgt acctcgtcaa atccaaaac 36420
 aatcctatg agtgagagct acttggtga ttccttct gtggctgct tagcaagta 36480
 tcaaagctta gtggctcaa acaacacata ttgcttatg ttgccagaga tcagaagtg 36540
 gagatgatt tccctgagc agggcggtgc tccctccggg actttaagg agaatccagt 36600
 tctcagctt tccacctc tggagctga ttccttgc ttttcaaag ccagcagcat 36660
 aacatctgc ctactggcc acttcactc cctactctgt gtccaatc ctttgcctc 36720
 tgtcttaca agagagagag cattacaag agggggcatt taaggacaa ctggataatc 36780
 caggataatc tccatctca agatccttca tttaggtg gcacgggtgc tcatgcctgt 36840
 aatcccagca cttgggagg ctgaggtggg tggatcacct gaggtcagga gttcaacacc 36900
 agcctggcca acatggtgaa agcccatct tactaaaaat acaaaaaaa aaaaaaatag 36960
 ccgggcatga ttgcaggctc ctgtaatccc agctactcg gaggtgaga caggagaatc 37020
 gcttgaacct gggaggcaga ggttcagtg agccgagatc gcaccactgc actccagcct 37080
 aggtgacaag agcgaaact catctaaaa aaaaaaaaa aatccttcat gtattcgcat 37140
 ctgcaaagag ctttccctag gggagtacta ggaggtaaag cagaaaagat attgataga 37200
 gtgccctgaa ttccagtca ataagttgg acttgatctt taatgggggc gtggggggca 37260
 ttaaaggtgt ttgggtacag gattggtctg ttgaaagtg tattttagga caatgagtt 37320
 aacagtgatg tgcccagac gggggtagg agagtgagga gatgcgatt tggctgccac 37380
 aataacact gtgcgagta ggtggggctg tacatatgt tcttcaatca gcattttcc 37440

tctaaaaacc ttaagcaatc ctggctatgc agggagatgt ctggcgggtg cgtaactcac 37500
 acccagcagc catagagact gtcccttgtt gatccttcag ggcggaataa tgggcatiga 37560
 gatctactgg gactgcaacc tagaccgttg gtccatcac tgccgtccca aatacagttt 37620
 ccgtgcctt gacgacaaga ccaccaacgt gtccttgatc cctggctaca acttcaggta 37680
 actccaaggc ccaggtaaaa ctacccaggt ggctgaatcg cattcccagg aactggtagg 37740
 actaatttg gttccaagg caacaagatg aatgaaaaa gactttctct aagaactagg 37800
 tgataactga atttttcca taattttta aaatttcaa aagagatgca cactctttat 37860
 tttttactt attttttt ttgtaaatg gagtctcact ctgtcaccca ggctgaagtg 37920
 cagtggcgcc atctcagta ctgcaaactt ccgcctccca ggticaagcg actctcctgc 37980
 ctacgcctcc caagtagctg agattacagg cggaatgaca ctgtttataa aacaaaacta 38040
 ttgggaaaca gaaaagcata gagggggatc aaaatcaccc ataattcccc taccctgaaa 38100
 taatcaataa caaccctcgg gggaatttc ctcatctgta ccaattttt catacagctc 38160
 ctatgagata atagcatata tatatatatc ttgtggtatt ctgcaggggt ttcatacca 38220
 cagccactca aaattcttg taaccatcac attaatgac ataacattcc atttttagg 38280
 tgaacaaata acaactgcta caattcagga agtgttttct ttcttttct ttcttttct 38340
 tttttttt tagatggagt cacactctgc ttgccaggc tggagtgcag tggcatgac 38400
 tcagtcact gcaacctctg cctcctaggt ccaagcgatc ctcccaccc ccaagtttct 38460
 gggaccacag gcatgtgcca ccacaccag ctaattttg tatattcagt agagatgggg 38520
 ttctactgtg ttggccagtc tggctcgaa ctctgacct caagtgatct tcccacctg 38580
 gtttccaaa gtgtaggat tacaggcatg agccactgtg cctggcccaa ggagggttt 38640
 ccatatacca agcactcccc atgccatcc ctaaatctcc caacaacct ggaaggaaga 38700
 tattgtttct ggaagatgat ttgccaaga cccacagctg atagtacatg ttgcataatt 38760
 ctaaccacg tcactctgac cccacactca cactccatcc ctctctccc atctcatgat 38820
 ttctcacct acgcciccat gattgaatat ttgagttgct tccagtttt ctattacaag 38880
 taaccacagt gtgcatttt gcacataaac ttctcttga attccaggtt acttccttag 38940
 gataaattc tagacttatt gaatacaagg ttgtgaacat ttatcatat gctttttatt 39000
 tttaaaaata tctatagta taatgttca tttttttt tgagacagag tctcactctg 39060
 tcaccaggt tggagtggac cgggtgcaat tatagctcac tgcaacctct gcctcccagg 39120
 cccaagtgt cctccacct cagcctcctg agtagctagg actacagggt cagccacca 39180
 tgcccagcta atttttaaaa ttttttga gagicgggt ctactatat tgcccaggct 39240
 ggctcacaac tctggctca agcaatccgc ctgccttggc ctctaaagc gtggggattg 39300
 caggtgtaag ccactgcacc tggcctataa ttctattc taggatttt attggtgct 39360
 tttaaaagt catctattat tgttcagtg agtccattc ttaccttaag gatcctac 39420
 ctctgtcca ttctactgta tcatctttt cataccgacc tattatcta agtaacttg 39480
 gtgggagtc tctctgtgg ctgtgaaata ctgtcttcag tagaaaagat ctatgcaa 39540

gtctgcatg tgctgtgtg tatggagtat tccttatata ttactcaaag cagctccata 39600
 aactggcagg cagcccccta aatgtgttc tgttctgtt ttttgttt ttggtctgca 39660
 tgatgttta aaacttgaga caggccaggt gcggtggctt atgcctgtaa tcccagcact 39720
 ttgggaggcc aaggtgggag gatcactga acccgggggg ttgaggctag cctgggcaac 39780
 atagggagac cccatctcta caaataataa ttttaaaac atcagccagg tgtgggggca 39840
 tgcacctctg gtcccageta ctccaggagg tgaggcagga ggattgctg aacctgggag 39900
 gtggaggctg cagtgaaccg tgggtgtgcc actgcactcc agcctgggtg acagagttag 39960
 acccttcat ggtggccact ggctggagct gattatcagt ggtcctattt agaaaggggc 40020
 tgggctttct ggttcacac tgcctccgc actcctagt gcttatact ggcccacac 40080
 actcattct gtcactgcc tggccctgt gtagacattt gatttgaac cccttgactc 40140
 aaaggcaggc tgatgcttt tgcctcctt ggatcaatgg aattcatcc aaggcaatgg 40200
 aaccagctcc taattgat aacttttga tcatcccctg cactgtgtt gaataggctt 40260
 atggccatta agaagagagg acaggatgaa tggctccctg actgactgcc ctccagggt 40320
 gtttttag ctgtcagcag aagcatcgg ggcatgtac caatcagggt tcagcaagt 40380
 tcttccagc gactgagtg agcacaaaa catctgtcc tggagagacc tgagccctt 40440
 gaaggcctca gccagtatg atgtaatgt tcttttagaa caaagtggta gacattgt 40500
 tctctggaat gagcaccaga agaatctct ggccaaaaat aaattgtga gggccgggag 40560
 cagtggctta tacctgtaat cccaacactt tgggaggccg aagcaggagg atcacctag 40620
 gtcaggagtt cgagaccagc ctggccaaca tggtgaaacc ctgtcttcc taaaaataca 40680
 aaaattagct ggacgtggg acaggtgcct gtaatacagc tactgggagg ctgaggcagg 40740
 gagaactgct tgaacccgga aggtggaggt tgcatgagc cggggtcatg ccacttact 40800
 ccagcctggg caacagagca aagctctgtc tttaaaaa acttaataa ataataaat 40860
 tgttgaggtc tgatgagtaa gtggacaagt tatttcag cagacacaca aaagagaagg 40920
 aaattacagg ttatcagg tatttcagaa aataaactt tctaaacat aggaagtga 40980
 agaagtgtat cacattacag aattctgtt tttagaaat gacctgtggc gaaatgtct 41040
 tattcagtga ataggtgatt ccgttatgc acgacctgt tgaagtggat caggccacc 41100
 agaatgcag acgcgttct caggccagc aggagtatgt gtctgtgta atttctgtg 41160
 gctattatga ctaattgcca caaatgtgtt ggctaaaac aacagaaatt aatttctta 41220
 tagttctgga agccagaagt ttggaatcaa gatgtcagca gggccacact cgctctgatg 41280
 ctctacggga gggctctctc ttgctcttc cagcgtctgg tggctccagg cattccataa 41340
 ctttatagca gcgtcccaca aatctctgcc tccatctca catggccttc tcactgtgt 41400
 ctctatgtct tcaatctct tttttttt tttttttt gaggcagggt ttactccag 41460
 tctcctagac tgaagtgaac tggcgtaatt tcgggtcact gcaacctctg cctccgggc 41520
 tcaagcatt tgatctctc ttatcttat aaagatacta gtcattggat ttggggctta 41580
 ccctaaatcc aggataatct catcttgaga tgttaactt aattatatct gcaaacactg 41640

tatttccaaa taagggtcata tcacagccac tagggattag atacttgaac atatcttatt 41700
 tgggggctca acccattcca gtgtacgaaa aacactcttg ttcaaggccc gatgtttctc 41760
 agggcatagc ccactgacta cctgcatcag aataatcact tggtagctgt actggaaata 41820
 cagactccta gaaacatctc agagcttctg caaccactct ttgagtgagg ggctcaggag 41880
 tctgcctctg aacacactca cccaagtga ttctttctt ttcttttt ttcttttt 41940
 ttgagatgg agtcttgctc tgtcccccag gctggagtgc agtggcgtga tctcggctca 42000
 ctgcaagctc cgctccccgg gttcacacga ttctcttgcc tcagcctccc gagtagctgg 42060
 gactacaggt gcccgcacc accccggct aatttttgc atttttagta cagacggggg 42120
 ttaccatgt tagccaggat ggtctcgatc tcctgacctc gtgatctgcc tgcctcggcc 42180
 tcccaaagtg ctgggattac aggcagtagc caccgcgccc agccatcacc ccaagtatt 42240
 ctgacctca gtttaagagc cactgtaaca agactatgga agcagaaatt cacgtgctta 42300
 ctacacaatg ttaacctcc caggcaaacc aactcacata gggagataat gccaatccca 42360
 gggcaggcag tggcaatgca tgcttgctg cgaattaaaa acaaatcac tgcctgggca 42420
 cgttggctca tgcctataat cccaacaatt tgggaggctg aggtgggtgc atcacttgag 42480
 ctgaggagtt caagccagcc tgggcaacac agtgagacca tcactctat aaaaaaatt 42540
 taaaaattag tgggtgtaa tggtttgac ctgtagtccc agctgccggg aggtgaagt 42600
 gggaggatga ctgagcccc ggaggcggag gccacagtga gctgtgtca agccactgca 42660
 ctccagcctg gatgacagag ccagatcctg tctcaaaaaa acaaaaaaca acaacaaca 42720
 acaaagataa atcactcaat acatcagcaa gagaaaaagc tctctgaaa tagtcacatg 42780
 caaagaaatt gaattccctc cagtcagaaa gagccactaa agtcctgag aatatctgat 42840
 cgattcaat gtcaggttt gagaggttt ttaaaaacag tticagatgt ttcttactat 42900
 ttttggcag aacatctgcc atctgcttc ttctctcct acatcttga actagacggg 42960
 gaattaataa ctacagaaaa ataacatgt atatgtaact ttactaaaa agaatacatca 43020
 aagtgtggtg agaagaagg gcagatttaa aagtttatg aagcgttcat tttaagcctc 43080
 ctaattatt ctgaaaaac aaaacacacc acttttctg actgcagcac tggtagggg 43140
 tgcataatcac gggtgactgt gatgattgt gcctggcgct taattttaa agttagtact 43200
 gagtgatgac agagagaggt caatgccact caaagaatat ttgtttgtt tgtttgtt 43260
 ttgtttgtt tgagaggcag tcttactctc tcactcaggc cggagtgcag tgggtgata 43320
 tcggctcact gcaacctctg cctcctgggt tcaagtatt ctctgcctc agcttctga 43380
 gtagctggga ttacaggcgt ctgccaccac atccagctaa ttttgtatt tttagtagag 43440
 atggggtttc atcatgttg ccaggctggg ctcaaactcc tgacctgagg tgatctgct 43500
 acctggcct ccaaagtgc tgggattaca ggcgtgagcc attgcacctg gccagaaga 43560
 atttattaca tactttccc aagagatggg gccacaccac accataccag gccacaccag 43620
 gccatgccat gtagacagg gccacaggag gaagtaccag atccgctcaa gaggcagaat 43680
 aaagggtaaa gaatgtcca gagctttatt gtgttactca gtggaagggc aaggcaggac 43740

tagggaaaca gcttaggggtt aactactttg aataatgccca gtaggttctg agttacagga 43800
 atggtctcta aatgtctggc cctcacccta cctgtcccta aggagaaata ctggagagtt 43860
 agaaaaggag gtggttgagg gattggttgg agggtttga atatgattt cacaccctca 43920
 caaaagctgg attgcagagg agatgtaaac aacttcagcc ttgggaggcc aagatgaaag 43980
 gatggctga agccaggagt tcaaggctgc aacaagctat gattacacca ctgcactcca 44040
 gcctgggtga aggaatgaga tcctgattct acaaaacatt ttgaaaaaa actttttat 44100
 tttttttt cttttttt tttttttt ttttgagac agtcttgc tcaccaccag 44160
 gctagagcgc agtgatacga tctcggctca ctgcgacctc catctcctgg gttaagtga 44220
 ttctcgtgcc tcagcgtcct gagtagctgg gattacaggt gcctgccacc atgcacggct 44280
 aatttttga ttttagtag agatgggggt tcacatggt gtccgggctg gtctgaact 44340
 cctgacctca agtgatccac ccgctcggc ctcccaaagt gctgggatta cagggtgtgag 44400
 ccacgggtcc cgacaaaaa atttaaaat aaaaattagc cacctgtagt ggtgcacgct 44460
 ttagtccca ccagctact tgggaggctg aggtgggagt atcacttggg cccaggaggt 44520
 ggaggctgca gtgagctctg atcatgccag tgctctcaa cctgggtgac agaacaagac 44580
 cccacctcaa aacaaaacaa acaaaaaaac acaactcag ccttaggtt ggccctgtga 44640
 ttaatgattg ccaaataggc atacacagaa tctaagaaaa tacagtgtgc tgaggtgtgc 44700
 ctgtctccag tccagtaatt agtatgcaag atttaccacc gccactccc actttgtct 44760
 agatgccaaa cctctcttc ccccttaagg aatagtcata ttgctgaag tttttttt 44820
 ttaatttctc tgcctgtgt taactcctgt ttggtttaaa atgtgcattt taactttaag 44880
 cgacaagctg atttccgtc actctgagat gatgcaggta cagggtgtga catggggagg 44940
 ggggactgct ctttggtcc aggtggttgg agagagacc agggcttgg attgttcc 45000
 ttccctgcc acctgtcaca gagccagggg acagaccagg aggactagac aggccactgt 45060
 ttggctttt cttccataa aataccagca tttttgcca gatgcagtgg ctcatgcctg 45120
 taaccccaaa actttgggag cctgaggcag gcaaatcact tgagctcagg agttcgagac 45180
 cagcctgggc aacatggcaa aagccgtct ctacaaaaa tacaaaaat tagccgagca 45240
 cgggtgtgca tgcctgtagt ccagctacc tgggagactg aggtgggaga atcatctgag 45300
 cctagcaagt caaggctgca gtgagtgtga atcctgtac tgactccag cctgggtgac 45360
 agcgtgagac cctgtctca aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa ccccaaaacc 45420
 cagcacttcc aaagggatct tacaataca gatcctttc ttctacaga tacgcaagt 45480
 actacaagga aaacaatgtt gagaaacgga ctctgataaa agtcttcggg atccgtttg 45540
 acatcctggt tttggcacc gtaagtctg ttcccagct ccgggcaccg gcatcctatg 45600
 actgtgtcct aattactgct gtggggcctc catggaggga agggttttg tctagcctt 45660
 cagctagcac tgggcatttc gcacatggga taaaagagga agaagatgtt ctcgggctgc 45720
 tgccctgaa tgagtcattt gacgagtaca ggagtgggcc tggaaagcaa ttctaactt 45780
 cggctttaaa aatactctga tacttaacaa gagagagaaa gaaatctctc tgtaaaatag 45840

ttgtaagta acagaatcca attcaaatg gcttaagggg aaaaaaaaaa gttttgagtg 45900
 ggctttttg gggtcacgta actgaaaaaa tccaggggtt attttaagc tacaagcatg 45960
 gttggatcca gggtgtcaaa caacaacatc atcatcaaga acttgccacc ttccgtgtct 46020
 tatatactgc ctccittggtc ttggcagcca tctctctatg aaggatcatg gcagcacctt 46080
 cgaatcgacc agtttagcaa cctcagtga aagaaaacac ctctttgccg gaagggtcag 46140
 caaagggtc agacaaatgt cattggctct gattggccca ctatggatga catgcccact 46200
 gctgaactaa tcaatgtgac cagggtcatg ccatactccc agtggttagt gtcgaaccag 46260
 ccctaacctg gccagggggc cagcagagag gccaacaccc cagccccttc aacacacaca 46320
 cacacacaca cacacacaca cactacatgg actgaaaaga aaattggagt 46380
 gtgccttct gaagatggaa ggaaaagatg ccagctatgc aaaaatcaac atatccacg 46440
 acatcctgtc caagagcatg ttatctatgt aacacagtga taagcaggac tagaagcaat 46500
 aagatatagc tgagagaatg caaagaccgc tgggaataaa aaccacaagg ctgggccggg 46560
 cagggtggct catgcctgta atcccagctc ttggggaggc tgaggcagggt ggatcacaag 46620
 gtcaggagct caagaccagc ctggccaaca tggtgaaacc ctgtctctac taaaaatgca 46680
 aaaaattagc tgggcgtggc ggtgggtact tgaatccca gctacttggg aggctgaggc 46740
 aggagtatct cttgaaccca ggaggcggag cttgcagtga gccgagattg cgccactgca 46800
 ctccagcctg ggcaaaagtg cgagactctg tctcaaaaaa aaaaaaaaaa aaaaaccac 46860
 gagggccacc agacattctt tccctctgct tctgaagtaa tgtcaaacct ttggaggag 46920
 aagcttgtct gaaacctca cctgtatgg agaaatgata gcccttcagt gggtctcccc 46980
 ttaatgtgtg ggccccagtg cttgctcatg taaaacctt atgggtccaa gcacaccctg 47040
 cacggctgaa gcaggatgcc tgagagtcag ttccagtctg cgtagtctct gccicagcca 47100
 gcactgaac gcatctacc aagtcacagc atgaggctcc gctccctgat agaaccaaca 47160
 attgcacgtt gaagcaaaag agcgtgtctc tgaattcac ctgagtaaac tctccactc 47220
 tgttttagg gaggaatg tgacattatc cagctgggtg tgtacatcg ctaaccctc 47280
 tctacttcg gtctggaag agattctctt ttccatgctt taggaaaatg gtttgagaa 47340
 ggaagtgact aacgcagcgc ttgtctgat tctccccagg ccgtgtgtt catcgactc 47400
 ctcatcgaca ctactccag taactgtgt cgtcccata ttatccctg gtgcaagtgc 47460
 gtgcagccct gtgtggtaaa cgaatactac tacaggaaga agtcgagtc cattgtggag 47520
 ccaagccgg tgaggccgt gtgttcacag gacaccaaga catggagaga ttccatgaaa 47580
 tcactcagaa atgcacgaaa attaggccca aatcacaggc ttatcctgt agtgagatg 47640
 tcgtgggtt ctacccgat caaccaactc tcagataaat ttttggtct tagagaagaa 47700
 tggaacaaa aatggagggg caagatagag ggaaggcaaa ttttatgtc taggacttg 47760
 caatttgc-attattat ttattatt attattat ttattatt attatttc 47820
 agatggagtc ttgtctatc acccaggctg gactgcagag gcatgatctc ggctcactgc 47880
 agcctctgcc tccagggtt aagtattct cctgcctcaa cctccaagt agctgggact 47940

acaggggcac accaccacac ctggctaatt tttgtatit tagtagagat ggtgtttage 48000
 tatgttgcc aggctggtct caaactcctg acttcaaat atccaccac tggcctccc 48060
 acagtgtgg gattataagt gtgagccact gagcccagcc tgtttgtca ttattaaat 48120
 tggtagacc aaaaaagaaa aaagaaaaga aaaataacaa ctttgagaa caatttgca 48180
 gtgactaatg tttaaatggg acatacttta cagcctggca tttcagttc tccatacctg 48240
 ccctagagaa acactcacat gagtaccgc agtacctgag tacaagatgt tcaagcagg 48300
 attgttatt aaattatcaa taatatatga ttattaacag tgaagaaata gcactaacc 48360
 aaatatcaa cagggtgaatg gtgaactat ggtaaatcca tagaaaggaa taccaggcac 48420
 cagcttaaaa aaatgagata gataataata atggcaaaca ttacacagc acttctaat 48480
 gctttatcta ttaactcaat ctccacaaca acctgatgta gacagatgct actattatct 48540
 cttctatat atagggaaat tgagccacag aaagggtaaa taattggccc aaggctgggc 48600
 acagtggctc acgcctataa tccaacact ttgggaggcc gaggcaggca gatcacttga 48660
 ggtcaagagt tcgagaccag cctggccaac atggtgaac cccatctcta ctaaaatac 48720
 aaaaattagc caggcgtggt ggtgcatgcc tgaagtcca gctactggg aggctgaggc 48780
 aggagaattg ctgaacctg gtaggcagag gttgcagtga gctgagatca tgccattgca 48840
 ctccagcctg agtgacagag ggaactctg tctcaaaaa aaaaaaaa aaaaattat 48900
 aggactgac aagagcagat ctccaagaca tttgttaa agaaaaaag caaactgtag 48960
 aacaatatat gttatatgat atataggta aaaaaaatc actagacaca gaagtaac 49020
 tgtatattt ctgcatgtat attctatata tacacaagt tacacacaca cacacataca 49080
 tatacatatg ttgggtgta tatatgta caaaattga ccagtggctg cttctgaaa 49140
 ggagcttagg gtgggggagt aggagtagc aaagagatt tagcgtatc tgtattgtt 49200
 tgattgatt aattcagact ttatcaagca ggtcctctgc gtcaactcc atgatttcc 49260
 ccaaagataa atctctgta ctaaaaaa aagacgatt gctagacgtg gtggctcacg 49320
 cctgtagtcc cagtacttg ggaggccaag gaggtggat cacctgaggt caggagtta 49380
 aaaccagcct ggccaacatg gagaacccc gtctctacta aaaatacaa aaattagcca 49440
 ggcatgggtg cgcatgcctg taatccagc ttcttagag gctgaggcag gagaatcgct 49500
 tgaaccagg aggtggaggt tttggtgagc caggattgca ccattgcact ccagcctggg 49560
 caacaagagc gaaactcct ctcaaaaa aaaaaaac aaagacgatt tcttgtctt 49620
 tccctatcc aagaacatga ttgtcctgtt ccagcagctg atgcacaatt cactgtccat 49680
 tgtatatgca ttcaaatg gaaataaag ttcatcttg cagctaaaac taatcaccac 49740
 ttcatggccc aagatgagat gaaattaac aaacatgta ataattaa ttgcaatagt 49800
 acaaattct ggagatactg aatctagagt tactgaaat gacagaatac acaaagaaa 49860
 tttatgcag caactggggg gtccaatga aaacattaa gcagtaagct gtggctgtgt 49920
 tgaattaca agtaagatg catgggggtc gcctggcgt ggtggctcac tctgtaac 49980
 ccagcacttt gtagggccaa ggcgggcgga tcaactgagg tcaggagttc aagaccagcc 50040

tgaccaacat ggagaaaccc cgtcttact aaaaatacaa aattagctgg gcatgatggt 50100
 gcatgcctgt aatccagct actcaggagg ctgaggcagg agaatactt gaacctggga 50160
 ggaggagggt gcagttagcc aagatcatgc cattgcactc cagcctggca acaagagtga 50220
 actccaactc aaaaaaaaaa aaaaaaaaaa agcatggggt tccattctg attatctt 50280
 agactcagaa atcattaatt ctgggtaat gagagtttg agccagcttg ttaatagtc 50340
 tatcatttg caaataggaa ttacagtgc cttagatag gcaattctg ataattctg 50400
 acaaaaatgg gtaaacctt aaaccatct ttcttagaca ttaaagtatg tgcctttg 50460
 ggatgaatcc cacattagga tgggaacca gcagctacta gggagaagtc tgcaagatgt 50520
 caagggccaa gaagtccag taagttaaatt cttttgtct tttttttt ttaagaaa 50580
 ttactgtta aatataaca catctagaaa ctgtacaaa tcaaaactga tggatttaa 50640
 caaagtaaac atactcatat aaccggcact cagattaaac aattgaaaat tactagcagg 50700
 agtcccttt atgccccct ccaatcacta cctctcttt ctctttttt aatttttaa 50760
 atttggttg gcacggtggc tcacacctgc aatccagca cttgggagg ccaagggtgg 50820
 tggatcact gagtcagga gtcgagacc aacctagcca acatggtgaa acctcgtctc 50880
 tactaaaaat acaaaaatta gccgggtggt gtggtggcac acatgtaac ccagctactc 50940
 agaagactga ggcaggagaa ctgctgaac ccaggagggtg gaggttcag tgagccgaga 51000
 ttgtgccatt gcactccagc ctgggtgaca gagcaagact ccatctcaa aaaaaaaag 51060
 aaaaaaatgt atattctaa attacaaacg aggtctcact atgtgtcca ggctggtctc 51120
 aagcagtcct ccacctcag ctgtaccaag cccaccaact gccctctctt aaaagtagtc 51180
 attatctgt ttcaaagat taattttact ttggctagaa ttctaaaa actgaatcac 51240
 gtagtatgta agcggtatat acgcggtga gtgtctggt tctttactc aacattatt 51300
 ttgtgagagt tgtcatgat gccatgata gttcattctc attgtataat tctgtttat 51360
 aaatatcaa cttatcagc catctactg ttgatggaca ttgggtagt gtccagttg 51420
 gggctaagc caataacgct gctatgctca acatagga ctctactgga cagttacct 51480
 agagtggaa ttgtgagta taaggcagac atatgttcgg tttaggaga tactaaciaa 51540
 cgggtctgaa aaatggtgt tcacattgc actctccca gcagttctg ttgtgggca 51600
 tctcaattt ctagggctg aattaccaca aactaagtgg cttaaaaca cagaaatggc 51660
 caggcatggt ggctcatgcc tgtaatcga gcactttggg aggtgaggc aggcagatca 51720
 cctgaggtca ggagtttag accagctga ccaacatgga gaaacccat ctctactaaa 51780
 aataaaaaa aatagccagg catggtgga ggcgactga gtccagcca ctcaggaggc 51840
 tgaggcagga gaatggcatg aacctgggag gcagagctg cagttagcca agattgcacc 51900
 actttgagag gccagggtg gcggatcacc tgaggtcagg agttcaggaa cagcctggcc 51960
 aacatggca aacctgtct ctattaaaa taaaaaatt agccaggcgt ggtggtgac 52020
 acctgtaac ccagctactc aggacgtga ggtgggagaa ttgctgaac ctgggaggca 52080
 gaggctcgg ttgctgaga tcacacctc gcactctagc ctgggcaaca gaacaagtc 52140

catctaaaaa aaaaaaaaaa aaaaagtctt ttctgctcct ttgaaatat cagttagttt 52200
tctcttttt ttctgttagt tgaattgtac tgattgattt tcaaatatta agccaaactt 52260
gcattcctga agtaaaccca atttgaatgt gttgtactat tctttgtatt tattgtctgaa 52320
ttccattcac taatatttag gatatttaca tctcttcttg agaaagactg accaaagtgt 52380
ttccattctt gtaatgttct tgttgattt gtgtatgaag tgaactacag tcatgcatca 52440
ctaataatg gggatatgtt ctgaaaaacg catcagtagc tgattctgtg gttgtctgaa 52500
tatcatggac tctatttaca caaacctaaa tagaatagcc tattatactt aggttatatg 52560
gtgtagtcta ttgctcctag gctgcaaacc tgtacagcat gttactgtac tgaatacggg 52620
aagcaactgt aacagaatgg taagatttgt atatctaaat atagaaaagg tacagtgaag 52680
alatgatata aaagcttaaa aatggtagat ctgcataggg cacttaccat gaatggagct 52740
tgtaggactg aaagtgtctc tgggtgagtc agtgagtagt gagtgaatgt gaaagcctag 52800
gtgtttaccg tgcactacag tagacttcat aaacactgta cacttaggct acactaaatt 52860
tacttcaaaa tattttactt tcttcaataa taaattaatc ttacttact gtgactgttt 52920
tactttataa atttttaaat tttttaaact ctgtacagtg gtataaaaat attttcttc 52980
tcaccgggag tgggtggctca tgcctgtaat cccatcactt tgggaggccg aggcaggcgg 53040
atcacaagat caggagattg agaccatcct ggccaacatg gtgaaacccc atctctacta 53100
aaaatacaaa aaaatagcca ggcatgggtg caggcgctg tagtcccagc tactcgggag 53160
gctgaggcag gagaatggcg tgaaccagg aggcagagct tgcagtgagc caagatcaca 53220
ccactgcact ccagcctggg cgacagagca agactccatc tcaaaaaaaaa aaaaattctt 53280
tctctatc cttattctat atacttttt ctattttta catttttat tttattttt 53340
actttttaa tatttttgtt aaaaactaag tcatggccgg gtgcagtggc tcacgcttgt 53400
aatctcagta ctttgggagg ctgagggtgg tggatcactc gaggtcagga gttcaacacc 53460
agcctggcca atatggtgaa actctgtctc tactaaaaat ataaaaatta gccgggtgtg 53520
gtggtgcacg cctgtagtcc cagctactca ggaggctgag gcagaagaat cgcttgaacc 53580
caggaggcgg aggttgagct aagccaagat catgccactg cactccagcc tgggagacag 53640
agcaagactc catctcaaaa aaaaaaaca acaacaaca cacacacaca cacacacaca 53700
aaaaccattt ccaatagtgc taagtcttat gaagaatgtg gaataccaca gtgtgataag 53760
ggaatcatgt gggagaaaag ctgctagata ggggtgtcag gacaagaggt gacatctcaa 53820
cagaggcctg gcttttctga atttcattt ccaaaatctg taaaataggc caggtgcagt 53880
ggctcatgcc tgtaatccca gcactttgag aggctgaggc aggtggatca cctgaggtca 53940
ggagttccaa acaagcctgg ccaacatgtt gaaatctgt ttactactaga aatacgaag 54000
aattagctgg gcatgggtgc atgcacctgt aataccagct acttaggagg ttaaggcatg 54060
aaaattgtct gaacctggga ggtgaaagt gcagtgagcc aagatcacac cactgtgctc 54120
cagcctgtgc gacagagtga gacctctct caaaaaaaaa aaaaaaatct gtaaaataaa 54180
gacaaggata cattatctca caagcgtctt caaaggcctg aatgaggcaa tgcttacaga 54240

acacatgcat ggtcctgata tctacaccta ataatgacg gctactataa atcatgtaat 54300
 attaaacgta actttataag ttaataaaat taaagaacct agaacctgag ggcttgcat 54360
 ggctaatagg ttggaaact tgcttttca gagacctgcg atggacttca cagattgtc 54420
 caggctgccc ctggccctcc atgacacacc cccgattcct ggacaaccag aggagataca 54480
 gctgcttaga aaggaggcga ctccatgac caggatagc cccgtctgt gccagtgtgg 54540
 aagctgcctc ccatctcaac tcccfagag ccacaggcgc ctggaggagc tgtgctgccg 54600
 gaaaaagccg ggggcctgca tcaccacctc agagctgttc aggaagctgg tcctgtccag 54660
 acacgtcctg cagttctcc tgctctacca ggagcccttg ctggcgtgg atgtggattc 54720
 caccaacagc cggctgcggc actgtgcta caggtgtac gccacctggc gcttcggctc 54780
 ccaggacatg gctgacttg ccaacctgcc cagctgtgc cgctggagga tccggaaaga 54840
 gttccgaag agtgaaggc agtacagtgg ctcaagagt ccttactgaa gccaggcacc 54900
 gtggctcacg tctgtaatcc cagcgcttg ggaggccgag gcaggcagat cacctgagat 54960
 cgggagtgg agacccgcct ggctaacaag gcgaaatcct gtctgtacta aaaatacaaa 55020
 aatcagccag acatgggtgc atgcacctgc aatcccagct actcgggagg ctgaggcaca 55080
 agaatactt gaacccggga ggcagagggt gtagtgagcc cagattgtc cactgtctc 55140
 cagcctggga ggcacagcaa actgtccca aaaaaaaaaa aagatcctt accaatagca 55200
 ggggctgcag tagccatgtt aacatgacat ttaccagcaa ctgaactt acctgcaaag 55260
 ctctgtggc acatttcag ccaaggga atatgcttc atctctgtt gctctgtg 55320
 tctgagagca aagtacctg gttaaaaaa ccagaatccc tctacatgga ctgagaaaa 55380
 agagattgag atgtaagtct caactgtgc ccaggaagt tgttgaccc taggcctc 55440
 acctgtgc ctctgtcc ttgtgccca actactatct cagagatatt gtgaggacaa 55500
 attgagacag tgcacatgaa ctgtcttta atgtgtaaag atctacatga atgcaaaaca 55560
 ttcatatg aggtcagact aggataatgt ccaactaaaa acaaacctt tcatcctg 55620
 ctggagaatg tggagaacta aagggtggca caaattctt gacactcaag tccccaaaga 55680
 cctaagggtt ttatctctc ccttgaata tgggtggctc tgattgctt atcaaaaagt 55740
 ggaagtgaca ttgtgtcagt ttcagatcct gatcttaaga ggctgacgc ttctactgc 55800
 tgcccttg aactctgt atcggggaag ccagacgcca tttaaagtc tgcctatcct 55860
 ggccagggtg ggtggctcac acctgtaac ccagcactt gggagacaa ggcggcgga 55920
 tcaattaaag tcaggagtcc aagaccagac tcgccaacat ggtgaaaccg tatcttaat 55980
 aaaaatacaa aaattgctg ggcattgtgc gggcacctgt agtctagct atcaagaggc 56040
 ttagacagga gaaacactg aacctgggag gtggagggtg cattgagctg agatcgtgcc 56100
 actgcactcc aggtgggtg acagagcgag actccatctc aaaaaaaaaa aaaaagaaa 56160
 aaaaaatgt ctgcctatcc tgagactgcc ctgctgtgag gaagcccaag cagtcactg 56220
 gacagtgcct gaccagcccc agcttcaag ccatccaagc ccagtcacca aacatgagag 56280
 agaagaagcc ttcaggtgat tctggactcc actaacatat gactgatacc gcatgataca 56340

tcccaagtga gaactgcccc ataatccag aaaaccacat tgctatctta agtccctaag 56400
 ttgggggctt attgttcca cagcaacagg taactggaac agagggaag cctgatgaat 56460
 gggcacacag actcagccca taccttcctt ggttctaag ttctcagga gcccgacca 56520
 accctgggag cctcaggaac ttaggttcc actggacagt tctagaagg ctatagacca 56580

<210> 2
 <211> 2164
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (79)..(1863)
 <223>

<400> 2
 ggcacgaggg ctgtgtgg ccctgtcagg aagagtagag ctctgtcca gtcgcgca 60

gggagggagg ctgtcacc atg ccg gcc tgc tgc agc tgc agt gat gtt ttc 111
 Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe
 1 5 10

cag tat gag acg aac aaa gtc act cgg atc cag agc atg aat tat ggc 159
 Gln Tyr Glu Thr Asn Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly
 15 20 25

acc att aag tgg ttc ttc cac gtg atc atc ttt tcc tac gtt tgc ttt 207
 Thr Ile Lys Trp Phe Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe
 30 35 40

gct ctg gtg agt gac aag ctg tac cag cgg aaa gag cct gtc atc agt 255
 Ala Leu Val Ser Asp Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser
 45 50 55

tct gtg cac acc aag gtg aag ggg ata gca gag gtg aaa gag gag atc 303
 Ser Val His Thr Lys Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile
 60 65 70 75

gtg gag aat gga gtg aag aag ttg gtg cac agt gtc ttt gac acc gca 351
 Val Glu Asn Gly Val Lys Lys Leu Val His Ser Val Phe Asp Thr Ala
 80 85 90

gac tac acc ttc cct ttg cag ggg aac tct ttc ttc gtg atg aca aac 399
 Asp Tyr Thr Phe Pro Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn
 95 100 105

ttt ctc aaa aca gaa ggc caa gag cag cgg ttg tgt ccc gag tat ccc 447
 Phe Leu Lys Thr Glu Gly Gln Glu Arg Leu Cys Pro Glu Tyr Pro
 110 115 120

acc cgc agg acg ctc tgt tcc tct gac cga ggt tgt aaa aag gga tgg 495
 Thr Arg Arg Thr Leu Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp
 125 130 135

atg gac ccg cag agc aaa gga att cag acc gga agg tgt gta gtg cat 543
 Met Asp Pro Gln Ser Lys Gly Ile Gln Thr Gly Arg Cys Val Val His
 140 145 150 155

gaa ggg aac cag aag acc tgt gaa gtc tct gcc tgg tgc ccc atc gag 591
 Glu Gly Asn Gln Lys Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu
 160 165 170

gca gtg gaa gag gcc ccc cgg cct gct ctc ttg aac agt gcc gaa aac 639
 Ala Val Glu Glu Ala Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn
 175 180 185

ttc act gtg ctc atc aag aac aat atc gac ttc ccc ggc cac aac tac 687
 Phe Thr Val Leu Ile Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr
 190 195 200

acc acg aga aac atc ctg cca ggt tta aac atc act tgt acc ttc cac 735
 Thr Thr Arg Asn Ile Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His
 205 210 215

aag act cag aat cca cag tgt ccc att ttc cga cta gga gac atc ttc 783
 Lys Thr Gln Asn Pro Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe
 220 225 230 235

cga gaa aca ggc gat aat ttt tca gat gtg gca att cag ggc gga ata 831
 Arg Glu Thr Gly Asp Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile
 240 245 250

atg ggc att gag atc tac tgg gac tgc aac cta gac cgt tgg ttc cat 879
 Met Gly Ile Glu Ile Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His
 255 260 265

cac tgc cat ccc aaa tac agt ttc cgt cgc ctt gac gac aag acc acc 927
 His Cys His Pro Lys Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr
 270 275 280

aac gtg tcc ttg tac cct ggc tac aac ttc aga tac gcc aag tac tac 975
 Asn Val Ser Leu Tyr Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr
 285 290 295

aag gaa aac aat gtt gag aaa cgg act ctg ata aaa gtc ttc ggg atc 1023
 Lys Glu Asn Asn Val Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile
 300 305 310 315

cgt ttt gac atc ctg gtt ttt ggc acc gga gga aaa ttt gac att atc 1071
 Arg Phe Asp Ile Leu Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile
 320 325 330

cag ctg gtt gtg tac atc ggc tca acc ctc tcc tac ttc ggt ctg gcc 1119
 Gln Leu Val Val Tyr Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala
 335 340 345

gct gtg ttc atc gac ttc ctc atc gac act tac tcc agt aac tgc tgt 1167
 Ala Val Phe Ile Asp Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys
 350 355 360

cgc tcc cat att tat ccc tgg tgc aag tgc tgt cag ccc tgt gtg gtc 1215
 Arg Ser His Ile Tyr Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val
 365 370 375

aac gaa tac tac tac agg aag aag tgc gag tcc att gtg gag cca aag 1263
 Asn Glu Tyr Tyr Tyr Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys
 380 385 390 395

ccg aca tta aag tat gtg tcc ttt gtg gat gaa tcc cac att agg atg 1311
 Pro Thr Leu Lys Tyr Val Ser Phe Val Asp Glu Ser His Ile Arg Met
 400 405 410

gtg aac cag cag cta cta ggg aga agt ctg caa gat gtc aag ggc caa 1359
 Val Asn Gln Gln Leu Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln
 415 420 425

gaa gtc cca aga cct cgg atg gac ttc aca gat ttg tcc agg ctg ccc 1407
 Glu Val Pro Arg Pro Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro
 430 435 440

ctg gcc ctc cat gac aca ccc ccg att cct gga caa cca gag gag ata 1455
 Leu Ala Leu His Asp Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile
 445 450 455

cag ctg ctt aga aag gag ggc act cct aga tcc agg gat agc ccc gtc 1503

Gln Leu Leu Arg Lys Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val
460 465 470 475

tgg tgc cag tgt gga agc tgc ctc cca tct caa ctc cct gag agc cac 1551
Trp Cys Gln Cys Gly Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His
480 485 490

agg tgc ctg gag gag ctg tgc tgc cgg aaa aag ccg ggg gcc tgc atc 1599
Arg Cys Leu Glu Glu Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile
495 500 505

acc acc tca gag ctg ttc agg aag ctg gtc ctg tcc aga cac gtc ctg 1647
Thr Thr Ser Glu Leu Phe Arg Lys Leu Val Leu Ser Arg His Val Leu
510 515 520

cag ttc ctc ctg ctc tac cag gag ccc ttg ctg gcg ctg gat gtg gat 1695
Gln Phe Leu Leu Leu Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp
525 530 535

tcc acc aac agc cgg ctg cgg cac tgt gcc tac agg tgc tac gcc acc 1743
Ser Thr Asn Ser Arg Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr
540 545 550 555

tgg cgc ttc ggc tcc cag gac atg gct gac ttt gcc atc ctg ccc agc 1791
Trp Arg Phe Gly Ser Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser
560 565 570

tgc tgc cgc tgg agg atc cgg aaa gag ttt ccg aag agt gaa ggg cag 1839
Cys Cys Arg Trp Arg Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln
575 580 585

tac agt ggc ttc aag agt cct tac tgaagccagg caccgtggct cagctctgta 1893
Tyr Ser Gly Phe Lys Ser Pro Tyr
590 595

atcccagcgc ttggggaggc cgaggcaggc agatcacctg aggtcgggag ttggagaccc 1953

gcctggctaa caaggcgaaa tctgtctgt actaaaaata caaaaatcag ccagacatgg 2013

tggcatgcac ctgcaatccc agctactcgg gaggctgagg cacaagaatc acttgaaccc 2073

gggaggcaga ggtgtagtgt agcccagatt gtgccactgc tctccagcct gggaggcaca 2133

gcaaaactgtc ccaaaaaaaaa aaaaaaaaaa a 2164

<210> 3
<211> 595
<212> PRT
<213> Homo sapiens

<400> 3

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val

65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys His Pro Lys
260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
 420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
 435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
 515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
 530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
 580 585 590

Ser Pro Tyr
 595

<210> 4
 <211> 595
 <212> PRT
 <213> Homo sapiens

<220>
 <221> DOMAIN
 <222> (1)..(20)
 <223> Intracellular domain

<220>
 <221> TRANSMEM
 <222> (21)..(46)
 <223>

<220>
 <221> DOMAIN
 <222> (47)..(320)
 <223> Extracellular domain

<220>
 <221> TRANSMEM
 <222> (321)..(356)
 <223>

<220>
 <221> DOMAIN
 <222> (357)..(595)
 <223> Intracellular domain

<400> 4

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
580 585 590

Ser Pro Tyr
595

<210> 5
<211> 595
<212> PRT
<213> Homo sapiens

<400> 5

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro

85

90

95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110

Gly Gln Glu Gln Trp Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
 290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
 340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
 420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
 435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
 515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
 530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
 580 585 590

Ser Pro Tyr
 595

<210> 6

<211> 595

<212> PRT

<213> Homo sapiens

<400> 6

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
130 135 140

Lys Gly Ile Gln Thr Arg Arg Cys Val Val His Glu Gly Asn Gln Lys
145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
 290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
 340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
 420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
 435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
 515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
 530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
 580 585 590

Ser Pro Tyr
 595

<210> 7
 <211> 595
 <212> PRT
 <213> Homo sapiens

<400> 7

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Lys Asn Phe Thr Val Leu Ile
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro

210

215

220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
 290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
 340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
 420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
 435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
 515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
 530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
 580 585 590

Ser Pro Tyr
 595

<210> 8
 <211> 595
 <212> PRT
 <213> Homo sapiens

<400> 8

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Pro Ile
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
 290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
 340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
 420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
 435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
 515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
 530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
 580 585 590

Ser Pro Tyr
 595

<210> 9
 <211> 595
 <212> PRT
 <213> Homo sapiens

<400> .9

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Cys Pro Lys
260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp

47

340

345

350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
580 585 590

Ser Pro Tyr
595

<210> 10
<211> 595
<212> PRT

<213> Homo sapiens

<400> 10

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
 290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
 340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
 420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
 435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
 450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
 465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
 485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
 500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
 515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
 530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
 545 550 555 560

Gln Asp Met Ala Asp Phe Ala Asn Leu Pro Ser Cys Cys Arg Trp Arg
 565 570 575

Ile Arg Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
 580 585 590

Ser Pro Tyr
 595

<210> 11
 <211> 588
 <212> PRT
 <213> Homo sapiens

<400> 11

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
405 410 415

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly

465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Glu Glu Leu Cys Cys Arg Lys Lys Pro
 485 490 495

Gly Ala Cys Ile Thr Thr Ser Glu Leu Phe Arg Lys Leu Val Leu Ser
 500 505 510

Arg His Val Leu Gln Phe Leu Leu Leu Tyr Gln Glu Pro Leu Leu Ala
 515 520 525

Leu Asp Val Asp Ser Thr Asn Ser Arg Leu Arg His Cys Ala Tyr Arg
 530 535 540

Cys Tyr Ala Thr Trp Arg Phe Gly Ser Gln Asp Met Ala Asp Phe Ala
 545 550 555 560

Ile Leu Pro Ser Cys Cys Arg Trp Arg Ile Arg Lys Glu Phe Pro Lys
 565 570 575

Ser Glu Gly Gln Tyr Ser Gly Phe Lys Ser Pro Tyr
 580 585

<210> 12
 <211> 595
 <212> PRT
 <213> Homo sapiens

<400> 12

Met Pro Ala Cys Cys Ser Cys Ser Asp Val Phe Gln Tyr Glu Thr Asn
 1 5 10 15

Lys Val Thr Arg Ile Gln Ser Met Asn Tyr Gly Thr Ile Lys Trp Phe
 20 25 30

Phe His Val Ile Ile Phe Ser Tyr Val Cys Phe Ala Leu Val Ser Asp
 35 40 45

Lys Leu Tyr Gln Arg Lys Glu Pro Val Ile Ser Ser Val His Thr Lys
 50 55 60

Val Lys Gly Ile Ala Glu Val Lys Glu Glu Ile Val Glu Asn Gly Val
 65 70 75 80

Lys Lys Leu Val His Ser Val Phe Asp Thr Ala Asp Tyr Thr Phe Pro
 85 90 95

Leu Gln Gly Asn Ser Phe Phe Val Met Thr Asn Phe Leu Lys Thr Glu
 100 105 110

Gly Gln Glu Gln Arg Leu Cys Pro Glu Tyr Pro Thr Arg Arg Thr Leu
 115 120 125

Cys Ser Ser Asp Arg Gly Cys Lys Lys Gly Trp Met Asp Pro Gln Ser
 130 135 140

Lys Gly Ile Gln Thr Gly Arg Cys Val Val His Glu Gly Asn Gln Lys
 145 150 155 160

Thr Cys Glu Val Ser Ala Trp Cys Pro Ile Glu Ala Val Glu Glu Ala
 165 170 175

Pro Arg Pro Ala Leu Leu Asn Ser Ala Glu Asn Phe Thr Val Leu Ile
 180 185 190

Lys Asn Asn Ile Asp Phe Pro Gly His Asn Tyr Thr Thr Arg Asn Ile
 195 200 205

Leu Pro Gly Leu Asn Ile Thr Cys Thr Phe His Lys Thr Gln Asn Pro
 210 215 220

Gln Cys Pro Ile Phe Arg Leu Gly Asp Ile Phe Arg Glu Thr Gly Asp
 225 230 235 240

Asn Phe Ser Asp Val Ala Ile Gln Gly Gly Ile Met Gly Ile Glu Ile
 245 250 255

Tyr Trp Asp Cys Asn Leu Asp Arg Trp Phe His His Cys Arg Pro Lys
 260 265 270

Tyr Ser Phe Arg Arg Leu Asp Asp Lys Thr Thr Asn Val Ser Leu Tyr
 275 280 285

Pro Gly Tyr Asn Phe Arg Tyr Ala Lys Tyr Tyr Lys Glu Asn Asn Val
 290 295 300

Glu Lys Arg Thr Leu Ile Lys Val Phe Gly Ile Arg Phe Asp Ile Leu
 305 310 315 320

Val Phe Gly Thr Gly Gly Lys Phe Asp Ile Ile Gln Leu Val Val Tyr
 325 330 335

Ile Gly Ser Thr Leu Ser Tyr Phe Gly Leu Ala Ala Val Phe Ile Asp
 340 345 350

Phe Leu Ile Asp Thr Tyr Ser Ser Asn Cys Cys Arg Ser His Ile Tyr
 355 360 365

Pro Trp Cys Lys Cys Cys Gln Pro Cys Val Val Asn Glu Tyr Tyr Tyr
 370 375 380

Arg Lys Lys Cys Glu Ser Ile Val Glu Pro Lys Pro Thr Leu Lys Tyr
 385 390 395 400

Val Ser Phe Val Asp Glu Ser His Ile Arg Met Val Asn Gln Gln Leu
 405 410 415

54

Leu Gly Arg Ser Leu Gln Asp Val Lys Gly Gln Glu Val Pro Arg Pro
420 425 430

Ala Met Asp Phe Thr Asp Leu Ser Arg Leu Pro Leu Ala Leu His Asp
435 440 445

Thr Pro Pro Ile Pro Gly Gln Pro Glu Glu Ile Gln Leu Leu Arg Lys
450 455 460

Glu Ala Thr Pro Arg Ser Arg Asp Ser Pro Val Trp Cys Gln Cys Gly
465 470 475 480

Ser Cys Leu Pro Ser Gln Leu Pro Glu Ser His Arg Cys Leu Glu Glu
485 490 495

Leu Cys Cys Arg Lys Lys Pro Gly Ala Cys Ile Thr Thr Ser Glu Leu
500 505 510

Phe Arg Lys Leu Val Leu Ser Arg His Val Leu Gln Phe Leu Leu Leu
515 520 525

Tyr Gln Glu Pro Leu Leu Ala Leu Asp Val Asp Ser Thr Asn Ser Arg
530 535 540

Leu Arg His Cys Ala Tyr Arg Cys Tyr Ala Thr Trp Arg Phe Gly Ser
545 550 555 560

Gln Asp Met Ala Asp Phe Ala Ile Leu Pro Ser Cys Cys Arg Trp Arg
565 570 575

Ile Gln Lys Glu Phe Pro Lys Ser Glu Gly Gln Tyr Ser Gly Phe Lys
580 585 590

Ser Pro Tyr
595

<210> 13
<211> 20
<212> DNA
<213> Homo sapiens

<400> 13
gtaacctcac tgagaacaga

20

<210> 14
<211> 20
<212> DNA
<213> Homo sapiens

<400> 14
tttggtaggc gaaggcgggc

20

<210> 15
<211> 21

| | |
|------------------------|----|
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 15 | |
| ccttgatgg gcttcgtgc t | 21 |
| <210> 16 | |
| <211> 21 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 16 | |
| ttgttatcc gtccatcagc t | 21 |
| <210> 17 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 17 | |
| actgccatat gtactattcc | 20 |
| <210> 18 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 18 | |
| aaatctaccc agccacttag | 20 |
| <210> 19 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 19 | |
| tgcatgtgcc cacagcacag | 20 |
| <210> 20 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 20 | |
| catcctccaa tgcctgcatc | 20 |
| <210> 21 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 21 | |
| ttcccatgt ttgccattta | 20 |
| <210> 22 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 22 | |
| caagagcagt ggttgtgtcc | 20 |

<210> 23
<211> 20
<212> DNA
<213> Homo sapiens

<400> 23
gagccacag atatctactg 20

<210> 24
<211> 20
<212> DNA
<213> Homo sapiens

<400> 24
tgcctgcctc agtctctggt 20

<210> 25
<211> 20
<212> DNA
<213> Homo sapiens

<400> 25
tctctcgccc aggttgagt 20

<210> 26
<211> 20
<212> DNA
<213> Homo sapiens

<400> 26
acgtcctctc cgcagttctt 20

<210> 27
<211> 20
<212> DNA
<213> Homo sapiens

<400> 27
attcagaccg aaaggtgtgt 20

<210> 28
<211> 20
<212> DNA
<213> Homo sapiens

<400> 28
tgcctggtgt cccatcgagg 20

<210> 29
<211> 20
<212> DNA
<213> Homo sapiens

<400> 29
gttgaacac tcctgtacca 20

<210> 30
<211> 20
<212> DNA
<213> Homo sapiens

<400> 30
aacagtgccca aaaacttcac 20

<210> 31
<211> 20
<212> DNA
<213> Homo sapiens

<400> 31
tcactgtgcc catcaagaac 20

<210> 32
<211> 20
<212> DNA
<213> Homo sapiens

<400> 32
tgctgcgtgc ttctggctca 20

<210> 33
<211> 20
<212> DNA
<213> Homo sapiens

<400> 33
ctggagaacc ctttccaag 20

<210> 34
<211> 20
<212> DNA
<213> Homo sapiens

<400> 34
caggaggcag ggatcattg 20

<210> 35
<211> 20
<212> DNA
<213> Homo sapiens

<400> 35
tccaccgct acgctaaggt 20

<210> 36
<211> 20
<212> DNA
<213> Homo sapiens

<400> 36
tgcgctaagg actttaccta 20

<210> 37
<211> 20
<212> DNA
<213> Homo sapiens

<400> 37
tttacctacc ctacctgctc 20

<210> 38
<211> 20
<212> DNA
<213> Homo sapiens

<400> 38
ccatcactgc tgtcccaaat

20

<210> 39
<211> 20
<212> DNA
<213> Homo sapiens

<400> 39
tcgccttgat gacaagacca

20

<210> 40
<211> 20
<212> DNA
<213> Homo sapiens

<400> 40
gtaaactctt ccactctgtt

20

<210> 41
<211> 20
<212> DNA
<213> Homo sapiens

<400> 41
accaagacac ggagagattc

20

<210> 42
<211> 20
<212> DNA
<213> Homo sapiens

<400> 42
taatattaaa tgtaacttta

20

<210> 43
<211> 20
<212> DNA
<213> Homo sapiens

<400> 43
aatattaaac ataacttat

20

<210> 44
<211> 20
<212> DNA
<213> Homo sapiens

<400> 44
ccatctcaac tggaggagct

20

<210> 45
<211> 20
<212> DNA
<213> Homo sapiens

<400> 45
gactttgcca tcctgcccaag

20

<210> 46
<211> 20
<212> DNA

<213> Homo sapiens
 <400> 46
 tggaggatcc agaaagagtt 20

 <210> 47
 <211> 20
 <212> DNA
 <213> Homo sapiens

 <400> 47
 aatcccagca ctttgggagg 20

 <210> 48
 <211> 20
 <212> DNA
 <213> Homo sapiens

 <400> 48
 aaactgtccc aaaaaaaaaa 20

 <210> 49
 <211> 20
 <212> DNA
 <213> Homo sapiens

 <400> 49
 aactgtcccc caaaaaaaaaa 20

 <210> 50
 <211> 20
 <212> DNA
 <213> Homo sapiens

 <400> 50
 ctgtccccac aaaaaaaaaa 20

 <210> 51
 <211> 20
 <212> DNA
 <213> Homo sapiens

 <400> 51
 caaggcgggt ggatcactta 20

 <210> 52
 <211> 19
 <212> DNA
 <213> Homo sapiens

 <400> 52
 cgtaggactt ggcgcttct 19

 <210> 53
 <211> 21
 <212> DNA
 <213> Homo sapiens

 <400> 53
 gagcacgtct cagattcgaa a 21

 <210> 54

<211> 23
 <212> DNA
 <213> Homo sapiens

<400> 54
 ccatgaggca ggtatgacta ttc

23

<210> 55
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 55
 ctcttgatc tcaccagtt

20

<210> 56
 <211> 23
 <212> DNA
 <213> Homo sapiens

<400> 56
 ctctgccagc ttgatatta agc

23

<210> 57
 <211> 22
 <212> DNA
 <213> Homo sapiens

<400> 57
 ggtccctagt gctagaacca ga

22

<210> 58
 <211> 18
 <212> DNA
 <213> Homo sapiens

<400> 58
 attcatccgt cagtgcc

18

<210> 59
 <211> 22
 <212> DNA
 <213> Homo sapiens

<400> 59
 gccatgtgaa ttctaccg at

22

<210> 60
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 60
 ttcgtgtgg ttaggatggg

20

<210> 61
 <211> 22
 <212> DNA
 <213> Homo sapiens

<400> 61
 caaggatgct caggtagta gc

22

| | |
|-----------------------------|----|
| <210> 62 | |
| <211> 25 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 62 | |
| cactaggtt gctgtatcca ttct | 25 |
| | |
| <210> 63 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 63 | |
| gcaactgtgt gagagcttg | 20 |
| | |
| <210> 64 | |
| <211> 19 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 64 | |
| tcaaccctgg tccagtgtg | 19 |
| | |
| <210> 65 | |
| <211> 25 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 65 | |
| caccaagtag ctctcactca taagg | 25 |
| | |
| <210> 66 | |
| <211> 24 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 66 | |
| caataacact tgtgcgagtt aggt | 24 |
| | |
| <210> 67 | |
| <211> 21 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 67 | |
| catctgttg ccttggaac c | 21 |
| | |
| <210> 68 | |
| <211> 23 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 68 | |
| gtgagtggtgta atcctgctac tgc | 23 |
| | |
| <210> 69 | |
| <211> 19 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 69 | |

aggcccactc ctgtactcg

19

<210> 70

<211> 19

<212> DNA

<213> Homo sapiens

<400> 70

ccaagtcaca gcatgaggc

19

<210> 71

<211> 18

<212> DNA

<213> Homo sapiens

<400> 71

acccagcgac gtatccac

18

<210> 72

<211> 19

<212> DNA

<213> Homo sapiens

<400> 72

aagcatgggg ttccattc

19

<210> 73

<211> 24

<212> DNA

<213> Homo sapiens

<400> 73

gcataaaaagg gactcctgct agta

24

<210> 74

<211> 21

<212> DNA

<213> Homo sapiens

<400> 74

gcttacagaa cacatgcatg g

21

<210> 75

<211> 19

<212> DNA

<213> Homo sapiens

<400> 75

gcacctgtag gcacagtgc

19

<210> 76

<211> 21

<212> DNA

<213> Homo sapiens

<400> 76

atcaccacct cagagctgtt.c

21

<210> 77

<211> 21

<212> DNA

<213> Homo sapiens

| | |
|---|----|
| <400> 77 gttaacatgg ctactgcagc c | 21 |
| <210> 78 <211> 21 <212> DNA <213> Homo sapiens | |
| <400> 78 gcttagaaag gaggcgactc c | 21 |
| <210> 79 <211> 22 <212> DNA <213> Homo sapiens | |
| <400> 79 ttgtgacatt tgcaaggctg cc | 22 |
| <210> 80 <211> 20 <212> DNA <213> Homo sapiens | |
| <400> 80 tctgaagctc tgctcctgag | 20 |
| <210> 81 <211> 20 <212> DNA <213> Homo sapiens | |
| <400> 81 ctcaccttct ggcttcagt | 20 |
| <210> 82 <211> 20 <212> DNA <213> Homo sapiens | |
| <400> 82 cttaccactc ccaggactaa | 20 |
| <210> 83 <211> 20 <212> DNA <213> Homo sapiens | |
| <400> 83 gtctgcctgt tcactgccat | 20 |
| <210> 84 <211> 21 <212> DNA <213> Homo sapiens | |
| <400> 84 cagagacctt cagaaacttc g | 21 |
| <210> 85 <211> 20 | |

<212> DNA
<213> Homo sapiens

<400> 85
agatcaccag ggacacagtg 20

<210> 86
<211> 20
<212> DNA
<213> Homo sapiens

<400> 86
ctcaactcca ctttctcgg 20

<210> 87
<211> 23
<212> DNA
<213> Homo sapiens

<400> 87
cctttcactt ttttgtctc atg 23

<210> 88
<211> 21
<212> DNA
<213> Homo sapiens

<400> 88
gggagaattc tgaaaatgcc c 21

<210> 89
<211> 20
<212> DNA
<213> Homo sapiens

<400> 89
ggaccagagc tctactcttc 20

<210> 90
<211> 20
<212> DNA
<213> Homo sapiens

<400> 90
aggatcataga tcgacctgcc 20

<210> 91
<211> 20
<212> DNA
<213> Homo sapiens

<400> 91
aagaagcgcc aagtcctacg 20

<210> 92
<211> 21
<212> DNA
<213> Homo sapiens

<400> 92
gcaatccaga ctgaagtga c 21

| | |
|----------------------------|----|
| <210> 93 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 93 | |
| actctggtct gcagttggtg | 20 |
| | |
| <210> 94 | |
| <211> 24 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 94 | |
| cctttaaaat cagagacctt caga | 24 |
| | |
| <210> 95 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 95 | |
| gccccatcctc tgaacaccat | 20 |
| | |
| <210> 96 | |
| <211> 21 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 96 | |
| cccttggaac tcttgctatc g | 21 |
| | |
| <210> 97 | |
| <211> 21 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 97 | |
| ggcagtacag tggctcaag a | 21 |
| | |
| <210> 98 | |
| <211> 21 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 98 | |
| gtgggacagt tigtgtgcc t | 21 |
| | |
| <210> 99 | |
| <211> 20 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 99 | |
| gagtccttac caatagcagg | 20 |
| | |
| <210> 100 | |
| <211> 21 | |
| <212> DNA | |
| <213> Homo sapiens | |
| <400> 100 | |
| gtcaaagaat ttgggccac c | 21 |

<210> 101
 <211> 26
 <212> DNA
 <213> Homo sapiens

<400> 101
 catgaactgt ctttaatgt gtaaag 26

<210> 102
 <211> 22
 <212> DNA
 <213> Homo sapiens

<400> 102
 gagatacggt ttcacatgt tg 22

<210> 103
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 103
 aattagctgg gcatggtgcg 20

<210> 104
 <211> 21
 <212> DNA
 <213> Homo sapiens

<400> 104
 ttgagatgga gtctcgctct g 21

<210> 105
 <211> 20
 <212> DNA
 <213> Homo sapiens

<400> 105
 cactgtccac gtgactgctt 20

<210> 106
 <211> 25
 <212> DNA
 <213> Homo sapiens

<400> 106
 tcctacttcg gtctggtaag agatt 25

<210> 107
 <211> 19
 <212> DNA
 <213> Homo sapiens

<400> 107
 gggcctaatt ttcgtgcat 19

<210> 108
 <211> 23
 <212> DNA
 <213> Homo sapiens

<400> 108
aagaacctag aacctgaggg ctt

23

<210> 109
<211> 19
<212> DNA
<213> Homo sapiens

<400> 109
ttgagatggg aggcagctt

19

<210> 110
<211> 17
<212> DNA
<213> Homo sapiens

<400> 110
ttcggctccc aggacat

17

<210> 111
<211> 20
<212> DNA
<213> Homo sapiens

<400> 111
cacagagctt tgcaggtgaa

20

PCT/EP2004/004076

